

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-Chloro-2-methyl-3-nitrobenzene

Matthew A. Pearce and Joseph M. Tanski*

Department of Chemistry, Vassar College, Poughkeepsie, NY 12604, USA

Correspondence e-mail: jotanski@vassar.edu


Received 27 January 2011; accepted 6 February 2011

Key indicators: single-crystal X-ray study; $T = 125$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.031; wR factor = 0.090; data-to-parameter ratio = 22.5.

In the title compound, $\text{C}_7\text{H}_6\text{ClNO}_2$, the chloro, methyl and nitro substituents are situated next to each other in this order on the benzene ring, with the mean plane of the nitro group twisted away from the mean plane of the benzene ring by 38.81 (5)°.

Related literature

For information on industrial chemicals, see: Chloronitrotoluenes (2010). For the use of the title compound as a starting material in the synthesis of 7-chlorovasicine (pyrrolo[2,1-*b*]quinazolin-3-ol, 8-chloro-1,2,3,9-tetrahydro), see: Southwick & Cremer (1959). For the toxic effects of the title compound on *D. magna*, see: Ramos *et al.* (2001) and on *T. pyriformis*, see: Schultz (1999); Katritzky *et al.* (2003). For a related structure, see: Liu & Du (2008).



Experimental

Crystal data

$\text{C}_7\text{H}_6\text{ClNO}_2$	$V = 1484.29$ (17) Å ³
$M_r = 171.58$	$Z = 8$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
$a = 7.3061$ (5) Å	$\mu = 0.46$ mm ⁻¹
$b = 13.8392$ (9) Å	$T = 125$ K
$c = 14.6799$ (10) Å	$0.32 \times 0.20 \times 0.10$ mm

Data collection

Bruker APEXII CCD diffractometer	22310 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2007)	2271 independent reflections
$T_{\min} = 0.868$, $T_{\max} = 0.956$	1963 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	101 parameters
$wR(F^2) = 0.090$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\max} = 0.35$ e Å ⁻³
2271 reflections	$\Delta\rho_{\min} = -0.38$ e Å ⁻³

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by Vassar College. X-ray facilities were provided by the US National Science Foundation (grant No. 0521237 to JMT).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2075).

References

- Bruker (2007). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chloronitrotoluenes (2010). *Environment Agency*. Environment Agency Wales, UK.
- Katritzky, A. R., Oliferenko, P., Oliferenko, A., Lomaka, A. & Karelson, M. (2003). *J. Phys. Org. Chem.* **16**, 811–817.
- Liu, H.-L. & Du, Z.-Q. (2008). *Acta Cryst.* **E64**, o536.
- Ramos, E. U., Vaal, M. A. & Hermens, J. L. M. (2001). *Environ. Toxicol. Pharmacol.* **11**, 149–158.
- Schultz, T. W. (1999). *Chem. Res. Toxicol.* **12**, 1262–1267.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Southwick, P. L. & Cremer, S. E. (1959). *J. Org. Chem.* **24**, 753–755.

supporting information

Acta Cryst. (2011). E67, o608 [doi:10.1107/S1600536811004466]

1-Chloro-2-methyl-3-nitrobenzene

Matthew A. Pearce and Joseph M. Tanski

S1. Comment

The title compound, C₇H₆ClNO₂, also known as 2-chloro-6-nitrotoluene, is a common intermediate in the synthesis of industrial chemicals (Chloronitrotoluenes, 2010), as well as in pharmaceuticals such as the bronchodilatory compound vasicine (Southwick *et al.*, 1959). 1-chloro-2-methyl-3-nitrobenzene is relatively toxic to biological species such as the freshwater flea *D. magna* and freshwater protozoa *T. pyriformis*, which suggests that this compound could become a harmful pollutant (Katritzky *et al.*, 2003; Schultz, 1999; Ramos *et al.*, 2001; Chloronitrotoluenes, 2010).

C₇H₆ClNO₂, (I), contains an aromatic ring with chloro, methyl and nitro substituents arranged in this order next to one another. The C—Cl and C—N bond lengths and angles in (I) are very close to those found in a similar structure (Liu & Du, 2008). The central methyl group interacts sterically with the neighboring chloro and nitro groups, as evidenced by the N—C3—C4 and Cl—C1—C6 angles of 115.0 (1)° and 116.78 (8)°, respectively. These angles are compressed from the ideal *sp*²-hybridized carbon atom. The mean plane of the nitro group is twisted away from the mean plane of the aromatic ring by 38.81 (5)°.

S2. Experimental

Crystalline 1-chloro-2-methyl-3-nitrobenzene was purchased from Aldrich Chemical Company, USA.

S3. Refinement

All non-hydrogen atoms were refined anisotropically. Hydrogen atoms on carbon were included in calculated positions and refined using a riding model at C—H = 0.95 or 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5 \times U_{\text{eq}}(\text{C})$ of the aryl and methyl C-atoms, respectively. The extinction parameter (EXTI) refined to zero and was removed from the refinement.




Figure 1

A view of the title compound, with displacement ellipsoids shown at the 50% probability level.

1-Chloro-2-methyl-3-nitrobenzene

Crystal data

$C_7H_6ClNO_2$

$M_r = 171.58$

Orthorhombic, *Pbca*

Hall symbol: $-P\ 2ac\ 2ab$

$a = 7.3061\ (5)\ \text{\AA}$

$b = 13.8392\ (9)\ \text{\AA}$

$c = 14.6799\ (10)\ \text{\AA}$

$V = 1484.29\ (17)\ \text{\AA}^3$

$Z = 8$

$F(000) = 704$

$D_x = 1.536\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9979 reflections

$\theta = 2.8\text{--}30.5^\circ$

$\mu = 0.46\ \text{mm}^{-1}$

$T = 125\ \text{K}$

Block, colorless

$0.32 \times 0.20 \times 0.10\ \text{mm}$

Data collection

Bruker APEXII CCD diffractometer	22310 measured reflections
Radiation source: fine-focus sealed tube	2271 independent reflections
Graphite monochromator	1963 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.038$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$\theta_{\text{max}} = 30.5^\circ$, $\theta_{\text{min}} = 2.8^\circ$
$T_{\text{min}} = 0.868$, $T_{\text{max}} = 0.956$	$h = -10 \rightarrow 10$
	$k = -19 \rightarrow 19$
	$l = -20 \rightarrow 20$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.031$	H-atom parameters constrained
$wR(F^2) = 0.090$	$w = 1/[\sigma^2(F_o^2) + (0.0501P)^2 + 0.4184P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
2271 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
101 parameters	$\Delta\rho_{\text{max}} = 0.35 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.38 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl	0.02248 (4)	0.79411 (2)	0.19541 (2)	0.02921 (10)
O1	0.18412 (16)	0.46030 (7)	0.46513 (7)	0.0383 (2)
O2	0.27741 (17)	0.60583 (8)	0.49519 (6)	0.0414 (3)
N	0.21538 (15)	0.54450 (7)	0.44328 (7)	0.0269 (2)
C1	0.09994 (14)	0.68212 (7)	0.23325 (8)	0.0200 (2)
C2	0.11332 (14)	0.66404 (7)	0.32692 (7)	0.0202 (2)
C3	0.18229 (14)	0.57217 (8)	0.34792 (7)	0.0198 (2)
C4	0.22965 (15)	0.50274 (8)	0.28388 (8)	0.0207 (2)
H4A	0.2734	0.4412	0.3026	0.025*
C5	0.21212 (15)	0.52466 (8)	0.19236 (7)	0.0218 (2)
H5A	0.2439	0.4783	0.1474	0.026*
C6	0.14754 (15)	0.61522 (8)	0.16681 (7)	0.0218 (2)
H6A	0.1360	0.6313	0.1041	0.026*
C7	0.05257 (18)	0.73682 (9)	0.39679 (9)	0.0294 (3)
H7A	-0.0502	0.7744	0.3725	0.044*
H7B	0.1544	0.7804	0.4111	0.044*

H7C 0.0139 0.7032 0.4523 0.044*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl	0.02669 (16)	0.01774 (15)	0.04320 (19)	-0.00006 (9)	-0.00430 (11)	0.00518 (11)
O1	0.0576 (6)	0.0269 (5)	0.0303 (5)	-0.0020 (4)	0.0027 (4)	0.0077 (4)
O2	0.0613 (7)	0.0399 (6)	0.0232 (4)	-0.0141 (5)	-0.0070 (4)	-0.0023 (4)
N	0.0321 (5)	0.0269 (5)	0.0218 (4)	-0.0029 (4)	0.0012 (4)	0.0015 (4)
C1	0.0176 (4)	0.0150 (4)	0.0274 (5)	-0.0011 (4)	-0.0013 (4)	0.0010 (4)
C2	0.0185 (5)	0.0174 (5)	0.0248 (5)	-0.0030 (4)	0.0024 (4)	-0.0032 (4)
C3	0.0208 (5)	0.0197 (5)	0.0189 (5)	-0.0033 (4)	0.0009 (4)	0.0007 (4)
C4	0.0207 (5)	0.0159 (4)	0.0256 (5)	-0.0006 (4)	0.0003 (4)	-0.0015 (4)
C5	0.0221 (5)	0.0202 (5)	0.0232 (5)	-0.0012 (4)	0.0013 (4)	-0.0054 (4)
C6	0.0222 (5)	0.0226 (5)	0.0207 (5)	-0.0027 (4)	-0.0013 (4)	-0.0003 (4)
C7	0.0326 (6)	0.0234 (5)	0.0324 (6)	-0.0011 (5)	0.0074 (5)	-0.0095 (5)

Geometric parameters (Å, °)

Cl—C1	1.7410 (11)	C4—C5	1.3832 (16)
O1—N	1.2300 (13)	C4—H4A	0.9500
O2—N	1.2275 (14)	C5—C6	1.3907 (16)
N—C3	1.4713 (14)	C5—H5A	0.9500
C1—C6	1.3891 (15)	C6—H6A	0.9500
C1—C2	1.4010 (15)	C7—H7A	0.9800
C2—C3	1.4018 (15)	C7—H7B	0.9800
C2—C7	1.5045 (15)	C7—H7C	0.9800
C3—C4	1.3882 (15)		
O2—N—O1	124.17 (11)	C3—C4—H4A	120.6
O2—N—C3	118.13 (10)	C4—C5—C6	119.41 (10)
O1—N—C3	117.66 (10)	C4—C5—H5A	120.3
C6—C1—C2	123.55 (10)	C6—C5—H5A	120.3
C6—C1—Cl	116.78 (8)	C1—C6—C5	119.74 (10)
C2—C1—Cl	119.66 (8)	C1—C6—H6A	120.1
C1—C2—C3	113.76 (10)	C5—C6—H6A	120.1
C1—C2—C7	121.93 (10)	C2—C7—H7A	109.5
C3—C2—C7	124.29 (10)	C2—C7—H7B	109.5
C4—C3—C2	124.64 (10)	H7A—C7—H7B	109.5
C4—C3—N	115.04 (10)	C2—C7—H7C	109.5
C2—C3—N	120.29 (10)	H7A—C7—H7C	109.5
C5—C4—C3	118.87 (10)	H7B—C7—H7C	109.5
C5—C4—H4A	120.6		
C6—C1—C2—C3	0.94 (15)	O1—N—C3—C4	-38.35 (15)
Cl—C1—C2—C3	-178.30 (7)	O2—N—C3—C2	-38.54 (16)
C6—C1—C2—C7	-177.26 (11)	O1—N—C3—C2	143.61 (12)
Cl—C1—C2—C7	3.50 (15)	C2—C3—C4—C5	1.31 (17)

supporting information

C1—C2—C3—C4	-1.68 (15)	N—C3—C4—C5	-176.63 (10)
C7—C2—C3—C4	176.46 (11)	C3—C4—C5—C6	-0.09 (17)
C1—C2—C3—N	176.17 (9)	C2—C1—C6—C5	0.13 (17)
C7—C2—C3—N	-5.69 (16)	Cl—C1—C6—C5	179.39 (8)
O2—N—C3—C4	139.50 (12)	C4—C5—C6—C1	-0.59 (17)
