

catena-Poly[[dimethylbis(thiocyanato- κN)tin(IV)]- μ -(4,4'-bipyridine- $\kappa^2 N:N'$)]

Ezzatollah Najafi,^a Mostafa M. Amini^a and Seik Weng Ng^{b*}

^aDepartment of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 5 February 2011; accepted 14 February 2011

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.023; wR factor = 0.061; data-to-parameter ratio = 16.7.

The title dimethyltin diisothiocyanate adduct of 4,4'-bipyridine, $[\text{Sn}(\text{CH}_3)_2(\text{NCS})_2(\text{C}_{10}\text{H}_8\text{N}_2)]_n$, adopts a chain motif in which the N -heterocycle functions as a bridge to adjacent all-*trans* octahedrally coordinated tin atoms. The Sn^{IV} atom lies on a special position of $2/m$ site symmetry, the methyl C atom on a special position of 2 site symmetry, and the thiocyanate and 4,4'-bipyridine on a special position of m site symmetry.

Related literature

For the 4,4'-bipyridine adducts of diorganotin dichlorides, see: Ma *et al.* (2004); Ng (1998). For the dimethyltin di(isothiocyanate) adduct of 1,10-phenanthroline, see: Najafi *et al.* (2011).



Experimental

Crystal data

$[\text{Sn}(\text{CH}_3)_2(\text{NCS})_2(\text{C}_{10}\text{H}_8\text{N}_2)]$
 $M_r = 421.10$
Monoclinic, $C2/m$
 $a = 10.8697 (8)\text{ \AA}$
 $b = 7.7741 (6)\text{ \AA}$
 $c = 11.3979 (8)\text{ \AA}$
 $\beta = 115.817 (1)^\circ$

$V = 867.0 (1)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.71\text{ mm}^{-1}$
 $T = 295\text{ K}$
 $0.30 \times 0.20 \times 0.10\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.628$, $T_{\max} = 0.847$

4033 measured reflections
1066 independent reflections
1064 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.061$
 $S = 1.08$
1066 reflections

64 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.49\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.82\text{ e \AA}^{-3}$

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2334).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ma, C.-L., Zhang, J.-H. & Zhang, R.-F. (2004). *Heteroatom. Chem.* **15**, 338–346.
- Najafi, E., Amini, M. M. & Ng, S. W. (2011). *Acta Cryst. E67*. Submitted (si2333).
- Ng, S. W. (1998). *Acta Cryst. C54*, 1393–1395.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, m350 [doi:10.1107/S1600536811005459]

catena-Poly[[dimethylbis(thiocyanato- κN)tin(IV)]- μ -(4,4'-bipyridine- $\kappa^2 N:N'$)]

Ezzatollah Najafi, Mostafa M. Amini and Seik Weng Ng

S1. Comment


The 4,4'-bipyridine ligand forms a number of adducts with diorganotin dihalides; the adducts adopt linear chain structures as the ligand functions in a bridging mode. The organotin dihalides include dimethyltin dichloride (Ng, 1998), dibutyltin dichloride and dibenzyltin dichloride (Ma *et al.*, 2004); no pseudohalides have been reported. The dimethyltin diisothiocyanate adduct similarly adopts a chain motif (Scheme I, Fig. 1). Polymeric $[Sn(NCS)_2(CH_3)_2(C_{10}H_8N_2)_2]_n$ has the *N*-heterocycle functioning as a bridging to adjacent all-*trans* octahedrally coordinated tin atoms. The tin atom lies on a special position of $2/m$ site symmetry, the methyl carbon on a special position of 2 site symmetry, and the isothiocyanate and 4,4'-bipyridine on a special position of m site symmetry. The geometry of the tin atom in the dimethyltin di(isothiocyanate) adduct with 1,10-phenanthroline is a *cis*-octahedron (Najafi *et al.*, 2011).

S2. Experimental

Dimethyltin diisothiocyanate (1 mmol, 0.26 g) and 4,4'-bipyridine (1 mmol, 0.16 g) were loaded into a convection tube. The tube was filled with acetonitrile and methanol (v:v / 9:1) and kept at 333 K. Colorless crystals were collected from the side arm after several days.

S3. Refinement

Hydrogen atoms were placed in calculated positions (C–H 0.93–0.96 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of a portion of the $\text{Sn}(\text{NCS})_2(\text{CH}_3)_2(\text{C}_{10}\text{H}_8\text{N}_2)$ chain at the 50% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

catena-Poly[[dimethylbis(thiocyanato- κN)tin(IV)]- μ -(4,4'-bipyridine- $\kappa^2\text{N:N'}$)]

Crystal data

$[\text{Sn}(\text{CH}_3)_2(\text{NCS})_2(\text{C}_{10}\text{H}_8\text{N}_2)]$

$M_r = 421.10$

Monoclinic, $C2/m$

Hall symbol: -C 2y

$a = 10.8697(8)$ Å

$b = 7.7741(6)$ Å

$c = 11.3979(8)$ Å

$\beta = 115.817(1)^\circ$

$V = 867.0(1)$ Å³

$Z = 2$

Data collection

Bruker SMART APEX

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.628$, $T_{\max} = 0.847$

$F(000) = 416$

$D_x = 1.613 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3765 reflections

$\theta = 3.8\text{--}28.3^\circ$

$\mu = 1.71 \text{ mm}^{-1}$

$T = 295$ K

Prism, colorless

$0.30 \times 0.20 \times 0.10$ mm

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.023$

$wR(F^2) = 0.061$

$S = 1.08$

1066 reflections

64 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.048P)^2 + 0.1158P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.49 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.82 \text{ e \AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Sn1	0.5000	0.5000	0.5000	0.03309 (11)	
S1	0.6834 (2)	0.5000	0.18029 (16)	0.0967 (5)	
N1	0.3011 (3)	0.5000	0.2981 (2)	0.0404 (5)	
N2	0.6379 (4)	0.5000	0.3960 (4)	0.0763 (12)	
C1	0.5000	0.2290 (5)	0.5000	0.0713 (12)	
H1A	0.5254	0.1878	0.5869	0.107*	0.50
H1B	0.5644	0.1878	0.4697	0.107*	0.50
H1C	0.4103	0.1878	0.4434	0.107*	0.50
C2	0.1780 (4)	0.5000	0.2946 (3)	0.0713 (14)	
H2	0.1717	0.5000	0.3734	0.086*	
C3	0.0584 (4)	0.5000	0.1812 (3)	0.0721 (15)	
H3	-0.0254	0.5000	0.1849	0.087*	
C4	0.0628 (3)	0.5000	0.0623 (3)	0.0388 (6)	
C5	0.1893 (4)	0.5000	0.0663 (4)	0.102 (3)	
H5	0.1985	0.5000	-0.0112	0.122*	
C6	0.3047 (4)	0.5000	0.1840 (4)	0.099 (2)	
H6	0.3896	0.5000	0.1826	0.119*	
C7	0.6565 (3)	0.5000	0.3071 (4)	0.0501 (8)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.02266 (15)	0.04968 (17)	0.02072 (15)	0.000	0.00366 (10)	0.000
S1	0.1031 (11)	0.1490 (14)	0.0594 (8)	0.000	0.0554 (8)	0.000
N1	0.0246 (11)	0.0643 (15)	0.0225 (11)	0.000	0.0011 (9)	0.000
N2	0.0413 (17)	0.146 (4)	0.0449 (19)	0.000	0.0216 (16)	0.000
C1	0.064 (3)	0.0532 (19)	0.068 (3)	0.000	0.002 (2)	0.000
C2	0.0298 (16)	0.154 (5)	0.0201 (15)	0.000	0.0018 (13)	0.000
C3	0.0252 (16)	0.157 (5)	0.0280 (17)	0.000	0.0057 (14)	0.000
C4	0.0231 (13)	0.0624 (16)	0.0215 (13)	0.000	0.0009 (12)	0.000
C5	0.0260 (17)	0.253 (8)	0.0205 (17)	0.000	0.0053 (14)	0.000
C6	0.0210 (16)	0.244 (8)	0.0231 (17)	0.000	0.0016 (14)	0.000
C7	0.0300 (15)	0.077 (2)	0.0387 (17)	0.000	0.0110 (13)	0.000

Geometric parameters (\AA , $^\circ$)

Sn1—C1 ⁱ	2.107 (4)	C1—H1B	0.9600
Sn1—C1	2.107 (4)	C1—H1C	0.9600
Sn1—N2 ⁱ	2.280 (3)	C2—C3	1.378 (5)
Sn1—N2	2.280 (3)	C2—H2	0.9300

Sn1—N1 ⁱ	2.374 (2)	C3—C4	1.376 (4)
Sn1—N1	2.374 (2)	C3—H3	0.9300
S1—C7	1.595 (4)	C4—C5	1.356 (5)
N1—C6	1.318 (5)	C4—C4 ⁱⁱ	1.480 (5)
N1—C2	1.321 (5)	C5—C6	1.381 (5)
N2—C7	1.116 (5)	C5—H5	0.9300
C1—H1A	0.9600	C6—H6	0.9300
C1 ⁱ —Sn1—C1	180.0	H1A—C1—H1B	109.5
C1 ⁱ —Sn1—N2 ⁱ	90.0	Sn1—C1—H1C	109.5
C1—Sn1—N2 ⁱ	90.000 (1)	H1A—C1—H1C	109.5
C1 ⁱ —Sn1—N2	90.000 (1)	H1B—C1—H1C	109.5
C1—Sn1—N2	90.0	N1—C2—C3	123.9 (3)
N2 ⁱ —Sn1—N2	180.000 (1)	N1—C2—H2	118.1
C1 ⁱ —Sn1—N1 ⁱ	90.000 (1)	C3—C2—H2	118.1
C1—Sn1—N1 ⁱ	90.000 (1)	C4—C3—C2	120.0 (3)
N2 ⁱ —Sn1—N1 ⁱ	91.34 (12)	C4—C3—H3	120.0
N2—Sn1—N1 ⁱ	88.66 (12)	C2—C3—H3	120.0
C1 ⁱ —Sn1—N1	90.0	C5—C4—C3	115.9 (3)
C1—Sn1—N1	90.000 (1)	C5—C4—C4 ⁱⁱ	122.0 (3)
N2 ⁱ —Sn1—N1	88.66 (12)	C3—C4—C4 ⁱⁱ	122.1 (4)
N2—Sn1—N1	91.34 (12)	C4—C5—C6	120.7 (3)
N1 ⁱ —Sn1—N1	180.0	C4—C5—H5	119.6
C6—N1—C2	115.8 (3)	C6—C5—H5	119.6
C6—N1—Sn1	123.4 (2)	N1—C6—C5	123.6 (3)
C2—N1—Sn1	120.8 (2)	N1—C6—H6	118.2
C7—N2—Sn1	153.1 (3)	C5—C6—H6	118.2
Sn1—C1—H1A	109.5	N2—C7—S1	179.8 (4)
Sn1—C1—H1B	109.5		
C1 ⁱ —Sn1—N1—C6	90.0	N1—Sn1—N2—C7	0.000 (2)
C1—Sn1—N1—C6	-90.0	C6—N1—C2—C3	0.0
N2 ⁱ —Sn1—N1—C6	180.0	Sn1—N1—C2—C3	180.0
N2—Sn1—N1—C6	0.0	N1—C2—C3—C4	0.0
C1 ⁱ —Sn1—N1—C2	-90.0	C2—C3—C4—C5	0.0
C1—Sn1—N1—C2	90.0	C2—C3—C4—C4 ⁱⁱ	180.0
N2 ⁱ —Sn1—N1—C2	0.0	C3—C4—C5—C6	0.0
N2—Sn1—N1—C2	180.0	C4 ⁱⁱ —C4—C5—C6	180.0
C1 ⁱ —Sn1—N2—C7	-90.000 (1)	C2—N1—C6—C5	0.0
C1—Sn1—N2—C7	90.000 (1)	Sn1—N1—C6—C5	180.0
N1 ⁱ —Sn1—N2—C7	180.000 (2)	C4—C5—C6—N1	0.0

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y+1, -z$.