

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Acetylacetonato- κ^2O,O')(2-bromo-4-chloro-6-{[2-(dimethylamino)ethyl-imino]methyl}phenolato- κ^3N,N',O)-oxidovanadium(IV)

Fu-Ming Wang

Department of Chemistry, Dezhou University, Dezhou Shandong 253023, People's Republic of China

Correspondence e-mail: wfm99999@126.com


Received 22 April 2011; accepted 24 April 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.009$ Å; R factor = 0.047; wR factor = 0.100; data-to-parameter ratio = 16.8.

The V^{IV} atom in the title complex, $[V(C_{11}H_{13}BrClN_2O)(C_5H_7O_2)O]$, is six-coordinated by one phenolate O, one imino N and one amino N atom of the tridentate anionic Schiff base ligand, by one oxide O atom, and by two O atoms of an acetylacetonate anion, forming a distorted *cis*- VN_2O_4 octahedral coordination geometry. The deviation of the V atom from the plane defined by the three donor atoms of the Schiff base ligand and one O atom of the acetylacetonate ligand towards the oxide O atom is 0.256 (2) Å.

Related literature

For background to oxidovanadium complexes, see: Hiromura *et al.* (2007); Seena *et al.* (2008); Rosenthal *et al.* (2008); Kurup *et al.* (2010). For similar oxidovanadium complexes with Schiff bases, see: Li *et al.* (1988); Cornman *et al.* (1992); Smith *et al.* (2000); Sarkar & Pal (2006).



Experimental

Crystal data

 $[V(C_{11}H_{13}BrClN_2O)(C_5H_7O_2)O]$ $M_r = 470.64$ Orthorhombic, $Pna2_1$ $a = 20.351$ (2) Å $b = 12.749$ (1) Å $c = 7.410$ (2) Å $V = 1922.6$ (6) Å³ $Z = 4$ Mo $K\alpha$ radiation
 $\mu = 2.76$ mm⁻¹ $T = 298$ K
 $0.37 \times 0.33 \times 0.32$ mm

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{min} = 0.429$, $T_{max} = 0.473$ 7060 measured reflections
3863 independent reflections
2284 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.052$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.100$
 $S = 0.93$
3863 reflections
230 parameters
1 restraintH-atom parameters constrained
 $\Delta\rho_{max} = 0.32$ e Å⁻³
 $\Delta\rho_{min} = -0.39$ e Å⁻³
Absolute structure: Flack (1983),
1475 Friedel pairs
Flack parameter: 0.028 (14)

Table 1

Selected geometric parameters (Å, °).

V1—O4	1.598 (4)	V1—N1	2.078 (5)
V1—O1	1.952 (4)	V1—O2	2.168 (4)
V1—O3	1.988 (4)	V1—N2	2.222 (5)
O4—V1—O1	100.0 (2)	O3—V1—O2	82.86 (15)
O4—V1—O3	98.39 (19)	N1—V1—O2	79.70 (17)
O1—V1—O3	88.92 (17)	O4—V1—N2	91.3 (2)
O4—V1—N1	99.13 (19)	O1—V1—N2	165.2 (2)
O1—V1—N1	88.30 (18)	O3—V1—N2	98.79 (17)
O3—V1—N1	162.47 (18)	N1—V1—N2	80.5 (2)
O4—V1—O2	173.0 (2)	O2—V1—N2	81.70 (17)
O1—V1—O2	86.82 (17)		

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported financially by Dezhou University, People's Republic of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5858).

References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cornman, C. R., Kampf, J., Lah, M. S. & Pecoraro, V. L. (1992). *Inorg. Chem.* **31**, 2035–2043.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Hiromura, M., Nakayama, A., Adachi, Y., Doi, M. & Sakurai, H. (2007). *J. Biol. Inorg. Chem.* **12**, 1275–1287.
- Kurup, M. R. P., Seena, E. B. & Kuriakose, M. (2010). *Struct. Chem.* **21**, 599–605.
- Li, X., Lah, M. S. & Pecoraro, V. L. (1988). *Inorg. Chem.* **27**, 4657–4664.
- Rosenthal, E. C. E., Cui, H. L. & Hummert, M. (2008). *Inorg. Chem. Commun.* **11**, 918–920.
- Sarkar, A. & Pal, S. (2006). *Polyhedron*, **25**, 1689–1694.
- Seena, E. B., Mathew, N., Kuriakose, M. & Kurup, M. R. P. (2008). *Polyhedron*, **27**, 1455–1462.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Smith, T. S., Root, C. A., Kampf, J. W., Rasmussen, P. G. & Pecoraro, V. L. (2000). *J. Am. Chem. Soc.* **122**, 767–775.

supporting information

Acta Cryst. (2011). E67, m650 [doi:10.1107/S1600536811015406]

(Acetylacetonato- κ^2O,O')(2-bromo-4-chloro-6-[[2-(dimethylamino)ethylimino]-methyl]phenolato- κ^3N,N',O)oxidovanadium(IV)

Fu-Ming Wang

S1. Comment

Oxovanadium complexes have received much attention due to their structures and biological properties (Hiromura *et al.*, 2007; Seena *et al.*, 2008; Rosenthal *et al.*, 2008; Kurup *et al.*, 2010). In this paper, the title new oxovanadium(IV) complex, (I), with a Schiff base ligand is reported.


The V^{IV} atom in the title complex, Fig. 1, is six-coordinated by one phenolic O, one imino N, and one amino N atoms of the Schiff base ligand, by one oxo O atom, and by two O atoms of an acetylacetonate ligand, forming a distorted octahedral geometry. The deviation of the V atom from the plane defined by the three donor atoms of the Schiff base ligand and one O atom of the acetylacetonate ligand towards the oxo O atom is 0.256 (2) Å. The coordinate bond lengths and angles (Table 1) are comparable with those observed in similar oxovanadium(IV) complexes with Schiff bases and acetylacetonate ligands (Li *et al.*, 1988; Cornman *et al.*, 1992; Smith *et al.*, 2000; Sarkar & Pal, 2006).

S2. Experimental

3-Bromo-5-chlorosalicylaldehyde (1 mmol, 0.23 g), *N,N*-dimethylethane-1,2-diamine (1 mmol, 0.09 g), and VO(acac)₂ (1 mmol, 0.26 g) were mixed in methanol (30 ml). The mixture was boiled under reflux for 2 h, then cooled to room temperature. Green blocks of (I) were formed after slow evaporation of the solution in air for a few days.

S3. Refinement

Hydrogen atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances of 0.93–0.97 Å, and with $U_{\text{iso}}(\text{H})$ set at $1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$.

**Figure 1**

The molecular structure of the title compound, showing displacement ellipsoids drawn at the 30% probability level.

(Acetylacetonato- κ^2O,O')(2-bromo-4-chloro-6-[[2-(dimethylamino)ethylimino]methyl]phenolato- κ^3N,N',O)oxidovanadium(IV)

Crystal data

$[V(C_{11}H_{13}BrClN_2O)(C_5H_7O_2)O]$

$M_r = 470.64$

Orthorhombic, $Pna2_1$

$a = 20.351(2) \text{ \AA}$

$b = 12.749(1) \text{ \AA}$

$c = 7.410(2) \text{ \AA}$

$V = 1922.6(6) \text{ \AA}^3$

$Z = 4$

$F(000) = 948$

$D_x = 1.626 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1129 reflections

$\theta = 2.5\text{--}24.5^\circ$

$\mu = 2.76 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Block, green

$0.37 \times 0.33 \times 0.32 \text{ mm}$

Data collection

Bruker SMART CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.429$, $T_{\max} = 0.473$

7060 measured reflections

3863 independent reflections

2284 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.6^\circ$

$h = -15 \rightarrow 26$

$k = -10 \rightarrow 16$

$l = -9 \rightarrow 9$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.100$
 $S = 0.93$
 3863 reflections
 230 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2)]$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983), 1475 Friedel
 pairs
 Absolute structure parameter: 0.028 (14)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
V1	0.63268 (4)	-0.04041 (7)	0.00361 (16)	0.0381 (3)
Br1	0.64969 (3)	0.21353 (5)	-0.47589 (10)	0.0583 (2)
Cl1	0.51964 (10)	0.50695 (15)	-0.0592 (3)	0.0756 (6)
N1	0.5724 (2)	0.0368 (4)	0.1869 (7)	0.0380 (12)
N2	0.6403 (2)	-0.1505 (4)	0.2353 (7)	0.0439 (14)
O1	0.62782 (18)	0.0834 (3)	-0.1510 (6)	0.0457 (11)
O2	0.70569 (18)	0.0442 (3)	0.1592 (5)	0.0476 (11)
O3	0.71095 (18)	-0.0874 (3)	-0.1350 (5)	0.0431 (10)
O4	0.57889 (17)	-0.1130 (3)	-0.0917 (6)	0.0538 (12)
C1	0.6018 (2)	0.1765 (5)	-0.1231 (8)	0.0343 (14)
C2	0.6071 (3)	0.2535 (5)	-0.2607 (8)	0.0376 (15)
C3	0.5833 (3)	0.3523 (6)	-0.2406 (9)	0.0450 (17)
H3	0.5897	0.4019	-0.3310	0.054*
C4	0.5496 (3)	0.3788 (5)	-0.0854 (10)	0.0501 (17)
C5	0.5398 (2)	0.3065 (5)	0.0468 (9)	0.0426 (15)
H5	0.5161	0.3248	0.1495	0.051*
C6	0.5652 (2)	0.2037 (4)	0.0296 (11)	0.0375 (12)
C7	0.5528 (3)	0.1311 (5)	0.1766 (8)	0.0402 (15)
H7	0.5280	0.1562	0.2728	0.048*
C8	0.5571 (3)	-0.0253 (5)	0.3450 (10)	0.0534 (19)
H8A	0.5201	-0.0709	0.3202	0.064*
H8B	0.5456	0.0203	0.4449	0.064*
C9	0.6162 (3)	-0.0896 (6)	0.3927 (9)	0.060 (2)

H9A	0.6509	-0.0436	0.4353	0.072*
H9B	0.6051	-0.1374	0.4898	0.072*
C10	0.7074 (3)	-0.1882 (6)	0.2724 (10)	0.072 (2)
H10A	0.7069	-0.2319	0.3779	0.108*
H10B	0.7359	-0.1294	0.2923	0.108*
H10C	0.7230	-0.2280	0.1711	0.108*
C11	0.5987 (3)	-0.2457 (5)	0.2077 (13)	0.076 (2)
H11A	0.5551	-0.2247	0.1734	0.113*
H11B	0.5967	-0.2852	0.3179	0.113*
H11C	0.6174	-0.2883	0.1142	0.113*
C12	0.8031 (3)	0.1328 (6)	0.2472 (10)	0.064 (2)
H12A	0.8117	0.0954	0.3572	0.097*
H12B	0.7777	0.1944	0.2733	0.097*
H12C	0.8439	0.1527	0.1923	0.097*
C13	0.7652 (3)	0.0633 (5)	0.1198 (9)	0.0422 (16)
C14	0.7966 (3)	0.0228 (5)	-0.0327 (9)	0.0459 (18)
H14	0.8394	0.0447	-0.0556	0.055*
C15	0.7685 (3)	-0.0473 (5)	-0.1516 (9)	0.0459 (17)
C16	0.8073 (3)	-0.0823 (6)	-0.3127 (10)	0.069 (2)
H16A	0.7848	-0.0620	-0.4209	0.104*
H16B	0.8121	-0.1572	-0.3101	0.104*
H16C	0.8500	-0.0501	-0.3101	0.104*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
V1	0.0346 (4)	0.0410 (6)	0.0388 (6)	0.0030 (4)	0.0017 (5)	-0.0003 (6)
Br1	0.0701 (4)	0.0680 (5)	0.0367 (3)	-0.0054 (3)	0.0078 (4)	-0.0003 (5)
Cl1	0.1159 (14)	0.0445 (11)	0.0665 (12)	0.0281 (11)	0.0116 (11)	0.0049 (10)
N1	0.031 (3)	0.041 (3)	0.042 (3)	-0.001 (2)	0.006 (2)	-0.001 (3)
N2	0.040 (3)	0.043 (4)	0.049 (3)	0.004 (3)	0.010 (2)	0.010 (3)
O1	0.057 (2)	0.043 (3)	0.038 (2)	0.014 (2)	0.018 (2)	0.004 (2)
O2	0.038 (2)	0.061 (3)	0.044 (3)	-0.005 (2)	0.0084 (19)	-0.012 (2)
O3	0.043 (2)	0.043 (3)	0.043 (3)	0.003 (2)	0.004 (2)	-0.006 (2)
O4	0.042 (2)	0.053 (3)	0.066 (3)	-0.002 (2)	-0.012 (2)	-0.005 (2)
C1	0.027 (3)	0.040 (4)	0.036 (4)	-0.002 (3)	-0.005 (2)	-0.001 (3)
C2	0.036 (3)	0.042 (4)	0.035 (4)	0.000 (3)	-0.002 (3)	0.002 (3)
C3	0.052 (4)	0.045 (5)	0.038 (4)	-0.004 (3)	-0.007 (3)	0.005 (3)
C4	0.053 (4)	0.042 (4)	0.056 (5)	0.005 (3)	-0.011 (3)	0.000 (4)
C5	0.047 (3)	0.049 (4)	0.033 (4)	0.004 (3)	-0.005 (3)	-0.003 (4)
C6	0.032 (2)	0.040 (3)	0.041 (3)	0.003 (2)	0.006 (4)	-0.001 (4)
C7	0.032 (3)	0.052 (5)	0.037 (4)	0.003 (3)	0.008 (3)	-0.005 (3)
C8	0.052 (4)	0.045 (4)	0.063 (5)	0.006 (3)	0.027 (3)	0.017 (4)
C9	0.068 (4)	0.062 (5)	0.051 (5)	0.012 (4)	0.020 (4)	0.021 (4)
C10	0.053 (4)	0.076 (6)	0.087 (6)	0.019 (4)	0.008 (4)	0.029 (5)
C11	0.089 (5)	0.054 (5)	0.084 (6)	-0.015 (4)	-0.001 (5)	0.006 (5)
C12	0.055 (4)	0.065 (6)	0.073 (6)	-0.008 (4)	-0.011 (4)	-0.017 (5)
C13	0.037 (4)	0.041 (4)	0.049 (4)	0.000 (3)	0.001 (3)	0.013 (3)

C14	0.032 (3)	0.050 (4)	0.056 (5)	-0.001 (3)	0.009 (3)	0.006 (3)
C15	0.051 (4)	0.045 (5)	0.042 (4)	0.017 (4)	0.007 (3)	0.019 (4)
C16	0.069 (5)	0.090 (6)	0.049 (4)	0.010 (4)	0.030 (4)	-0.001 (5)

Geometric parameters (Å, °)

V1—O4	1.598 (4)	C6—C7	1.452 (9)
V1—O1	1.952 (4)	C7—H7	0.9300
V1—O3	1.988 (4)	C8—C9	1.497 (8)
V1—N1	2.078 (5)	C8—H8A	0.9700
V1—O2	2.168 (4)	C8—H8B	0.9700
V1—N2	2.222 (5)	C9—H9A	0.9700
Br1—C2	1.885 (6)	C9—H9B	0.9700
C11—C4	1.755 (7)	C10—H10A	0.9600
N1—C7	1.268 (7)	C10—H10B	0.9600
N1—C8	1.448 (8)	C10—H10C	0.9600
N2—C10	1.473 (7)	C11—H11A	0.9600
N2—C9	1.484 (8)	C11—H11B	0.9600
N2—C11	1.494 (7)	C11—H11C	0.9600
O1—C1	1.316 (6)	C12—C13	1.507 (8)
O2—C13	1.269 (6)	C12—H12A	0.9600
O3—C15	1.285 (7)	C12—H12B	0.9600
C1—C6	1.399 (9)	C12—H12C	0.9600
C1—C2	1.420 (8)	C13—C14	1.397 (8)
C2—C3	1.358 (8)	C14—C15	1.379 (8)
C3—C4	1.381 (8)	C14—H14	0.9300
C3—H3	0.9300	C15—C16	1.499 (9)
C4—C5	1.360 (8)	C16—H16A	0.9600
C5—C6	1.414 (7)	C16—H16B	0.9600
C5—H5	0.9300	C16—H16C	0.9600
O4—V1—O1	100.0 (2)	C6—C7—H7	116.7
O4—V1—O3	98.39 (19)	N1—C8—C9	108.6 (5)
O1—V1—O3	88.92 (17)	N1—C8—H8A	110.0
O4—V1—N1	99.13 (19)	C9—C8—H8A	110.0
O1—V1—N1	88.30 (18)	N1—C8—H8B	110.0
O3—V1—N1	162.47 (18)	C9—C8—H8B	110.0
O4—V1—O2	173.0 (2)	H8A—C8—H8B	108.4
O1—V1—O2	86.82 (17)	N2—C9—C8	111.4 (6)
O3—V1—O2	82.86 (15)	N2—C9—H9A	109.3
N1—V1—O2	79.70 (17)	C8—C9—H9A	109.3
O4—V1—N2	91.3 (2)	N2—C9—H9B	109.3
O1—V1—N2	165.2 (2)	C8—C9—H9B	109.3
O3—V1—N2	98.79 (17)	H9A—C9—H9B	108.0
N1—V1—N2	80.5 (2)	N2—C10—H10A	109.5
O2—V1—N2	81.70 (17)	N2—C10—H10B	109.5
C7—N1—C8	120.0 (5)	H10A—C10—H10B	109.5
C7—N1—V1	126.5 (4)	N2—C10—H10C	109.5

C8—N1—V1	113.3 (4)	H10A—C10—H10C	109.5
C10—N2—C9	109.3 (5)	H10B—C10—H10C	109.5
C10—N2—C11	106.6 (5)	N2—C11—H11A	109.5
C9—N2—C11	110.2 (5)	N2—C11—H11B	109.5
C10—N2—V1	114.5 (4)	H11A—C11—H11B	109.5
C9—N2—V1	104.7 (3)	N2—C11—H11C	109.5
C11—N2—V1	111.6 (4)	H11A—C11—H11C	109.5
C1—O1—V1	131.1 (4)	H11B—C11—H11C	109.5
C13—O2—V1	128.8 (4)	C13—C12—H12A	109.5
C15—O3—V1	131.3 (4)	C13—C12—H12B	109.5
O1—C1—C6	124.5 (5)	H12A—C12—H12B	109.5
O1—C1—C2	118.7 (5)	C13—C12—H12C	109.5
C6—C1—C2	116.7 (6)	H12A—C12—H12C	109.5
C3—C2—C1	122.4 (6)	H12B—C12—H12C	109.5
C3—C2—Br1	120.5 (5)	O2—C13—C14	123.5 (6)
C1—C2—Br1	117.1 (5)	O2—C13—C12	117.1 (6)
C2—C3—C4	119.7 (6)	C14—C13—C12	119.3 (5)
C2—C3—H3	120.1	C15—C14—C13	124.5 (5)
C4—C3—H3	120.1	C15—C14—H14	117.7
C5—C4—C3	120.5 (6)	C13—C14—H14	117.7
C5—C4—Cl1	120.0 (5)	O3—C15—C14	125.1 (6)
C3—C4—Cl1	119.5 (6)	O3—C15—C16	116.0 (6)
C4—C5—C6	120.6 (6)	C14—C15—C16	118.9 (6)
C4—C5—H5	119.7	C15—C16—H16A	109.5
C6—C5—H5	119.7	C15—C16—H16B	109.5
C1—C6—C5	119.9 (6)	H16A—C16—H16B	109.5
C1—C6—C7	122.7 (5)	C15—C16—H16C	109.5
C5—C6—C7	117.4 (6)	H16A—C16—H16C	109.5
N1—C7—C6	126.5 (6)	H16B—C16—H16C	109.5
N1—C7—H7	116.7		
