

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Poly[$\{\mu_3\text{-}3\text{-}[4\text{-}(1H\text{-imidazol-}1\text{-ylmethyl})\text{-phenyl]prop-2-enoato-}\kappa\text{N}:\eta^2:\kappa\text{O}\}\text{-copper(I)}$]

Benyong Lou

Department of Chemistry and Chemical Engineering, Minjiang University, Fuzhou 350108, People's Republic of China

Correspondence e-mail: lby@mju.edu.cn


Received 19 April 2011; accepted 21 April 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; R factor = 0.042; wR factor = 0.105; data-to-parameter ratio = 12.9.

In the coordination polymer, $[\text{Cu}^1(\text{C}_{13}\text{H}_{11}\text{N}_2\text{O}_2)]_n$, the Cu^1 atom exists in a trigonal-planar geometry that is defined by the $\text{C}=\text{C}$ unit, the imidazole N atom and carboxylate O atoms from three different ozagrel ligands, resulting in the formation of a three-dimensional framework.

Related literature

For background to the design and construction of coordination polymers, see: Kitagawa *et al.* (2004); Zhao *et al.* (2008). For other olefin complexes, see: Kato *et al.* (1997); Wang *et al.* (2005, 2007); Young *et al.* (1998); Zhang *et al.* (2001).



Experimental

Crystal data

$[\text{Cu}(\text{C}_{13}\text{H}_{11}\text{N}_2\text{O}_2)]$
 $M_r = 290.78$
 Trigonal, $P3_1$
 $a = 9.7894(19)\text{ \AA}$
 $c = 10.483(2)\text{ \AA}$
 $V = 870.0(3)\text{ \AA}^3$

$Z = 3$
 Mo $K\alpha$ radiation
 $\mu = 1.88\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.20 \times 0.20 \times 0.20\text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer
 Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2000)
 $T_{\min} = 0.765$, $T_{\max} = 1.000$

6852 measured reflections
 2105 independent reflections
 1904 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.105$
 $S = 1.03$
 2105 reflections
 163 parameters
 1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.36\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32\text{ e \AA}^{-3}$
 Absolute structure: Flack (1983),
 773 Friedel pairs
 Flack parameter: 0.05 (3)

Table 1

Selected geometric parameters (\AA , $^\circ$).

Cu1—N1 ⁱ	1.962 (5)	Cu1—C3	2.030 (5)
Cu1—C2	2.000 (6)	C2—C3	1.381 (7)
Cu1—O2 ⁱⁱ	2.007 (4)		
N1 ⁱ —Cu1—C2	151.2 (2)	N1 ⁱ —Cu1—C3	111.1 (2)
N1 ⁱ —Cu1—O2 ⁱⁱ	104.12 (19)	C2—Cu1—C3	40.1 (2)
C2—Cu1—O2 ⁱⁱ	104.49 (19)		

Symmetry codes: (i) $-x + y, -x, z + \frac{2}{3}$; (ii) $-y + 1, x - y, z + \frac{1}{3}$.

Data collection: *CrystalClear* (Rigaku, 2000); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The author is grateful for grants from the National Natural Science Foundation of China (grant No. 20901037), Fujian Provincial Department of Education (grant No. JB09181) and the Program for New Century Excellent Talents in Fujian Province University (grant No. JK2010043).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5155).

References

- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Kato, H., Emura, S., Takeuchi, N., Enoki, M., Oogushi, K., Takashima, T., Ohmori, K. & Saito, I. (1997). *J. Int. Med. Res.* **25**, 108–111.
 Kitagawa, S., Kitaura, R. & Noro, S. (2004). *Angew. Chem. Int. Ed.* **43**, 2334–2375.
 Rigaku (2000). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, Y. T., Tang, G. M., Liu, Z. M. & Yi, X. H. (2007). *Cryst. Growth Des.* **7**, 2272–2275.
 Wang, X. S., Zhao, H., Li, Y. H., Xiong, R. G. & You, X. Z. (2005). *Top. Catal.* **35**, 43–61.
 Young, D. M., Geiser, U., Schultz, A. J. & Wang, H. H. (1998). *J. Am. Chem. Soc.* **120**, 1331–1332.
 Zhang, J., Xiong, R. G., Chen, X. T., Che, C. M., Xue, Z. L. & You, X. Z. (2001). *Organometallics*, **20**, 4118–4121.
 Zhao, J., Mi, L., Hu, J., Hou, H. & Fan, Y. (2008). *J. Am. Chem. Soc.* **130**, 15222–15223.

supporting information

Acta Cryst. (2011). E67, m640 [doi:10.1107/S1600536811015005]

Poly[$\{\mu_3$ -3-[4-(1*H*-imidazol-1-ylmethyl)phenyl]prop-2-enoato- $\kappa N:\eta^2:\kappa O$ }copper(I)]

Benyong Lou

S1. Comment

The design and construction of coordination polymers have been an area of explosive growth in recent years (Kitagawa *et al.*, 2004). Some active pharmaceutical ingredients (APIs), which contain carboxylic group, N-containing ring in the structures, have also utilized for constructing specific functional coordination polymers (Zhao *et al.*, 2008). The hydrophilic or hydrophobic groups in drug molecules may play an important role in the structures and properties of final metal-organic frameworks.

Ozagrel, (*E*)-3-[4-(1*H*-imidazol-1-ylmethyl)phenyl]-2-propenic acid, is a selective thromboxane A₂-synthetase inhibitor which is used for treating cerebrovascular disease (Kato *et al.*, 1997). It has a carboxylic group and an imidazole ring in the structure. The molecule is an ideal building block for constructing coordination polymers with specific structures. In this contribution, we report a Cu(I)-olefin coordination polymer of ozagrel, [(C₁₃H₁₁N₂O₂)Cu(I)] (I), which was obtained under solvothermal reaction conditions. In the structure, conjugated olefinic and carboxylic groups of ozagrel link metal centers into a 3-fold helical chain which is linked into a three-dimensional framework structure by metal-imidazole coordination interactions.

Compound I crystallizes in the space group P31 with a deprotonated ozagrel anion and a Cu(I) cation in the asymmetric unit (Fig.1). There exist obvious interactions between Cu(I) center and C=C moiety of the olefin of ozagrel (Cu1—C2, Cu1—C3, Table 1). The C=C bond distance (1.381 Å) of the coordinated olefin is longer than that in free ozagrel (1.324 Å) (Wang *et al.*, 2007). The lengthening of the C=C distance is typical for ethylene that is η^2 -bonded to low-valent, electron-rich, transition metals such as copper(I) (Young *et al.*, 1998). Cu(I) ion is nearly centered in a trigonal planar geometry, which is defined by C=C moiety, imidazole N atom and carboxylic O atom from three different ozagrel molecules. Interestingly, carboxylic group of ozagrel doesn't serve as bidentate moiety as does it in [Cu(3-PYA)]_n reported previously (Zhang *et al.*, 2001). But, conjugated olefinic and carboxylic groups as bidentate spacer link Cu(I) centers into a 3-fold helical chain along *c* axis (Fig.2). Cu(I)-imidazole interactions further link the one-dimensional helical chain into a three-dimensional framework structure (Fig.3). Thus, ozagrel anion acting as a tridentate linker is coordinated to Cu(I) ion generating a three-dimensional coordination polymer based on one-dimensional helical chain of Cu(I) centers.

Since Schultz synthesized the first air-stable Cu(I)-olefin coordination polymer based on fumarate ligand under hydrothermal conditions (Young *et al.*, 1998), some Cu(I)-olefin complexes with extended framework structures have been prepared by crystal engineering strategies (Wang *et al.*, 2005). Impressively, two luminescent two-dimensional layered copper(I)-olefin coordination polymers were constructed by the use of 3(2)-pyridylacrylic acid as tetradentate linkers (Zhang *et al.*, 2001). Therein, acrylic acid anions linked Cu(I) centers into a one-dimensional chain which was further linked into two-dimensional coordination layers by coordinated pyridyl rings. Otherwise from that in pyridylacrylic acid,

the acrylic acid anion in ozagrel acts as a bidentate spacer and links Cu(I) centers into a 3-fold helical chain which is further linked into a three-dimensional framework structure by coordinated imidazole ring. In other words, rigid 3(2)-pyridylacrylic acid resulted in two-dimensional coordination layers by metal coordination to Cu(I) ion while more flexible ozagrel gave rise to a three-dimensional coordination framework. The flexible molecular structure of ozagrel could play the subtle role in the final extended structure.

In conclusion, a Cu(I)-olefin coordination polymer based on ozagrel ligand was synthesized under solvothermal conditions. Conjugated olefinic and carboxylic groups of ozagrel as bidentate spacer link Cu(I) centers into a 3-fold helical chain which is linked into a three-dimensional framework structure by metal-imidazole coordination interactions.

S2. Experimental

Ozagrel (228 mg, 1 mmol) and $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (240 mg, 1 mmol) were suspended in 10 ml methanol and a few drops of triethylamine were added. The mixture was placed in a 23 ml Teflon-lined autoclave, sealed, and placed in a furnace at 130 °C for 2 days. Yellow block crystals were isolated. Element analysis for $\text{C}_{13}\text{H}_{11}\text{N}_2\text{O}_2 \text{Cu}_1$ (%), Calcd: C, 53.65; H, 3.22; N, 9.63; Found: C, 53.57; H, 3.89; N, 9.66.

S3. Refinement

H atoms were located geometrically ($\text{C}-\text{H} = 0.95-1.00 \text{ \AA}$) with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.




Figure 1

ORTEP of compound I with 30% thermal ellipsoids. $A = 1 - Y, X - Y, Z + 1/3$; $B = -X + Y, -X, Z + 2/3$





Figure 2

One-dimensional helical chain of Cu(I) along *c* axis in compound I. Imidazole group of ozagrel is omitted for clarity.

**Figure 3**

The three-dimensional structure of compound I viewed along *c* axis.

Poly[μ_3 -3-[4-(1*H*-imidazol-1-ylmethyl)phenyl]prop-2-enoato- $\kappa N:\eta^2:\kappa O$]copper(I)]

Crystal data

[Cu(C₁₃H₁₁N₂O₂)]

M_r = 290.78

Trigonal, *P*3₁

Hall symbol: P 31

a = 9.7894 (19) Å

c = 10.483 (2) Å

V = 870.0 (3) Å³

Z = 3

F(000) = 444

D_x = 1.665 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 987 reflections

θ = 2.4–27.4°

μ = 1.88 mm⁻¹

T = 293 K

Block, yellow

0.20 × 0.20 × 0.20 mm

Data collection

Rigaku Mercury CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 13.6612 pixels mm⁻¹

φ and ω scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2000)

T_{min} = 0.765, *T_{max}* = 1.000

6852 measured reflections

2105 independent reflections

1904 reflections with *I* > 2σ(*I*)

R_{int} = 0.053

θ_{max} = 27.5°, θ_{min} = 2.4°

h = -12→12

k = -12→12

l = -9→13

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.042

wR(*F*²) = 0.105

S = 1.03

2105 reflections

163 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.052*P*)² + 0.5211*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.36 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.32 \text{ e } \text{Å}^{-3}$

Absolute structure: Flack (1983), 773 Friedel
 pairs
 Absolute structure parameter: 0.05 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.44797 (7)	0.33783 (7)	0.62232 (6)	0.03118 (17)
O1	0.6474 (5)	0.5593 (5)	0.3892 (5)	0.0532 (12)
N1	-0.3953 (6)	-0.0278 (5)	0.1055 (5)	0.0353 (11)
C1	0.6055 (6)	0.4173 (6)	0.3901 (5)	0.0335 (11)
O2	0.6941 (4)	0.3626 (5)	0.3546 (4)	0.0413 (9)
N2	-0.3899 (5)	-0.0886 (6)	0.3067 (5)	0.0375 (10)
C2	0.4448 (6)	0.2963 (6)	0.4352 (5)	0.0312 (11)
H2	0.4069	0.1847	0.4093	0.037*
C3	0.3305 (6)	0.3354 (6)	0.4620 (5)	0.0330 (12)
H3	0.3597	0.4436	0.4320	0.040*
C4	0.1578 (6)	0.2269 (6)	0.4662 (5)	0.0311 (11)
C5	0.0875 (7)	0.0622 (7)	0.4627 (6)	0.0403 (13)
H5	0.1519	0.0150	0.4631	0.048*
C6	-0.0749 (7)	-0.0321 (7)	0.4588 (6)	0.0406 (13)
H6	-0.1213	-0.1436	0.4558	0.049*
C7	0.0604 (6)	0.2919 (7)	0.4678 (6)	0.0387 (13)
H7	0.1061	0.4034	0.4692	0.046*
C8	-0.1017 (7)	0.1980 (7)	0.4675 (6)	0.0417 (14)
H8	-0.1662	0.2449	0.4731	0.050*
C9	-0.1707 (6)	0.0355 (7)	0.4591 (6)	0.0357 (12)
C10	-0.3482 (7)	-0.0695 (9)	0.4438 (6)	0.0441 (15)
H10A	-0.3824	-0.1740	0.4824	0.053*
H10B	-0.4031	-0.0216	0.4884	0.053*
C11	-0.3627 (7)	0.0271 (7)	0.2233 (6)	0.0403 (13)
H11	-0.3250	0.1340	0.2462	0.048*
C12	-0.4460 (7)	-0.2242 (7)	0.2378 (6)	0.0449 (14)
H12	-0.4761	-0.3261	0.2701	0.054*
C13	-0.4504 (7)	-0.1860 (7)	0.1143 (6)	0.0392 (13)
H13	-0.4863	-0.2579	0.0448	0.047*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0309 (3)	0.0364 (4)	0.0284 (3)	0.0185 (3)	0.0010 (3)	0.0030 (3)
O1	0.038 (2)	0.035 (2)	0.082 (4)	0.0155 (18)	0.004 (2)	0.012 (2)
N1	0.041 (3)	0.030 (2)	0.036 (3)	0.019 (2)	-0.003 (2)	0.002 (2)
C1	0.029 (2)	0.043 (3)	0.029 (3)	0.018 (2)	0.002 (2)	0.005 (2)
O2	0.037 (2)	0.044 (2)	0.043 (2)	0.0197 (17)	0.0039 (17)	0.0035 (18)
N2	0.030 (2)	0.044 (3)	0.035 (3)	0.016 (2)	-0.0026 (19)	0.006 (2)
C2	0.030 (3)	0.029 (2)	0.030 (3)	0.011 (2)	0.002 (2)	0.003 (2)
C3	0.031 (3)	0.041 (3)	0.030 (3)	0.021 (2)	-0.005 (2)	0.005 (2)
C4	0.032 (3)	0.033 (3)	0.028 (3)	0.016 (2)	-0.002 (2)	0.005 (2)
C5	0.041 (3)	0.047 (3)	0.043 (3)	0.030 (3)	-0.002 (3)	0.005 (3)
C6	0.036 (3)	0.036 (3)	0.042 (3)	0.012 (2)	-0.001 (2)	0.002 (2)
C7	0.029 (3)	0.031 (3)	0.052 (4)	0.011 (2)	-0.006 (2)	-0.003 (3)
C8	0.036 (3)	0.051 (3)	0.048 (4)	0.029 (3)	-0.007 (3)	0.001 (3)
C9	0.031 (3)	0.041 (3)	0.027 (3)	0.013 (2)	0.000 (2)	0.005 (2)
C10	0.030 (3)	0.054 (4)	0.036 (3)	0.012 (3)	0.001 (2)	0.008 (3)
C11	0.041 (3)	0.037 (3)	0.037 (3)	0.015 (2)	-0.002 (2)	0.004 (2)
C12	0.050 (3)	0.038 (3)	0.048 (4)	0.023 (3)	-0.002 (3)	0.000 (3)
C13	0.047 (3)	0.031 (3)	0.041 (3)	0.021 (3)	-0.003 (3)	-0.003 (2)

Geometric parameters (\AA , $^\circ$)

Cu1—N1 ⁱ	1.962 (5)	C4—C7	1.386 (8)
Cu1—C2	2.000 (6)	C4—C5	1.402 (8)
Cu1—O2 ⁱⁱ	2.007 (4)	C5—C6	1.383 (8)
Cu1—C3	2.030 (5)	C5—H5	0.9500
O1—C1	1.237 (7)	C6—C9	1.392 (8)
N1—C11	1.320 (8)	C6—H6	0.9500
N1—C13	1.364 (7)	C7—C8	1.380 (7)
N1—Cu1 ⁱⁱⁱ	1.962 (5)	C7—H7	0.9500
C1—O2	1.281 (6)	C8—C9	1.386 (8)
C1—C2	1.496 (7)	C8—H8	0.9500
O2—Cu1 ^{iv}	2.007 (4)	C9—C10	1.521 (8)
N2—C11	1.347 (7)	C10—H10A	0.9900
N2—C12	1.363 (8)	C10—H10B	0.9900
N2—C10	1.480 (8)	C11—H11	0.9500
C2—C3	1.381 (7)	C12—C13	1.354 (9)
C2—H2	1.0000	C12—H12	0.9500
C3—C4	1.481 (7)	C13—H13	0.9500
C3—H3	1.0000		
N1 ⁱ —Cu1—C2	151.2 (2)	C6—C5—C4	120.5 (5)
N1 ⁱ —Cu1—O2 ⁱⁱ	104.12 (19)	C6—C5—H5	119.7
C2—Cu1—O2 ⁱⁱ	104.49 (19)	C4—C5—H5	119.7
N1 ⁱ —Cu1—C3	111.1 (2)	C5—C6—C9	120.3 (5)
C2—Cu1—C3	40.1 (2)	C5—C6—H6	119.8

O2 ⁱⁱ —Cu1—C3	144.1 (2)	C9—C6—H6	119.8
C11—N1—C13	106.1 (5)	C8—C7—C4	121.3 (5)
C11—N1—Cu1 ⁱⁱⁱ	122.7 (4)	C8—C7—H7	119.3
C13—N1—Cu1 ⁱⁱⁱ	130.5 (4)	C4—C7—H7	119.3
O1—C1—O2	123.6 (5)	C7—C8—C9	120.2 (5)
O1—C1—C2	121.2 (5)	C7—C8—H8	119.9
O2—C1—C2	115.2 (5)	C9—C8—H8	119.9
C1—O2—Cu1 ^{iv}	104.3 (3)	C8—C9—C6	119.2 (5)
C11—N2—C12	106.8 (5)	C8—C9—C10	121.3 (5)
C11—N2—C10	126.7 (5)	C6—C9—C10	119.5 (5)
C12—N2—C10	126.1 (5)	N2—C10—C9	109.7 (5)
C3—C2—C1	121.5 (5)	N2—C10—H10A	109.7
C3—C2—Cu1	71.1 (3)	C9—C10—H10A	109.7
C1—C2—Cu1	104.2 (3)	N2—C10—H10B	109.7
C3—C2—H2	116.6	C9—C10—H10B	109.7
C1—C2—H2	116.6	H10A—C10—H10B	108.2
Cu1—C2—H2	116.6	N1—C11—N2	111.0 (5)
C2—C3—C4	126.9 (5)	N1—C11—H11	124.5
C2—C3—Cu1	68.8 (3)	N2—C11—H11	124.5
C4—C3—Cu1	114.9 (4)	C13—C12—N2	106.9 (5)
C2—C3—H3	112.8	C13—C12—H12	126.6
C4—C3—H3	112.8	N2—C12—H12	126.6
Cu1—C3—H3	112.8	C12—C13—N1	109.2 (5)
C7—C4—C5	118.2 (5)	C12—C13—H13	125.4
C7—C4—C3	118.2 (5)	N1—C13—H13	125.4
C5—C4—C3	123.5 (5)		

Symmetry codes: (i) $-x+y, -x, z+2/3$; (ii) $-\gamma+1, x-\gamma, z+1/3$; (iii) $-\gamma, x-\gamma, z-2/3$; (iv) $-x+y+1, -x+1, z-1/3$.