

# (1*S*,3*S*)-Methyl 6,7-dimethoxy-1-phenyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylate

Tricia Naicker,<sup>a</sup> Thavendran Govender,<sup>a</sup> Hendrik G. Kruger<sup>b</sup> and Glenn E. M. Maguire<sup>b\*</sup>

<sup>a</sup>School of Pharmacy and Pharmacology, University of KwaZulu Natal, Durban 4000, South Africa, and <sup>b</sup>School of Chemistry, University of KwaZulu Natal, Durban 4000, South Africa  
Correspondence e-mail: maguireg@ukzn.ac.za


Received 15 May 2011; accepted 17 May 2011

Key indicators: single-crystal X-ray study;  $T = 173\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.030;  $wR$  factor = 0.082; data-to-parameter ratio = 10.2.

In the title compound,  $\text{C}_{19}\text{H}_{21}\text{NO}_4$ , an organocatalyst with a tetrahydroisoquinoline backbone, the heterocyclic ring assumes a half-boat conformation. The dihedral angle between the aromatic rings is  $82.93(8)^\circ$ . In the crystal, molecules are linked via  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds, forming a layer parallel to  $(10\bar{1})$ .

## Related literature

For related structures, see: Naicker *et al.* (2010, 2011).



## Experimental

### Crystal data

$\text{C}_{19}\text{H}_{21}\text{NO}_4$	$V = 843.25(4)\text{ \AA}^3$
$M_r = 327.37$	$Z = 2$
Monoclinic, $P2_1$	$\text{Mo K}\alpha$ radiation
$a = 9.3841(3)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 6.3453(2)\text{ \AA}$	$T = 173\text{ K}$
$c = 14.2048(4)\text{ \AA}$	$0.90 \times 0.07 \times 0.06\text{ mm}$
$\beta = 94.475(2)^\circ$	

### Data collection

Nonius KappaCCD diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.923$ ,  $T_{\max} = 0.995$

4184 measured reflections  
2275 independent reflections  
2138 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.010$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.082$   
 $S = 1.05$   
2275 reflections  
222 parameters  
2 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.13\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N $\cdots$ O3 <sup>i</sup>	0.91 (2)	2.27 (1)	3.0918 (17)	149 (2)
C1—H1 $\cdots$ O3 <sup>ii</sup>	1.00	2.55	3.503 (2)	160
C19—H19B $\cdots$ O2 <sup>iii</sup>	0.98	2.53	3.270 (2)	132

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z + 2$ ; (ii)  $x, y + 1, z$ ; (iii)  $-x, y - \frac{1}{2}, -z + 1$ .

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *DENZO-SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *SHELXL97*.

The authors wish to thank Dr Hong Su from the Chemistry Department of the University of Cape Town for her assistance with the crystallographic data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2714).

## References

- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Naicker, T., Govender, T., Kruger, H. G. & Maguire, G. E. M. (2011). *Acta Cryst. E* **67**, o1403.
- Naicker, T., Petzold, K., Singh, T., Arvidsson, P. I., Kruger, H. G., Maguire, G. E. M. & Govender, T. (2010). *Tetrahedron Asymmetry*, **21**, 2859–2867.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2011). E67, o1501 [doi:10.1107/S1600536811018782]

## (*1S,3S*)-Methyl 6,7-dimethoxy-1-phenyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylate

Tricia Naicker, Thavendran Govender, Hendrik G. Kruger and Glenn E. M. Maguire

### S1. Comment

The title compound is a novel chiral organocatalyst containing a tetrahydroisoquinoline (TIQ) framework. We have recently reported the use of similar TIQ derivatives as organocatalysts in the Diels-Alder cycloaddition between alpha, beta-unsaturated aldehydes and cyclopentadiene (Naicker *et al.*, 2010).

Diastereomers formed during the synthesis of the title compound were easily separated using column chromatography to yield the TIQ derivative with the stereochemistry as illustrated in Fig. 1. The absolute stereochemistry was confirmed to be *S,S* at the C1 and C9 positions, respectively, by proton NMR spectroscopy.

The *N*-containing six-membered ring assumes a half-boat conformation [ $Q = 0.5537$  (16) Å,  $\theta = 53.94$  (16)° and  $\varphi = 335.3$  (2)°]. This observation is similar to a related structure that we recently reported (Naicker *et al.*, 2011). The molecules are linked through N1—H1N···O3<sup>i</sup> and C1—H1···O3<sup>ii</sup> hydrogen bonds (Table 1) into a column stacked along the *b* axis. The columns are further connected by C19—H19B···O2<sup>iii</sup> hydrogen bonds, forming a layer parallel to the (10<sup>−1</sup>) plane (Fig. 2).

### S2. Experimental


To a stirred solution of 1:1 methanol: methylene chloride (6.0 ml) with 4 Å molecular sieves, (*S*)-methyl 2-amino-3-(3,4-dimethoxyphenyl)propanoate (1.0 g, 4.2 mmol) and benzaldehyde (1.1 eq.) was added under an inert atmosphere. The reaction mixture was allowed to stir for 1.5 h. Thereafter the reaction mixture was filtered and the solvents was removed *in vacuo* to yield the intermediate imine which was left on a high vacuum pump to remove any residual water for 2 h. The residue was then dissolved in trifluoroacetic acid (20 ml) and refluxed for 3 h. The reaction mixture was then neutralized with a saturated sodium bicarbonate solution and extracted with ethylacetate (4 × 20 ml). The organic extracts were combined and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and the solvent was removed *in vacuo*. The crude product (diastereomers) was purified by column chromatography (50:50 EtOAc/Hexane, *R*<sub>f</sub> 1/2) to afford the product 1.20 g (88%) as a white solid. Melting point 370–372 K. IR (neat): 2928, 2600, 1746, 1516, 1250, 1123, 727 cm<sup>−1</sup> [ $\alpha$ ]<sup>20</sup><sub>D</sub> = +15.38 (*c* 0.26 in CHCl<sub>3</sub>) <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.33 – 7.11 (m, 5H), 6.57 (s, 1H), 6.10 (s, 1H), 5.02 (s, 1H), 3.79 (s, 4H), 3.70 (s, 3H), 3.52 (s, 3H), 3.01 (s, 2H). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 172.96, 147.76, 147.41, 143.87, 130.22, 129.04, 128.59, 127.84, 126.07, 111.31, 110.56, 62.85, 56.54, 55.89, 55.84, 52.18, 32.22.

Recrystallization from ethyl acetate at room temperature afforded crystals suitable for X-ray analysis.

### S3. Refinement


All hydrogen atoms, except H1N on N1, were placed in idealized positions and refined as riding, with  $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5U_{\text{eq}}(\text{C})$ . The position of H1N was located in a difference electron density map and refined with a bond length restraint of N—H = 0.95 (3) Å. With unmerged data, the Flack *x* parameter refines to -0.5475 with e.s.d. 0.6554, and the absolute

structure cannot be determined reliably. The final refinements were performed with merged data.



**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 40% probability level. Hydrogen atoms have been omitted for clarity.

**Figure 2**

A partial projection of the title compound, viewed along the  $a$  axis.

### (1*S*,3*S*)-Methyl 6,7-dimethoxy-1-phenyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylate

#### Crystal data

$C_{19}H_{21}NO_4$   
 $M_r = 327.37$   
Monoclinic,  $P2_1$   
Hall symbol: P 2yb  
 $a = 9.3841 (3) \text{ \AA}$   
 $b = 6.3453 (2) \text{ \AA}$   
 $c = 14.2048 (4) \text{ \AA}$   
 $\beta = 94.475 (2)^\circ$   
 $V = 843.25 (4) \text{ \AA}^3$   
 $Z = 2$

$F(000) = 348$   
 $D_x = 1.289 \text{ Mg m}^{-3}$   
Melting point: 371 K  
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
Cell parameters from 4184 reflections  
 $\theta = 2.2\text{--}28.3^\circ$   
 $\mu = 0.09 \text{ mm}^{-1}$   
 $T = 173 \text{ K}$   
Needle, colourless  
 $0.90 \times 0.07 \times 0.06 \text{ mm}$

*Data collection*

Nonius KappaCCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.923$ ,  $T_{\max} = 0.995$

4184 measured reflections  
2275 independent reflections  
2138 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.010$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.2^\circ$   
 $h = -12 \rightarrow 12$   
 $k = -8 \rightarrow 8$   
 $l = -18 \rightarrow 18$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.082$   
 $S = 1.05$   
2275 reflections  
222 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods  
Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0478P)^2 + 0.1004P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$   
Extinction correction: *SHELXL97* (Sheldrick,  
2008),  $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$   
Extinction coefficient: 0.014 (4)

*Special details*

**Experimental.** Half sphere of data collected using *COLLECT* strategy (Nonius, 2000). Crystal to detector distance = 33 mm; combination of  $\varphi$  and  $\omega$  scans of  $1.0^\circ$ , 60 s per  $^\circ$ , 2 iterations.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.32761 (13)	0.8193 (2)	0.52227 (9)	0.0421 (3)
O2	0.14691 (13)	0.5304 (2)	0.46573 (8)	0.0377 (3)
O3	0.39059 (11)	-0.1210 (2)	0.96117 (8)	0.0372 (3)
O4	0.15273 (11)	-0.0802 (2)	0.93763 (8)	0.0359 (3)
N1	0.44475 (13)	0.2219 (2)	0.85516 (9)	0.0286 (3)
H1N	0.4969 (18)	0.214 (4)	0.9118 (11)	0.033 (5)*
C1	0.45806 (15)	0.4396 (3)	0.82212 (10)	0.0276 (3)
H1	0.4271	0.5382	0.8715	0.033*
C2	0.36095 (14)	0.4684 (2)	0.73185 (10)	0.0260 (3)
C3	0.38575 (16)	0.6398 (3)	0.67243 (11)	0.0303 (3)
H3	0.4558	0.7420	0.6921	0.036*
C4	0.30923 (16)	0.6610 (3)	0.58571 (11)	0.0309 (3)
C5	0.20784 (15)	0.5077 (3)	0.55602 (10)	0.0299 (3)

C6	0.17693 (14)	0.3466 (3)	0.61690 (10)	0.0273 (3)
H6	0.1036	0.2485	0.5984	0.033*
C7	0.25316 (14)	0.3266 (2)	0.70613 (10)	0.0250 (3)
C8	0.21706 (15)	0.1462 (3)	0.76978 (10)	0.0274 (3)
H8A	0.2446	0.0110	0.7415	0.033*
H8B	0.1127	0.1433	0.7759	0.033*
C9	0.29613 (15)	0.1715 (3)	0.86760 (10)	0.0259 (3)
H9	0.2524	0.2909	0.9011	0.031*
C10	0.28876 (15)	-0.0258 (3)	0.92677 (10)	0.0273 (3)
C11	0.1344 (2)	-0.2585 (3)	0.99879 (13)	0.0423 (4)
H11A	0.0322	-0.2859	1.0025	0.063*
H11B	0.1788	-0.2277	1.0621	0.063*
H11C	0.1799	-0.3829	0.9733	0.063*
C12	0.61296 (15)	0.4843 (3)	0.80537 (10)	0.0291 (3)
C13	0.68782 (16)	0.3487 (3)	0.74980 (11)	0.0374 (4)
H13	0.6419	0.2270	0.7229	0.045*
C14	0.82974 (17)	0.3906 (4)	0.73342 (12)	0.0426 (4)
H14	0.8798	0.2980	0.6951	0.051*
C15	0.89763 (17)	0.5663 (4)	0.77280 (13)	0.0431 (4)
H15	0.9951	0.5924	0.7631	0.052*
C16	0.82347 (18)	0.7041 (3)	0.82631 (13)	0.0419 (4)
H16	0.8695	0.8269	0.8520	0.050*
C17	0.68109 (16)	0.6639 (3)	0.84278 (11)	0.0339 (3)
H17	0.6307	0.7594	0.8796	0.041*
C18	0.43336 (19)	0.9743 (3)	0.54802 (15)	0.0450 (4)
H18A	0.4362	1.0782	0.4972	0.067*
H18B	0.5270	0.9062	0.5585	0.067*
H18C	0.4097	1.0450	0.6061	0.067*
C19	0.0553 (2)	0.3649 (3)	0.43094 (12)	0.0480 (5)
H19A	0.0173	0.3981	0.3664	0.072*
H19B	-0.0238	0.3498	0.4715	0.072*
H19C	0.1094	0.2328	0.4308	0.072*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0381 (6)	0.0362 (7)	0.0523 (7)	-0.0028 (6)	0.0043 (5)	0.0163 (6)
O2	0.0426 (6)	0.0405 (7)	0.0296 (5)	-0.0005 (6)	-0.0009 (4)	0.0067 (5)
O3	0.0353 (6)	0.0404 (7)	0.0356 (6)	0.0093 (6)	0.0002 (4)	0.0060 (5)
O4	0.0313 (5)	0.0380 (7)	0.0380 (6)	-0.0025 (5)	-0.0002 (4)	0.0088 (5)
N1	0.0229 (6)	0.0322 (7)	0.0299 (6)	0.0000 (5)	-0.0032 (5)	0.0002 (6)
C1	0.0234 (6)	0.0302 (8)	0.0290 (7)	-0.0003 (6)	0.0007 (5)	-0.0056 (6)
C2	0.0214 (6)	0.0254 (7)	0.0315 (7)	0.0018 (6)	0.0031 (5)	-0.0026 (6)
C3	0.0237 (6)	0.0251 (7)	0.0422 (8)	0.0006 (6)	0.0023 (5)	-0.0008 (7)
C4	0.0263 (6)	0.0281 (8)	0.0390 (8)	0.0036 (6)	0.0077 (6)	0.0067 (7)
C5	0.0266 (7)	0.0337 (8)	0.0296 (7)	0.0040 (6)	0.0030 (5)	0.0018 (6)
C6	0.0237 (6)	0.0286 (7)	0.0296 (7)	0.0000 (6)	0.0012 (5)	-0.0001 (6)
C7	0.0218 (6)	0.0257 (7)	0.0276 (6)	0.0017 (6)	0.0027 (5)	-0.0004 (6)

C8	0.0243 (6)	0.0284 (7)	0.0289 (7)	-0.0023 (6)	-0.0012 (5)	0.0021 (6)
C9	0.0240 (6)	0.0271 (7)	0.0267 (6)	0.0021 (6)	0.0017 (5)	-0.0016 (6)
C10	0.0302 (7)	0.0291 (8)	0.0223 (6)	0.0015 (6)	0.0008 (5)	-0.0040 (6)
C11	0.0477 (9)	0.0388 (10)	0.0404 (9)	-0.0081 (8)	0.0035 (7)	0.0082 (8)
C12	0.0232 (6)	0.0350 (8)	0.0284 (7)	-0.0005 (6)	-0.0017 (5)	-0.0012 (6)
C13	0.0290 (7)	0.0449 (10)	0.0381 (8)	0.0022 (8)	0.0010 (6)	-0.0077 (8)
C14	0.0299 (7)	0.0590 (12)	0.0394 (8)	0.0068 (8)	0.0068 (6)	-0.0014 (9)
C15	0.0247 (7)	0.0602 (12)	0.0445 (9)	-0.0010 (8)	0.0029 (6)	0.0106 (9)
C16	0.0314 (8)	0.0473 (11)	0.0460 (9)	-0.0093 (8)	-0.0031 (7)	0.0041 (9)
C17	0.0286 (7)	0.0370 (9)	0.0357 (8)	-0.0045 (7)	0.0002 (6)	-0.0014 (7)
C18	0.0414 (9)	0.0287 (9)	0.0673 (12)	-0.0008 (8)	0.0203 (8)	0.0072 (9)
C19	0.0671 (12)	0.0411 (11)	0.0331 (8)	0.0001 (10)	-0.0126 (8)	0.0006 (8)

*Geometric parameters (Å, °)*

O1—C4	1.3696 (19)	C8—H8B	0.9900
O1—C18	1.424 (2)	C9—C10	1.512 (2)
O2—C5	1.3704 (18)	C9—H9	1.0000
O2—C19	1.421 (2)	C11—H11A	0.9800
O3—C10	1.2017 (18)	C11—H11B	0.9800
O4—C10	1.3429 (18)	C11—H11C	0.9800
O4—C11	1.445 (2)	C12—C17	1.392 (2)
N1—C9	1.4552 (18)	C12—C13	1.394 (2)
N1—C1	1.467 (2)	C13—C14	1.395 (2)
N1—H1N	0.909 (14)	C13—H13	0.9500
C1—C12	1.5178 (19)	C14—C15	1.380 (3)
C1—C2	1.525 (2)	C14—H14	0.9500
C1—H1	1.0000	C15—C16	1.382 (3)
C2—C7	1.381 (2)	C15—H15	0.9500
C2—C3	1.407 (2)	C16—C17	1.398 (2)
C3—C4	1.383 (2)	C16—H16	0.9500
C3—H3	0.9500	C17—H17	0.9500
C4—C5	1.403 (2)	C18—H18A	0.9800
C5—C6	1.384 (2)	C18—H18B	0.9800
C6—C7	1.4118 (18)	C18—H18C	0.9800
C6—H6	0.9500	C19—H19A	0.9800
C7—C8	1.513 (2)	C19—H19B	0.9800
C8—C9	1.5317 (19)	C19—H19C	0.9800
C8—H8A	0.9900		
C4—O1—C18	117.28 (14)	C8—C9—H9	108.9
C5—O2—C19	116.41 (13)	O3—C10—O4	123.84 (15)
C10—O4—C11	115.40 (13)	O3—C10—C9	124.95 (14)
C9—N1—C1	110.63 (12)	O4—C10—C9	111.20 (12)
C9—N1—H1N	109.5 (12)	O4—C11—H11A	109.5
C1—N1—H1N	106.5 (15)	O4—C11—H11B	109.5
N1—C1—C12	109.42 (13)	H11A—C11—H11B	109.5
N1—C1—C2	108.75 (12)	O4—C11—H11C	109.5

C12—C1—C2	111.26 (12)	H11A—C11—H11C	109.5
N1—C1—H1	109.1	H11B—C11—H11C	109.5
C12—C1—H1	109.1	C17—C12—C13	119.00 (14)
C2—C1—H1	109.1	C17—C12—C1	120.67 (14)
C7—C2—C3	119.79 (13)	C13—C12—C1	120.31 (14)
C7—C2—C1	121.43 (13)	C12—C13—C14	120.43 (18)
C3—C2—C1	118.75 (13)	C12—C13—H13	119.8
C4—C3—C2	120.72 (14)	C14—C13—H13	119.8
C4—C3—H3	119.6	C15—C14—C13	120.18 (18)
C2—C3—H3	119.6	C15—C14—H14	119.9
O1—C4—C3	125.04 (15)	C13—C14—H14	119.9
O1—C4—C5	115.36 (14)	C14—C15—C16	119.84 (16)
C3—C4—C5	119.55 (14)	C14—C15—H15	120.1
O2—C5—C6	124.75 (14)	C16—C15—H15	120.1
O2—C5—C4	115.61 (14)	C15—C16—C17	120.37 (18)
C6—C5—C4	119.64 (13)	C15—C16—H16	119.8
C5—C6—C7	120.76 (14)	C17—C16—H16	119.8
C5—C6—H6	119.6	C12—C17—C16	120.14 (16)
C7—C6—H6	119.6	C12—C17—H17	119.9
C2—C7—C6	119.20 (13)	C16—C17—H17	119.9
C2—C7—C8	121.88 (12)	O1—C18—H18A	109.5
C6—C7—C8	118.89 (13)	O1—C18—H18B	109.5
C7—C8—C9	110.34 (12)	H18A—C18—H18B	109.5
C7—C8—H8A	109.6	O1—C18—H18C	109.5
C9—C8—H8A	109.6	H18A—C18—H18C	109.5
C7—C8—H8B	109.6	H18B—C18—H18C	109.5
C9—C8—H8B	109.6	O2—C19—H19A	109.5
H8A—C8—H8B	108.1	O2—C19—H19B	109.5
N1—C9—C10	109.60 (12)	H19A—C19—H19B	109.5
N1—C9—C8	108.29 (11)	O2—C19—H19C	109.5
C10—C9—C8	112.19 (12)	H19A—C19—H19C	109.5
N1—C9—H9	108.9	H19B—C19—H19C	109.5
C10—C9—H9	108.9		
C9—N1—C1—C12	-177.34 (12)	C5—C6—C7—C8	-179.14 (14)
C9—N1—C1—C2	-55.63 (14)	C2—C7—C8—C9	9.86 (19)
N1—C1—C2—C7	16.42 (18)	C6—C7—C8—C9	-171.91 (13)
C12—C1—C2—C7	137.00 (14)	C1—N1—C9—C10	-163.99 (11)
N1—C1—C2—C3	-161.44 (13)	C1—N1—C9—C8	73.33 (15)
C12—C1—C2—C3	-40.86 (19)	C7—C8—C9—N1	-46.80 (16)
C7—C2—C3—C4	-4.0 (2)	C7—C8—C9—C10	-167.89 (11)
C1—C2—C3—C4	173.89 (14)	C11—O4—C10—O3	3.2 (2)
C18—O1—C4—C3	-0.3 (2)	C11—O4—C10—C9	-175.80 (13)
C18—O1—C4—C5	-177.71 (14)	N1—C9—C10—O3	2.7 (2)
C2—C3—C4—O1	-178.59 (14)	C8—C9—C10—O3	123.00 (16)
C2—C3—C4—C5	-1.2 (2)	N1—C9—C10—O4	-178.38 (12)
C19—O2—C5—C6	-5.8 (2)	C8—C9—C10—O4	-58.05 (15)
C19—O2—C5—C4	173.61 (15)	N1—C1—C12—C17	-130.54 (15)

O1—C4—C5—O2	3.6 (2)	C2—C1—C12—C17	109.27 (16)
C3—C4—C5—O2	-174.04 (14)	N1—C1—C12—C13	51.37 (18)
O1—C4—C5—C6	-177.03 (13)	C2—C1—C12—C13	-68.82 (19)
C3—C4—C5—C6	5.4 (2)	C17—C12—C13—C14	1.2 (3)
O2—C5—C6—C7	175.00 (14)	C1—C12—C13—C14	179.29 (16)
C4—C5—C6—C7	-4.4 (2)	C12—C13—C14—C15	0.5 (3)
C3—C2—C7—C6	5.0 (2)	C13—C14—C15—C16	-1.9 (3)
C1—C2—C7—C6	-172.82 (13)	C14—C15—C16—C17	1.6 (3)
C3—C2—C7—C8	-176.77 (13)	C13—C12—C17—C16	-1.4 (2)
C1—C2—C7—C8	5.4 (2)	C1—C12—C17—C16	-179.55 (15)
C5—C6—C7—C2	-0.9 (2)	C15—C16—C17—C12	0.1 (3)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···O3 <sup>i</sup>	0.91 (2)	2.27 (1)	3.0918 (17)	149 (2)
C1—H1···O3 <sup>ii</sup>	1.00	2.55	3.503 (2)	160
C19—H19B···O2 <sup>iii</sup>	0.98	2.53	3.270 (2)	132

Symmetry codes: (i)  $-x+1, y+1/2, -z+2$ ; (ii)  $x, y+1, z$ ; (iii)  $-x, y-1/2, -z+1$ .