

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Poly[(μ_5 -3,5-dinitrobenzoato)-rubidium]

Yanqing Miao,* Xiaoqing Zhang and Chunye Liu

Xi'an Medical University, Department of Pharmacy, Hanguang Road No.137, Xi'an 710021, Shaanxi, People's Republic of China

Correspondence e-mail: miaoyanqing66@163.com


Received 11 May 2011; accepted 14 June 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.033; wR factor = 0.078; data-to-parameter ratio = 11.9.

The asymmetric unit of the title compound, $[\text{Rb}(\text{C}_7\text{H}_3\text{N}_2\text{O}_6)]_n$, comprises an Rb cation and a 3,5-dinitrobenzoate anion. The Rb cation is eight-coordinated by O atoms from five 3,5-dinitrobenzoate anions. On the other hand, each 3,5-dinitrobenzoate anion links five Rb cations with the carboxylate groups as μ_5 -bridging. The metal atom is firstly linked by the carboxylate groups into a chain along the c -axis direction, which is further linked by bonds between the Rb and nitro O atoms, giving a three-dimensional framework.

Related literature

For 3,5-dinitrobenzoate complexes, see: Askarinejad *et al.* (2007); Madej *et al.* (2007). For Rb—O bond lengths, see: Cametti *et al.* (2005).



Experimental

Crystal data

 $[\text{Rb}(\text{C}_7\text{H}_3\text{N}_2\text{O}_6)]$
 $M_r = 296.58$

 Monoclinic, $I2/a$
 $a = 7.2789$ (15) Å
 $b = 18.072$ (4) Å
 $c = 7.3652$ (14) Å
 $\beta = 91.70$ (3)°

 $V = 968.4$ (3) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 5.13$ mm⁻¹
 $T = 293$ K
 $0.64 \times 0.40 \times 0.14$ mm

Data collection

 Bruker SMART CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.396$, $T_{\max} = 1.000$

 4663 measured reflections
 896 independent reflections
 760 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.078$
 $S = 1.06$
 896 reflections

 75 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29$ e Å⁻³
 $\Delta\rho_{\min} = -0.55$ e Å⁻³

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported financially by grants from the Scientific Research Plan Projects of Shaanxi Province Department of Health (2010D54), the Natural Science Research Plan Projects of Shaanxi Science and Technology Department (SJ08B19) and the Scientific Research Plan Projects of Shaanxi Education Department (09 J K709). Jing You is thanked for helping to improve the English.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2012).

References

- Askarinejad, A., Fadaei, M. R., Morsali, A. & Mahjoub, A. R. (2007). *J. Coord. Chem.* **60**, 753–761.
- Bruker (2002). SMART and SAINT. Bruker AXS Inc, Madison, Wisconsin, USA.
- Cametti, M., Nissinen, M., Cort, A. D., Mandolini, L. & Rissanen, K. (2005). *J. Am. Chem. Soc.* **127**, 3831–3837.
- Madej, A., Oleksyn, B. J. & Śliwiński, J. (2007). *Pol. J. Chem.* **81**, 1201–1218.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, m1002 [doi:10.1107/S1600536811023026]

Poly[(μ_5 -3,5-dinitrobenzoato)rubidium]**Yanqing Miao, Xiaoqing Zhang and Chunye Liu****S1. Comment**

In the coordination chemistry of 3,5-dinitrobenzoic acid, it has been found that the 3,5-dinitrobenzoate moiety functions as a multidentate ligand (Askarinejad *et al.*, 2007; Madej *et al.*, 2007) with versatile binding and coordination modes. In this paper, we report the crystal structure of the title compound, a new Rb complex obtained by the reaction of 3,5-dinitrobenzoic acid and RbOH in water.

The asymmetric unit of the title compound comprises a Rb cation and a 3,5-dinitrobenzoate anion. Rb cation lies on and the dinitrobenzoate is centred upon crystallographic twofold axes. The Rb cation is coordinated to eight O atoms from five 3,5-dinitrobenzoate anions (Fig. 1) with the Rb—O distances ranging from 2.761 (2) Å to 3.124 (4) Å, which are well within the range reported in the literature (Cametti *et al.*, 2005). The Rb centre is firstly linked by the carboxylic groups to give a one-dimensional chain along the *c*-axis direction, which is further linked by the phenyl groups to give a three-dimensional framework of the title compound (Fig. 2).

S2. Experimental

3,5-dinitrobenzoic acid and RbOH were commercially available and used without further purification. To a solution of 10 mmol 3,5-dinitrobenzoic acid in 30 ml bidistilled water, a solution of 10 mmol RbOH in 20 ml bidistilled water was added dropwise at room temperature. After vigorous stirring for 1 h, the resulting solution was then evaporated to a volume of about 15 ml in vacuum and filtered hot. The filtrate was then set aside for crystallization at room temperature. Three weeks later, colorless prism crystals of the title compound suitable for X-ray determination were isolated.

S3. Refinement

Carbon-bound H atoms were placed at calculated positions and were treated as riding on the parent C atoms with C—H = 0.93 Å, and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.




Figure 1

The structure of (I), showing the atomic numbering scheme. Non-H atoms are shown with 30% probability displacement ellipsoids. Symmetry codes: (i) $1.5 - x, y, 1 - z$; (ii) $x, 1.5 - y, -1/2 + z$; (iii) $0.5 - x, 1.5 - y, 1.5 - z$; (iv) $0.5 - x, y, 1 - z$; (v) $-1/2 + x, 1 - y, z$; (vi) $1 - x, 1 - y, 1 - z$.




Figure 2

The three-dimensional framework of (I).

Poly[(μ -3,5-dinitrobenzoato)rubidium]

Crystal data

[Rb(C₇H₃N₂O₆)]

$M_r = 296.58$

Monoclinic, $I2/a$

Hall symbol: $-I\ 2ya$

$a = 7.2789$ (15) Å

$b = 18.072$ (4) Å

$c = 7.3652$ (14) Å

$\beta = 91.70$ (3)°

$V = 968.4$ (3) Å³

$Z = 4$

$F(000) = 576$

$D_x = 2.034$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1610 reflections

$\theta = 3.0$ – 25.4 °

$\mu = 5.13$ mm⁻¹

$T = 293$ K

Prism, colorless

$0.64 \times 0.40 \times 0.14$ mm

Data collection

Bruker SMART CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.396$, $T_{\max} = 1.000$

4663 measured reflections

896 independent reflections

760 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.053$

$\theta_{\max} = 25.3$ °, $\theta_{\min} = 3.0$ °

$h = -8 \rightarrow 8$

$k = -21 \rightarrow 20$

$l = -8 \rightarrow 7$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.078$

$S = 1.06$

896 reflections

75 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0418P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.29$ e Å⁻³

$\Delta\rho_{\min} = -0.55$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Rb1	0.2500	0.54657 (2)	0.5000	0.0557 (2)
C1	0.7500	0.8732 (3)	0.5000	0.0493 (11)
H1A	0.7500	0.9246	0.5000	0.059*
C2	0.6036 (4)	0.83316 (17)	0.5606 (4)	0.0451 (8)

C3	0.5991 (4)	0.75684 (17)	0.5597 (4)	0.0421 (7)
H3A	0.4961	0.7316	0.5986	0.050*
C4	0.7500	0.7185 (2)	0.5000	0.0387 (10)
C5	0.7500	0.6342 (2)	0.5000	0.0398 (10)
N1	0.4460 (5)	0.87295 (16)	0.6344 (4)	0.0595 (8)
O1	0.6019 (3)	0.60328 (12)	0.5293 (4)	0.0649 (7)
O2	0.3212 (4)	0.83742 (15)	0.6978 (4)	0.0696 (7)
O3	0.4500 (5)	0.94073 (15)	0.6317 (5)	0.0913 (11)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Rb1	0.0317 (3)	0.0382 (3)	0.0985 (4)	0.000	0.0213 (2)	0.000
C1	0.066 (3)	0.040 (2)	0.043 (3)	0.000	0.004 (2)	0.000
C2	0.0482 (19)	0.0444 (17)	0.0429 (17)	0.0112 (15)	0.0053 (14)	0.0032 (14)
C3	0.0381 (17)	0.0441 (16)	0.0441 (17)	0.0020 (15)	0.0020 (13)	0.0032 (14)
C4	0.035 (2)	0.042 (2)	0.039 (2)	0.000	-0.0038 (18)	0.000
C5	0.028 (2)	0.035 (2)	0.056 (3)	0.000	-0.0009 (19)	0.000
N1	0.068 (2)	0.0532 (18)	0.0576 (18)	0.0181 (16)	0.0117 (15)	0.0056 (15)
O1	0.0359 (13)	0.0414 (12)	0.118 (2)	-0.0037 (11)	0.0118 (13)	0.0012 (13)
O2	0.0555 (16)	0.0686 (17)	0.0860 (18)	0.0089 (13)	0.0222 (14)	-0.0045 (14)
O3	0.113 (3)	0.0515 (15)	0.112 (2)	0.0326 (16)	0.049 (2)	0.0202 (16)

Geometric parameters (Å, °)

Rb1—O1	2.761 (2)	C1—C2	1.374 (4)
Rb1—O1 ⁱ	2.925 (2)	C1—C2 ⁱⁱⁱ	1.374 (4)
Rb1—O2 ⁱⁱ	3.113 (3)	C1—H1A	0.9300
Rb1—O3 ⁱⁱ	3.125 (3)	C2—C3	1.380 (4)
O1—Rb1—O1 ^{iv}	136.43 (9)	C1—C2—C3	122.9 (3)
O1—Rb1—O1 ⁱ	132.92 (7)	C1—C2—N1	118.9 (3)
O1 ^{iv} —Rb1—O1 ⁱ	90.36 (6)	C3—C2—N1	118.2 (3)
O1 ⁱ —Rb1—O1 ^v	44.43 (9)	C2—C3—C4	118.9 (3)
O1—Rb1—O2 ⁱⁱ	82.42 (8)	C2—C3—H3A	120.5
O1 ^{iv} —Rb1—O2 ⁱⁱ	68.41 (8)	C4—C3—H3A	120.5
O1 ⁱ —Rb1—O2 ⁱⁱ	120.08 (7)	C3 ⁱⁱⁱ —C4—C3	119.9 (4)
O1 ^v —Rb1—O2 ⁱⁱ	138.23 (8)	C3—C4—C5	120.07 (19)
O2 ⁱⁱ —Rb1—O2 ^{vi}	95.32 (11)	O1—C5—O1 ⁱⁱⁱ	126.3 (4)
O1—Rb1—O3 ⁱⁱ	111.10 (10)	O1—C5—C4	116.83 (19)
O1 ^{iv} —Rb1—O3 ⁱⁱ	65.51 (9)	O1 ⁱⁱⁱ —C5—C4	116.83 (19)
O1 ⁱ —Rb1—O3 ⁱⁱ	79.74 (8)	O2—N1—O3	123.5 (3)
O1 ^v —Rb1—O3 ⁱⁱ	108.31 (7)	O2—N1—C2	118.9 (3)
O2 ⁱⁱ —Rb1—O3 ⁱⁱ	40.35 (7)	O3—N1—C2	117.6 (3)
O2 ^{vi} —Rb1—O3 ⁱⁱ	131.54 (7)	C5—O1—Rb1	164.54 (19)
O2 ⁱⁱ —Rb1—O3 ^{vi}	131.54 (8)	C5—O1—Rb1 ^v	94.6 (2)
O3 ⁱⁱ —Rb1—O3 ^{vi}	171.57 (10)	Rb1—O1—Rb1 ^v	89.64 (6)
C2—C1—C2 ⁱⁱⁱ	116.5 (4)	N1—O2—Rb1 ⁱⁱ	93.5 (2)

C2—C1—H1A	121.8	N1—O3—Rb1 ⁱⁱ	92.7 (2)
C3—C4—C5—O1	10.3 (2)	C3—C2—N1—O2	2.7 (5)
C3 ⁱⁱⁱ —C4—C5—O1	-169.7 (2)	C1—C2—N1—O3	2.8 (4)
C1—C2—N1—O2	-175.8 (3)	C3—C2—N1—O3	-178.8 (3)

Symmetry codes: (i) $x-1/2, -y+1, z$; (ii) $-x+1/2, -y+3/2, -z+3/2$; (iii) $-x+3/2, y, -z+1$; (iv) $-x+1/2, y, -z+1$; (v) $-x+1, -y+1, -z+1$; (vi) $x, -y+3/2, z-1/2$.