

Bis(2-amino-6-methyl-1,3-benzothiazole- κ N³)bis(4-nitrobenzoato- κ O¹)zinc

Xue-Tong Sun, Xiu-Guang Wang and Xiao-Jun Zhao*

College of Chemistry, Tianjin Key Laboratory of Structure and Performance for Functional Molecules, Tianjin Normal University, Tianjin 300387, People's Republic of China

Correspondence e-mail: xiaojun_zhao15@yahoo.com.cn


Received 26 May 2011; accepted 9 June 2011

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.036; wR factor = 0.101; data-to-parameter ratio = 12.7.

In the title mononuclear complex, $[\text{Zn}(\text{C}_7\text{H}_4\text{NO}_4)_2(\text{C}_8\text{H}_8\text{N}_2\text{S})_2]$, the Zn^{II} atom is coordinated by two N atoms from two 2-amino-6-methyl-1,3-benzothiazole and by two carboxylate O atoms from two 4-nitrobenzoate ligands, adopting a slightly distorted tetrahedral coordination geometry. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds between the amino group of 2-amino-6-methyl-1,3-benzothiazole and the carboxylate group of 4-nitrobenzoate link these discrete mononuclear units into a one-dimensional supramolecular chain extending parallel to $[100]$.

Related literature

For the properties of metal complexes with aminobenzothiazole and its derivatives, see: Sun & Cui (2008); Chen *et al.* (2008); Kovalska *et al.* (2006); Batista *et al.* (2007); Marconato *et al.* (1998).



Experimental

Crystal data

 $[\text{Zn}(\text{C}_7\text{H}_4\text{NO}_4)_2(\text{C}_8\text{H}_8\text{N}_2\text{S})_2]$
 $M_r = 726.04$

 Monoclinic, $P2_1/c$
 $a = 13.2240$ (8) Å

 $b = 10.7369$ (7) Å
 $c = 21.8863$ (13) Å
 $\beta = 96.099$ (1)°
 $V = 3089.9$ (3) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.99$ mm⁻¹
 $T = 296$ K
 $0.05 \times 0.04 \times 0.03$ mm

Data collection

 Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.952$, $T_{\text{max}} = 0.971$

 15119 measured reflections
 5418 independent reflections
 4724 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.101$
 $S = 1.05$
 5418 reflections

 426 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.42$ e Å⁻³
Table 1

Selected bond lengths (Å).

Zn1—O6	1.9489 (18)	Zn1—N3	2.036 (2)
Zn1—O1	1.9717 (18)	Zn1—N1	2.054 (2)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2A \cdots O1	0.86	2.27	3.008 (3)	144
N2—H2B \cdots O5 ⁱ	0.86	2.05	2.845 (3)	153
N4—H4A \cdots O6	0.86	2.20	2.971 (3)	149
N4—H4B \cdots O2 ⁱⁱ	0.86	2.05	2.860 (3)	156

 Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2003); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008) and DIAMOND (Brandenburg & Berndt, 1999); software used to prepare material for publication: SHELXL97.

The authors gratefully acknowledge financial support from Tianjin Normal University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2014).

References

- Batista, R. M. F., Costa, S. P. G., Malheiro, E. L., Belsley, M. & Raposo, M. M. (2007). *Tetrahedron*, **63**, 4258–4265.
- Brandenburg, K. & Berndt, M. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2001). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2003). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, Q., Yang, E.-C., Zhang, R.-W., Wang, X.-G. & Zhao, X.-J. (2008). *J. Coord. Chem.* **12**, 1951–1962.
- Kovalska, V. B., Volkova, K. D., Losytskyy, M. Y., Tolmachev, O. I., Balanda, A. O. & Yarmoluk, S. M. (2006). *Spectrochim. Acta Part A*, **65**, 271–277.
- Marconato, J. C., Bulhoes, L. O. & Temperini, M. L. (1998). *Electrochim. Acta*, **43**, 771–780.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sun, Y. F. & Cui, Y. P. (2008). *Chin. J. Struct. Chem.* **27**, 1526–1532.

supporting information

Acta Cryst. (2011). E67, m933 [doi:10.1107/S1600536811022331]

Bis(2-amino-6-methyl-1,3-benzothiazole- κ N³)bis(4-nitrobenzoato- κ O¹)zinc

Xue-Tong Sun, Xiu-Guang Wang and Xiao-Jun Zhao

S1. Comment

Organic compounds containing benzothiazole and their metal complexes are of optical, biological, and pharmaceutical importance used extensively as electroluminescent device, fluorescent probes for DNA and corrosion inhibitors (Kovalska *et al.* 2006; Batista *et al.* 2007; Marconato *et al.* 1998). As our continuing investigation on the coordination behavior of the benzothiazole ligand (Chen *et al.* 2008), herein, we report the structure of a mononuclear complex, I.

The molecular structure of the title mononuclear complex is shown in Fig.1 and selected bond lengths and angles are listed in Table 1. The Zn^{II} atom in the mononuclear structure of I exhibits a slightly distorted tetrahedral coordination geometry involving two thiazole N atoms from two different 2-amino-6-methyl-1,3-benzothiazole ligands and two monodentate carboxylate O atoms from two separate 4-nitrobenzoate anions. Both ligands act as typically terminal ligands to coordinate to Zn^{II} ion in monodentate mode with the intramolecular N—H \cdots O hydrogen bonds between amino group of neutral 2-amino-6-methyl-1,3-benzothiazole and anionic 4-nitrobenzoate ligand stabilized the molecular structure.


In the crystal structure, intermolecular N—H \cdots O hydrogen bonds between amino group of neutral 2-amino-6-methyl-1,3-benzothiazole and anionic 4-nitrobenzoate ligands link the discrete mononuclear entities into a one-dimensional supramolecular chain (Fig.2 and Table 2).

S2. Experimental


To an ethanol solution (6.0 ml) containing 2-amino-6-methyl-1,3-benzothiazole (32.8 mg, 0.2 mmol) and 4-nitrobenzoic acid (33.4 mg, 0.2 mmol) was dropwise added an aqueous solution (4.0 ml) of ZnCl₂ (13.6 mg, 0.1 mmol) with constant stirring. The reaction was stirred at room temperature for about 20 min and the precipitate was filtered off, leaving the filtrate to evaporate at room temperature. Yellow block-shaped crystals were obtained with five days. Yield: 50% based on Zn^{II} salt. Anal. Calcd. for C₃₀H₂₄N₆O₈S₂Zn: C, 49.63; H, 3.33; N, 11.57%. Found: C, 49.60; H, 3.31; N, 11.61%.

S3. Refinement

Refinement of F2 against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on F2, conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F2*. The threshold expression of $F2 > 2\sigma(F2)$ is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on *F2* are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

**Figure 1**

The molecular structure of (I). Displacement ellipsoids are drawn at the 30% probability level. Dashed lines indicate hydrogen bonds.

**Figure 2**

Part of the one-dimensional chain of (I) formed by N—H...O hydrogen bonds interactions. Only H atoms involving hydrogen bonds are included.

Bis(2-amino-6-methyl-1,3-benzothiazole- κN^3)bis(4-nitrobenzoato- κO^1)zinc

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{NO}_4)_2(\text{C}_8\text{H}_8\text{N}_2\text{S})_2]$

$M_r = 726.04$

Monoclinic, $P2_1/c$

$a = 13.2240$ (8) Å

$b = 10.7369$ (7) Å

$c = 21.8863$ (13) Å

$\beta = 96.099$ (1)°

$V = 3089.9$ (3) Å³

$Z = 4$

$F(000) = 1488$

$D_x = 1.561$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7342 reflections

$\theta = 2.7\text{--}27.7^\circ$

$\mu = 0.99$ mm⁻¹

$T = 296$ K

Block, yellow

$0.05 \times 0.04 \times 0.03$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.952$, $T_{\max} = 0.971$

15119 measured reflections

5418 independent reflections

4724 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -15 \rightarrow 11$

$k = -10 \rightarrow 12$

$l = -22 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.101$

$S = 1.05$

5418 reflections

426 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0476P)^2 + 4.110P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 1.24$ e Å⁻³

$\Delta\rho_{\min} = -0.42$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

A small void of about 37 Å³ was ignored and was not dealt with SQUEEZE in *PLATON*, because it is so small that no solvent molecules can be kept in.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.26944 (2)	0.00802 (3)	0.480986 (13)	0.02098 (11)
S1	0.47182 (6)	0.30130 (7)	0.40799 (3)	0.03297 (19)
S2	-0.01802 (5)	0.24653 (7)	0.48211 (3)	0.02783 (17)
O1	0.34638 (13)	-0.01142 (17)	0.56266 (8)	0.0247 (4)
O2	0.19228 (14)	-0.0698 (2)	0.58293 (9)	0.0302 (5)
O3	0.5423 (2)	-0.2293 (3)	0.84770 (12)	0.0759 (10)
O4	0.40207 (17)	-0.3229 (2)	0.85604 (10)	0.0459 (6)
O5	0.30961 (15)	-0.27205 (19)	0.47934 (9)	0.0323 (5)
O6	0.20619 (13)	-0.12869 (17)	0.43234 (8)	0.0244 (4)
O7	-0.0259 (2)	-0.5840 (3)	0.24530 (15)	0.0772 (10)
O8	0.0226 (2)	-0.7233 (3)	0.31379 (15)	0.0694 (8)
N1	0.36676 (16)	0.1045 (2)	0.43113 (10)	0.0233 (5)
N2	0.49494 (17)	0.1583 (2)	0.50904 (10)	0.0294 (5)
H2A	0.4785	0.0988	0.5323	0.035*
H2B	0.5445	0.2072	0.5214	0.035*
N3	0.15791 (16)	0.1385 (2)	0.48208 (10)	0.0226 (5)
N4	0.02296 (17)	0.0328 (2)	0.42585 (11)	0.0272 (5)
H4A	0.0622	-0.0258	0.4156	0.033*
H4B	-0.0410	0.0308	0.4135	0.033*
N5	0.4538 (2)	-0.2505 (3)	0.83008 (12)	0.0405 (7)
N6	0.0226 (2)	-0.6164 (3)	0.29378 (15)	0.0495 (8)
C1	0.3274 (2)	0.1474 (3)	0.37255 (12)	0.0244 (6)
C2	0.3728 (2)	0.2560 (3)	0.35280 (13)	0.0284 (6)
C3	0.3393 (2)	0.3119 (3)	0.29694 (14)	0.0358 (7)
H3	0.3691	0.3853	0.2852	0.043*
C4	0.2610 (2)	0.2571 (3)	0.25899 (15)	0.0381 (7)
C5	0.2186 (2)	0.1459 (3)	0.27823 (14)	0.0345 (7)
H5	0.1673	0.1078	0.2525	0.041*
C6	0.2504 (2)	0.0908 (3)	0.33415 (13)	0.0282 (6)
H6	0.2208	0.0172	0.3458	0.034*
C7	0.44390 (19)	0.1748 (3)	0.45406 (12)	0.0257 (6)
C8	0.2202 (3)	0.3160 (4)	0.19792 (16)	0.0505 (9)
H8A	0.2012	0.4008	0.2046	0.076*
H8B	0.1617	0.2703	0.1804	0.076*

H8C	0.2718	0.3139	0.1703	0.076*
C9	0.1747 (2)	0.2472 (3)	0.51684 (12)	0.0243 (6)
C10	0.0875 (2)	0.3181 (3)	0.52264 (13)	0.0266 (6)
C11	0.0913 (2)	0.4282 (3)	0.55639 (13)	0.0308 (6)
H11	0.0323	0.4733	0.5602	0.037*
C12	0.1845 (2)	0.4698 (3)	0.58424 (14)	0.0339 (7)
C13	0.2717 (2)	0.3993 (3)	0.57735 (14)	0.0359 (7)
H13	0.3343	0.4279	0.5953	0.043*
C14	0.2679 (2)	0.2894 (3)	0.54497 (13)	0.0312 (7)
H14	0.3269	0.2437	0.5419	0.037*
C15	0.06120 (19)	0.1267 (2)	0.46081 (12)	0.0228 (6)
C16	0.1951 (3)	0.5880 (3)	0.62144 (16)	0.0459 (8)
H16A	0.1288	0.6198	0.6269	0.069*
H16B	0.2317	0.6487	0.6003	0.069*
H16C	0.2314	0.5709	0.6609	0.069*
C17	0.32872 (19)	-0.0954 (3)	0.66182 (12)	0.0228 (6)
C18	0.4305 (2)	-0.0734 (3)	0.68280 (12)	0.0267 (6)
H18	0.4710	-0.0274	0.6589	0.032*
C19	0.4716 (2)	-0.1198 (3)	0.73917 (13)	0.0306 (6)
H19	0.5390	-0.1041	0.7539	0.037*
C20	0.4098 (2)	-0.1901 (3)	0.77294 (12)	0.0300 (6)
C21	0.3074 (2)	-0.2099 (3)	0.75424 (13)	0.0305 (6)
H21	0.2670	-0.2555	0.7784	0.037*
C22	0.2671 (2)	-0.1600 (3)	0.69863 (13)	0.0277 (6)
H22	0.1982	-0.1697	0.6858	0.033*
C23	0.2843 (2)	-0.0555 (3)	0.59863 (12)	0.0236 (6)
C25	0.18095 (19)	-0.3394 (3)	0.40249 (12)	0.0230 (6)
C26	0.1996 (2)	-0.4654 (3)	0.41508 (13)	0.0271 (6)
H26	0.2463	-0.4882	0.4479	0.032*
C27	0.1490 (2)	-0.5567 (3)	0.37901 (14)	0.0336 (7)
H27	0.1611	-0.6407	0.3873	0.040*
C28	0.0805 (2)	-0.5203 (3)	0.33078 (15)	0.0364 (7)
C29	0.0623 (2)	-0.3960 (3)	0.31583 (14)	0.0364 (7)
H29	0.0177	-0.3741	0.2818	0.044*
C30	0.1120 (2)	-0.3053 (3)	0.35279 (13)	0.0285 (6)
H30	0.0993	-0.2215	0.3444	0.034*
C31	0.23768 (19)	-0.2413 (3)	0.44193 (12)	0.0231 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.01631 (17)	0.02239 (18)	0.02357 (18)	0.00094 (12)	-0.00095 (12)	-0.00048 (12)
S1	0.0306 (4)	0.0337 (4)	0.0346 (4)	-0.0124 (3)	0.0033 (3)	-0.0041 (3)
S2	0.0185 (3)	0.0242 (4)	0.0404 (4)	0.0044 (3)	0.0012 (3)	0.0016 (3)
O1	0.0204 (9)	0.0301 (11)	0.0233 (9)	0.0020 (8)	0.0007 (8)	0.0001 (8)
O2	0.0186 (10)	0.0376 (12)	0.0328 (11)	0.0003 (8)	-0.0050 (8)	0.0015 (9)
O3	0.0495 (16)	0.119 (3)	0.0528 (16)	-0.0286 (17)	-0.0247 (13)	0.0367 (17)
O4	0.0456 (13)	0.0576 (16)	0.0341 (12)	-0.0051 (12)	0.0026 (10)	0.0146 (11)

O5	0.0272 (10)	0.0308 (11)	0.0360 (11)	-0.0024 (9)	-0.0101 (9)	0.0026 (9)
O6	0.0221 (9)	0.0225 (10)	0.0284 (10)	-0.0001 (8)	0.0010 (8)	-0.0011 (8)
O7	0.082 (2)	0.065 (2)	0.075 (2)	0.0077 (16)	-0.0390 (17)	-0.0271 (16)
O8	0.0672 (19)	0.0440 (17)	0.095 (2)	-0.0024 (14)	-0.0005 (16)	-0.0192 (16)
N1	0.0185 (11)	0.0266 (12)	0.0244 (11)	-0.0008 (9)	0.0000 (9)	-0.0019 (9)
N2	0.0215 (11)	0.0353 (14)	0.0303 (13)	-0.0038 (10)	-0.0016 (10)	-0.0061 (11)
N3	0.0181 (11)	0.0233 (12)	0.0261 (11)	0.0027 (9)	0.0010 (9)	0.0011 (9)
N4	0.0157 (11)	0.0276 (13)	0.0373 (13)	0.0010 (9)	-0.0026 (9)	-0.0018 (10)
N5	0.0344 (15)	0.0565 (18)	0.0290 (14)	-0.0043 (13)	-0.0041 (11)	0.0049 (13)
N6	0.0456 (17)	0.0400 (18)	0.061 (2)	0.0002 (14)	-0.0041 (15)	-0.0193 (15)
C1	0.0217 (13)	0.0252 (14)	0.0268 (14)	0.0011 (11)	0.0049 (11)	-0.0009 (11)
C2	0.0268 (14)	0.0284 (15)	0.0305 (15)	-0.0046 (12)	0.0059 (12)	-0.0063 (12)
C3	0.0393 (17)	0.0306 (17)	0.0376 (17)	-0.0060 (14)	0.0049 (14)	0.0050 (13)
C4	0.0400 (17)	0.0360 (18)	0.0377 (17)	-0.0017 (14)	0.0010 (14)	0.0054 (14)
C5	0.0320 (16)	0.0353 (17)	0.0344 (16)	-0.0044 (13)	-0.0049 (13)	0.0024 (13)
C6	0.0258 (14)	0.0262 (15)	0.0316 (15)	-0.0035 (12)	-0.0010 (12)	0.0025 (12)
C7	0.0188 (13)	0.0304 (15)	0.0285 (14)	-0.0008 (11)	0.0057 (11)	-0.0068 (12)
C8	0.058 (2)	0.051 (2)	0.0410 (19)	-0.0048 (18)	0.0004 (17)	0.0123 (17)
C9	0.0240 (14)	0.0223 (14)	0.0265 (14)	0.0023 (11)	0.0018 (11)	0.0017 (11)
C10	0.0242 (14)	0.0240 (14)	0.0312 (15)	0.0035 (11)	0.0009 (11)	0.0040 (12)
C11	0.0322 (15)	0.0265 (15)	0.0334 (15)	0.0103 (12)	0.0024 (12)	0.0006 (12)
C12	0.0378 (17)	0.0258 (15)	0.0367 (17)	0.0067 (13)	-0.0034 (13)	-0.0024 (13)
C13	0.0318 (16)	0.0333 (17)	0.0402 (17)	0.0016 (13)	-0.0069 (13)	-0.0046 (14)
C14	0.0246 (14)	0.0326 (16)	0.0348 (16)	0.0072 (12)	-0.0036 (12)	-0.0041 (13)
C15	0.0209 (13)	0.0213 (14)	0.0263 (14)	0.0025 (11)	0.0028 (11)	0.0049 (11)
C16	0.051 (2)	0.0349 (18)	0.048 (2)	0.0092 (16)	-0.0126 (16)	-0.0103 (15)
C17	0.0212 (13)	0.0246 (14)	0.0225 (13)	0.0039 (11)	0.0014 (10)	-0.0048 (11)
C18	0.0223 (13)	0.0327 (16)	0.0252 (14)	0.0001 (12)	0.0038 (11)	-0.0009 (12)
C19	0.0219 (14)	0.0403 (17)	0.0283 (15)	-0.0017 (12)	-0.0031 (11)	-0.0027 (13)
C20	0.0325 (15)	0.0352 (17)	0.0216 (14)	0.0010 (13)	0.0000 (12)	0.0001 (12)
C21	0.0282 (15)	0.0379 (17)	0.0261 (14)	-0.0029 (13)	0.0052 (12)	-0.0007 (13)
C22	0.0222 (13)	0.0322 (16)	0.0281 (14)	-0.0008 (12)	-0.0005 (11)	-0.0039 (12)
C23	0.0227 (14)	0.0214 (14)	0.0258 (14)	0.0037 (11)	-0.0007 (11)	-0.0031 (11)
C25	0.0190 (13)	0.0241 (14)	0.0260 (14)	0.0006 (11)	0.0029 (10)	-0.0008 (11)
C26	0.0234 (14)	0.0285 (15)	0.0288 (15)	0.0040 (12)	0.0006 (11)	0.0007 (12)
C27	0.0309 (16)	0.0258 (15)	0.0434 (17)	0.0043 (13)	0.0009 (13)	-0.0067 (13)
C28	0.0322 (16)	0.0332 (17)	0.0423 (18)	0.0004 (13)	-0.0027 (14)	-0.0130 (14)
C29	0.0338 (16)	0.0376 (18)	0.0349 (16)	0.0050 (14)	-0.0103 (13)	-0.0052 (14)
C30	0.0263 (14)	0.0263 (15)	0.0317 (15)	0.0020 (12)	-0.0024 (12)	-0.0011 (12)
C31	0.0195 (13)	0.0263 (15)	0.0235 (13)	-0.0012 (11)	0.0020 (11)	0.0010 (11)

Geometric parameters (Å, °)

Zn1—O6	1.9489 (18)	C8—H8A	0.9600
Zn1—O1	1.9717 (18)	C8—H8B	0.9600
Zn1—N3	2.036 (2)	C8—H8C	0.9600
Zn1—N1	2.054 (2)	C9—C14	1.393 (4)
S1—C2	1.754 (3)	C9—C10	1.399 (4)

S1—C7	1.755 (3)	C10—C11	1.392 (4)
S2—C10	1.750 (3)	C11—C12	1.389 (4)
S2—C15	1.754 (3)	C11—H11	0.9300
O1—C23	1.287 (3)	C12—C13	1.401 (4)
O2—C23	1.238 (3)	C12—C16	1.507 (4)
O3—N5	1.214 (3)	C13—C14	1.375 (4)
O4—N5	1.216 (3)	C13—H13	0.9300
O5—C31	1.232 (3)	C14—H14	0.9300
O6—C31	1.289 (3)	C16—H16A	0.9600
O7—N6	1.230 (4)	C16—H16B	0.9600
O8—N6	1.229 (4)	C16—H16C	0.9600
N1—C7	1.324 (3)	C17—C22	1.390 (4)
N1—C1	1.409 (3)	C17—C18	1.396 (4)
N2—C7	1.327 (3)	C17—C23	1.506 (4)
N2—H2A	0.8600	C18—C19	1.387 (4)
N2—H2B	0.8600	C18—H18	0.9300
N3—C15	1.320 (3)	C19—C20	1.383 (4)
N3—C9	1.398 (3)	C19—H19	0.9300
N4—C15	1.332 (4)	C20—C21	1.388 (4)
N4—H4A	0.8600	C21—C22	1.384 (4)
N4—H4B	0.8600	C21—H21	0.9300
N5—C20	1.472 (4)	C22—H22	0.9300
N6—C28	1.474 (4)	C25—C30	1.392 (4)
C1—C6	1.389 (4)	C25—C26	1.397 (4)
C1—C2	1.401 (4)	C25—C31	1.510 (4)
C2—C3	1.391 (4)	C26—C27	1.386 (4)
C3—C4	1.388 (4)	C26—H26	0.9300
C3—H3	0.9300	C27—C28	1.373 (4)
C4—C5	1.403 (4)	C27—H27	0.9300
C4—C8	1.525 (4)	C28—C29	1.389 (4)
C5—C6	1.384 (4)	C29—C30	1.387 (4)
C5—H5	0.9300	C29—H29	0.9300
C6—H6	0.9300	C30—H30	0.9300
O6—Zn1—O1	124.45 (8)	C12—C11—H11	120.5
O6—Zn1—N3	104.78 (8)	C10—C11—H11	120.5
O1—Zn1—N3	111.52 (8)	C11—C12—C13	118.8 (3)
O6—Zn1—N1	110.27 (8)	C11—C12—C16	122.3 (3)
O1—Zn1—N1	104.06 (8)	C13—C12—C16	118.9 (3)
N3—Zn1—N1	98.91 (9)	C14—C13—C12	122.3 (3)
C2—S1—C7	89.64 (13)	C14—C13—H13	118.9
C10—S2—C15	89.45 (13)	C12—C13—H13	118.9
C23—O1—Zn1	106.99 (16)	C13—C14—C9	119.3 (3)
C31—O6—Zn1	120.34 (16)	C13—C14—H14	120.4
C7—N1—C1	110.8 (2)	C9—C14—H14	120.4
C7—N1—Zn1	125.95 (18)	N3—C15—N4	124.7 (2)
C1—N1—Zn1	117.15 (16)	N3—C15—S2	114.8 (2)
C7—N2—H2A	120.0	N4—C15—S2	120.4 (2)

C7—N2—H2B	120.0	C12—C16—H16A	109.5
H2A—N2—H2B	120.0	C12—C16—H16B	109.5
C15—N3—C9	111.4 (2)	H16A—C16—H16B	109.5
C15—N3—Zn1	127.29 (19)	C12—C16—H16C	109.5
C9—N3—Zn1	120.47 (17)	H16A—C16—H16C	109.5
C15—N4—H4A	120.0	H16B—C16—H16C	109.5
C15—N4—H4B	120.0	C22—C17—C18	119.9 (2)
H4A—N4—H4B	120.0	C22—C17—C23	118.6 (2)
O3—N5—O4	122.9 (3)	C18—C17—C23	121.5 (2)
O3—N5—C20	118.0 (3)	C19—C18—C17	120.3 (3)
O4—N5—C20	119.1 (2)	C19—C18—H18	119.8
O8—N6—O7	123.4 (3)	C17—C18—H18	119.8
O8—N6—C28	118.5 (3)	C20—C19—C18	118.2 (3)
O7—N6—C28	118.0 (3)	C20—C19—H19	120.9
C6—C1—C2	119.2 (3)	C18—C19—H19	120.9
C6—C1—N1	125.7 (2)	C19—C20—C21	122.6 (3)
C2—C1—N1	115.1 (2)	C19—C20—N5	119.4 (3)
C3—C2—C1	121.6 (3)	C21—C20—N5	117.9 (3)
C3—C2—S1	129.1 (2)	C22—C21—C20	118.2 (3)
C1—C2—S1	109.4 (2)	C22—C21—H21	120.9
C4—C3—C2	119.4 (3)	C20—C21—H21	120.9
C4—C3—H3	120.3	C21—C22—C17	120.5 (3)
C2—C3—H3	120.3	C21—C22—H22	119.7
C3—C4—C5	118.5 (3)	C17—C22—H22	119.7
C3—C4—C8	121.4 (3)	O2—C23—O1	123.2 (2)
C5—C4—C8	120.0 (3)	O2—C23—C17	119.7 (2)
C6—C5—C4	122.4 (3)	O1—C23—C17	117.0 (2)
C6—C5—H5	118.8	C30—C25—C26	119.7 (3)
C4—C5—H5	118.8	C30—C25—C31	120.5 (2)
C5—C6—C1	118.9 (3)	C26—C25—C31	119.8 (2)
C5—C6—H6	120.6	C27—C26—C25	120.6 (3)
C1—C6—H6	120.6	C27—C26—H26	119.7
N1—C7—N2	124.4 (3)	C25—C26—H26	119.7
N1—C7—S1	115.1 (2)	C28—C27—C26	118.4 (3)
N2—C7—S1	120.4 (2)	C28—C27—H27	120.8
C4—C8—H8A	109.5	C26—C27—H27	120.8
C4—C8—H8B	109.5	C27—C28—C29	122.6 (3)
H8A—C8—H8B	109.5	C27—C28—N6	119.0 (3)
C4—C8—H8C	109.5	C29—C28—N6	118.4 (3)
H8A—C8—H8C	109.5	C30—C29—C28	118.6 (3)
H8B—C8—H8C	109.5	C30—C29—H29	120.7
C14—C9—N3	126.5 (2)	C28—C29—H29	120.7
C14—C9—C10	118.8 (3)	C29—C30—C25	120.1 (3)
N3—C9—C10	114.7 (2)	C29—C30—H30	120.0
C11—C10—C9	121.8 (3)	C25—C30—H30	120.0
C11—C10—S2	128.6 (2)	O5—C31—O6	124.9 (2)
C9—C10—S2	109.6 (2)	O5—C31—C25	119.6 (2)
C12—C11—C10	119.1 (3)	O6—C31—C25	115.4 (2)

O6—Zn1—O1—C23	-61.77 (19)	S2—C10—C11—C12	-178.7 (2)
N3—Zn1—O1—C23	65.32 (18)	C10—C11—C12—C13	0.0 (5)
N1—Zn1—O1—C23	171.00 (17)	C10—C11—C12—C16	179.9 (3)
O1—Zn1—O6—C31	-22.1 (2)	C11—C12—C13—C14	-1.2 (5)
N3—Zn1—O6—C31	-151.97 (19)	C16—C12—C13—C14	178.9 (3)
N1—Zn1—O6—C31	102.47 (19)	C12—C13—C14—C9	1.3 (5)
O6—Zn1—N1—C7	-157.5 (2)	N3—C9—C14—C13	179.5 (3)
O1—Zn1—N1—C7	-21.9 (2)	C10—C9—C14—C13	-0.2 (4)
N3—Zn1—N1—C7	93.1 (2)	C9—N3—C15—N4	179.4 (2)
O6—Zn1—N1—C1	52.8 (2)	Zn1—N3—C15—N4	-11.3 (4)
O1—Zn1—N1—C1	-171.65 (18)	C9—N3—C15—S2	-0.7 (3)
N3—Zn1—N1—C1	-56.7 (2)	Zn1—N3—C15—S2	168.56 (13)
O6—Zn1—N3—C15	7.2 (2)	C10—S2—C15—N3	0.1 (2)
O1—Zn1—N3—C15	-129.9 (2)	C10—S2—C15—N4	-180.0 (2)
N1—Zn1—N3—C15	121.1 (2)	C22—C17—C18—C19	2.5 (4)
O6—Zn1—N3—C9	175.59 (19)	C23—C17—C18—C19	-174.8 (3)
O1—Zn1—N3—C9	38.5 (2)	C17—C18—C19—C20	1.4 (4)
N1—Zn1—N3—C9	-70.6 (2)	C18—C19—C20—C21	-3.7 (5)
C7—N1—C1—C6	175.9 (3)	C18—C19—C20—N5	174.2 (3)
Zn1—N1—C1—C6	-29.9 (3)	O3—N5—C20—C19	4.8 (5)
C7—N1—C1—C2	-3.3 (3)	O4—N5—C20—C19	-172.1 (3)
Zn1—N1—C1—C2	150.81 (19)	O3—N5—C20—C21	-177.2 (3)
C6—C1—C2—C3	3.2 (4)	O4—N5—C20—C21	5.9 (4)
N1—C1—C2—C3	-177.5 (3)	C19—C20—C21—C22	1.9 (5)
C6—C1—C2—S1	-176.4 (2)	N5—C20—C21—C22	-176.0 (3)
N1—C1—C2—S1	2.9 (3)	C20—C21—C22—C17	2.1 (4)
C7—S1—C2—C3	179.2 (3)	C18—C17—C22—C21	-4.3 (4)
C7—S1—C2—C1	-1.3 (2)	C23—C17—C22—C21	173.1 (3)
C1—C2—C3—C4	-1.8 (5)	Zn1—O1—C23—O2	-5.9 (3)
S1—C2—C3—C4	177.7 (2)	Zn1—O1—C23—C17	171.76 (18)
C2—C3—C4—C5	-0.5 (5)	C22—C17—C23—O2	7.0 (4)
C2—C3—C4—C8	178.8 (3)	C18—C17—C23—O2	-175.7 (3)
C3—C4—C5—C6	1.6 (5)	C22—C17—C23—O1	-170.7 (2)
C8—C4—C5—C6	-177.7 (3)	C18—C17—C23—O1	6.6 (4)
C4—C5—C6—C1	-0.3 (5)	C30—C25—C26—C27	-0.9 (4)
C2—C1—C6—C5	-2.1 (4)	C31—C25—C26—C27	-179.4 (2)
N1—C1—C6—C5	178.7 (3)	C25—C26—C27—C28	0.1 (4)
C1—N1—C7—N2	178.7 (2)	C26—C27—C28—C29	1.9 (5)
Zn1—N1—C7—N2	27.3 (4)	C26—C27—C28—N6	-177.2 (3)
C1—N1—C7—S1	2.3 (3)	O8—N6—C28—C27	13.7 (5)
Zn1—N1—C7—S1	-149.09 (14)	O7—N6—C28—C27	-168.6 (3)
C2—S1—C7—N1	-0.6 (2)	O8—N6—C28—C29	-165.4 (3)
C2—S1—C7—N2	-177.2 (2)	O7—N6—C28—C29	12.2 (5)
C15—N3—C9—C14	-178.7 (3)	C27—C28—C29—C30	-2.9 (5)
Zn1—N3—C9—C14	11.2 (4)	N6—C28—C29—C30	176.2 (3)
C15—N3—C9—C10	1.0 (3)	C28—C29—C30—C25	2.0 (5)
Zn1—N3—C9—C10	-169.03 (19)	C26—C25—C30—C29	-0.2 (4)

C14—C9—C10—C11	-1.0 (4)	C31—C25—C30—C29	178.3 (3)
N3—C9—C10—C11	179.3 (2)	Zn1—O6—C31—O5	-2.6 (4)
C14—C9—C10—S2	178.9 (2)	Zn1—O6—C31—C25	178.49 (16)
N3—C9—C10—S2	-0.9 (3)	C30—C25—C31—O5	-168.5 (3)
C15—S2—C10—C11	-179.8 (3)	C26—C25—C31—O5	10.0 (4)
C15—S2—C10—C9	0.4 (2)	C30—C25—C31—O6	10.5 (4)
C9—C10—C11—C12	1.0 (4)	C26—C25—C31—O6	-171.1 (2)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N2—H2A...O1	0.86	2.27	3.008 (3)	144
N2—H2B...O5 ⁱ	0.86	2.05	2.845 (3)	153
N4—H4A...O6	0.86	2.20	2.971 (3)	149
N4—H4B...O2 ⁱⁱ	0.86	2.05	2.860 (3)	156

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x, -y, -z+1$.