

{4-Hydroxy-*N'*-[(2*E*,3*Z*)-4-oxido-4-phenylbut-3-en-2-ylidene]benzohydrazidato}dimethyltin(IV)

Md. Abu Affan,^a† Norrihan B. Sam,^a Fasihuddin B. Ahmad,^a Fraser White^b and Edward R. T. Tiekink^{c*}

^aFaculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia, ^bAgilent Technologies UK Ltd, 10 Mead Road, Oxford Industrial Park, Oxford OX5 1QU, England, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekink@gmail.com


Received 15 June 2011; accepted 16 June 2011

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.032; wR factor = 0.081; data-to-parameter ratio = 13.6.

The Sn^{IV} atom in the title compound, [Sn(CH₃)₂-(C₁₇H₁₄N₂O₃)], is five-coordinated within a C₂N₂O donor set provided by the *N,N,O*-tridentate ligand and two methyl groups. The resultant coordination geometry is intermediate between trigonal-bipyramidal and square-pyramidal. In the crystal, supramolecular zigzag chains propagating along the *c*-axis direction are mediated by O—H···O hydrogen bonds, and weak C—H··· π interactions consolidate the packing.

Related literature

For background to the biological interest of related compounds, see: Affan *et al.* (2010). For related structures, see: Affan *et al.* (2009, 2011). For additional structural analysis, see: Addison *et al.* (1984).



Experimental

Crystal data

[Sn(CH₃)₂(C₁₇H₁₄N₂O₃)]
 $M_r = 443.06$
 Monoclinic, $P2_1/c$
 $a = 8.0784$ (2) Å
 $b = 20.5410$ (5) Å
 $c = 11.1678$ (3) Å
 $\beta = 93.025$ (2)°
 $V = 1850.58$ (8) Å³
 $Z = 4$
 Cu $K\alpha$ radiation
 $\mu = 11.15$ mm⁻¹
 $T = 150$ K
 $0.22 \times 0.16 \times 0.10$ mm

† Additional correspondence author, e-mail: maaffan@yahoo.com.

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
 Absorption correction: analytical (*CrysAlis PRO*; Agilent, 2011)
 $T_{\min} = 0.696$, $T_{\max} = 0.822$
 5718 measured reflections
 3124 independent reflections
 2690 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.081$
 $S = 1.00$
 3124 reflections
 230 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.76$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.72$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Sn—O1	2.156 (3)	Sn—C18	2.105 (4)
Sn—O3	2.099 (3)	Sn—C19	2.112 (4)
Sn—N2	2.148 (3)		
O1—Sn—O3	155.08 (10)	C18—Sn—C19	124.65 (18)

Table 2

Hydrogen-bond geometry (Å, °).

*Cg*1, *Cg*2 and *Cg*3 are the centroids of the C12–C17, Sn, O1, C1, N1, N2 and C2–C7 rings, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2 _o ···O1 ⁱ	0.84	1.91	2.702 (3)	156
C4—H4··· <i>Cg</i> 1 ⁱⁱ	0.95	2.91	3.624 (4)	133
C9—H9 _c ··· <i>Cg</i> 2 ⁱⁱⁱ	0.98	2.88	3.777 (4)	152
C16—H16··· <i>Cg</i> 3 ^{iv}	0.95	2.89	3.668 (4)	140

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x, -y, -z + 1$; (iv) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

We thank MOSTI (grant No. 06–01–09–SF0046) and the Universiti Malaysia Sarawak for support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5915).

References

- Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
 Affan, M. A., Foo, S. W., Jusoh, I., Hanapi, S. & Tiekink, E. R. T. (2009). *Inorg. Chim. Acta*, **362**, 5031–5037.
 Affan, M. A., Sam, N. B., Ahmad, F. B. & Tiekink, E. R. T. (2010). *Acta Cryst. E66*, m924.
 Affan, M. A., Sam, N. B., Ahmad, F. B., White, F. & Tiekink, E. R. T. (2011). *Acta Cryst. E67*, m963–m964.
 Agilent (2011). *CrysAlis PRO*. Agilent Technologies UK Ltd, Oxford, UK.
 Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
 Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, m965 [doi:10.1107/S1600536811023506]

{4-Hydroxy-*N'*-[(2*E*,3*Z*)-4-oxido-4-phenylbut-3-en-2-ylidene]benzohydrazidato}dimethyltin(IV)

Md. Abu Affan, Norrihan B. Sam, Fasihuddin B. Ahmad, Fraser White and Edward R. T. Tiekink

S1. Comment

The title compound, (I), was examined in connection with on-going structural studies (Affan *et al.*, 2010) of organotin derivatives of biological interest (Affan *et al.*, 2009), and compliments the structure of the diphenyltin analogue (Affan *et al.*, 2011).

The Sn atom in (I), Fig. 1, is five-coordinated by the tridentate ligand and two methyl groups, Table 1. The resulting C₂NO₂ donor set defines a coordination geometry intermediate between square pyramidal and trigonal bipyramidal geometry. This is quantified by the value of $\tau = 0.51$ which compare to the τ values of 0.0 and 1.0 for ideal square pyramidal and trigonal bipyramidal geometries, respectively (Addison *et al.*, 1984). For comparison, the values of τ for the two independent molecules in the structure of the diphenyltin analogue are 0.55 and 0.47 (Affan *et al.*, 2011).

While the five-membered SnCN₂O chelate ring is almost planar with a r.m.s. deviation = 0.063 Å [max. deviations of 0.039 (1) and -0.052 (2) Å for the Sn and O1 atoms, respectively], there is considerable distortion in the SnC₃NO six-membered chelate [r.m.s. deviation = 0.226 Å] with the Sn and O3 atoms lying -0.209 (1) and 0.245 (3) Å out of the least-squares plane. Each of the benzene rings is twisted out of the plane from the adjacent chelate ring as seen in the O1—C1—C2—C3 and O3—C11—C12—C13 torsion angles of 13.7 (5) and -150.4 (4)°, respectively. The dihedral angle between the two benzene rings is 68.14 (18)°, indicating a twist in the tridentate ligand.


The crystal packing is dominated by O—H...O hydrogen bonding, Table 2, which leads to a zigzag supramolecular chain along the *c* axis, Fig. 2. These are consolidated in the crystal structure by C—H... π interactions, Table 2.

S2. Experimental

Benzoylacetone 4-hydroxybenzhydrazone (0.59 g, 2 mmol) was dissolved in distilled methanol (20 ml) under a nitrogen atmosphere. Potassium hydroxide (0.23 g, 4 mmol) dissolved in methanol (10 ml) was added drop wise to the solution. The colour of the solution changed from yellow to orange. The resulting mixture was refluxed for 1 h and then treated with dimethyltin dichloride (0.439 g, 2 mmol) in distilled methanol (10 ml). The resulting mixture was heated under reflux conditions for 4 h and allowed to cool to room temperature. Potassium chloride (KCl) was removed *via* filtration. The filtrate was evaporated to dryness using a rotary evaporator to yield yellow microcrystals. The microcrystals were filtered off and washed with ethanol and dried *in vacuo* over P₂O₅ overnight. Yellow blocks of (I) were obtained by slow evaporation of its acetone solution at room temperature. Yield: 0.94 g, 75%. *M.pt.*: 504–506 K. IR (ν_{\max} , cm⁻¹, KBr): 3569 (OH), 1591 (C=N—N=C), 949 (N—N), 562 (Sn—C), 525 (Sn—O), 447 (Sn—N).

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (O—H = 0.84 Å; C—H = 0.95 to 0.98 Å) and were included in the refinement in the riding model approximation with $U_{\text{iso}}(\text{H})$ set to 1.2- $U_{\text{eq}}(\text{C})$ and 1.5- $U_{\text{eq}}(\text{O, methyl-C})$.

**Figure 1**

The molecular structure of (I) showing displacement ellipsoids at the 50% probability level.




Figure 2

A view of the supramolecular chain aligned along [001] in (I). The O—H···O hydrogen bonds are shown as orange dashed lines.




Figure 3

A view in projection down the c axis of the crystal packing in (I). The O—H...O hydrogen bonds and C—H... π interactions are shown as orange and purple dashed lines, respectively.

{4-Hydroxy-*N*-[(2*E*,3*Z*)-4-oxido-4-phenylbut-3-en-2-ylidene]benzohydrazidato}dimethyltin(IV)

Crystal data

[Sn(CH₃)₂(C₁₇H₁₄N₂O₃)]

$M_r = 443.06$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.0784$ (2) Å

$b = 20.5410$ (5) Å

$c = 11.1678$ (3) Å

$\beta = 93.025$ (2)°

$V = 1850.58$ (8) Å³

$Z = 4$

$F(000) = 888$

$D_x = 1.590$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Cell parameters from 3495 reflections

$\theta = 4.0$ – 74.2 °

$\mu = 11.15$ mm⁻¹

$T = 150$ K

Block, yellow

$0.22 \times 0.16 \times 0.10$ mm

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas detector
Radiation source: SuperNova (Cu) X-ray
Source
Mirror monochromator
 ω scans
Absorption correction: analytical
(*CrysAlis PRO*; Agilent, 2011)
 $T_{\min} = 0.696$, $T_{\max} = 0.822$

5718 measured reflections
3124 independent reflections
2690 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$
 $\theta_{\max} = 65.0^\circ$, $\theta_{\min} = 4.5^\circ$
 $h = -6 \rightarrow 9$
 $k = -24 \rightarrow 24$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.081$
 $S = 1.00$
3124 reflections
230 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0384P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.76 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.72 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Agilent Technologies (2011) *CrysAlis PRO* Software system, version 1.171.34.49, Agilent Technologies UK Ltd, Oxford, UK

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn	0.18670 (3)	0.011355 (13)	0.22279 (2)	0.02375 (11)
O1	0.0690 (3)	0.09638 (13)	0.2930 (2)	0.0267 (6)
O2	-0.0780 (3)	0.34947 (13)	0.5942 (2)	0.0309 (6)
H2o	-0.0354	0.3553	0.6636	0.046*
O3	0.3618 (3)	-0.06412 (14)	0.2237 (2)	0.0333 (7)
N1	0.1880 (4)	0.06513 (16)	0.4764 (3)	0.0242 (7)
N2	0.2394 (4)	0.01068 (16)	0.4134 (3)	0.0232 (7)
C1	0.1054 (4)	0.10630 (19)	0.4079 (3)	0.0230 (8)
C2	0.0519 (4)	0.16830 (19)	0.4610 (3)	0.0208 (8)
C3	-0.0628 (5)	0.2084 (2)	0.3999 (3)	0.0263 (9)
H3	-0.1131	0.1942	0.3256	0.032*
C4	-0.1051 (5)	0.2685 (2)	0.4452 (3)	0.0275 (9)
H4	-0.1835	0.2952	0.4020	0.033*
C5	-0.0331 (5)	0.28957 (19)	0.5535 (3)	0.0240 (8)

C6	0.0800 (4)	0.2492 (2)	0.6178 (3)	0.0250 (8)
H6	0.1282	0.2631	0.6929	0.030*
C7	0.1213 (4)	0.18929 (19)	0.5719 (3)	0.0245 (8)
H7	0.1975	0.1621	0.6159	0.029*
C8	0.2979 (5)	-0.0386 (2)	0.4775 (3)	0.0247 (8)
C9	0.3027 (5)	-0.0339 (2)	0.6122 (3)	0.0311 (9)
H9A	0.3736	0.0027	0.6385	0.047*
H9B	0.3474	-0.0744	0.6472	0.047*
H9C	0.1903	-0.0269	0.6385	0.047*
C10	0.3544 (5)	-0.0971 (2)	0.4265 (3)	0.0268 (9)
H10	0.3750	-0.1326	0.4797	0.032*
C11	0.3827 (5)	-0.1079 (2)	0.3075 (3)	0.0255 (9)
C12	0.4463 (4)	-0.1711 (2)	0.2669 (3)	0.0243 (8)
C13	0.4076 (5)	-0.2292 (2)	0.3228 (4)	0.0286 (9)
H13	0.3440	-0.2285	0.3919	0.034*
C14	0.4607 (5)	-0.2880 (2)	0.2787 (4)	0.0340 (10)
H14	0.4317	-0.3274	0.3169	0.041*
C15	0.5565 (5)	-0.2900 (2)	0.1787 (4)	0.0365 (10)
H15	0.5938	-0.3304	0.1489	0.044*
C16	0.5966 (5)	-0.2323 (2)	0.1232 (4)	0.0351 (10)
H16	0.6629	-0.2332	0.0554	0.042*
C17	0.5410 (5)	-0.1731 (2)	0.1658 (4)	0.0295 (9)
H17	0.5674	-0.1339	0.1260	0.035*
C18	0.3273 (6)	0.0647 (2)	0.1034 (4)	0.0379 (11)
H18A	0.4406	0.0701	0.1380	0.057*
H18B	0.2767	0.1076	0.0896	0.057*
H18C	0.3302	0.0413	0.0271	0.057*
C19	-0.0261 (6)	-0.0439 (2)	0.1720 (4)	0.0451 (12)
H19A	-0.0003	-0.0739	0.1074	0.068*
H19B	-0.1158	-0.0146	0.1439	0.068*
H19C	-0.0611	-0.0688	0.2411	0.068*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn	0.02822 (16)	0.02438 (17)	0.01860 (15)	0.00018 (11)	0.00060 (10)	-0.00033 (10)
O1	0.0341 (15)	0.0238 (15)	0.0217 (14)	0.0056 (12)	-0.0027 (10)	-0.0038 (11)
O2	0.0410 (17)	0.0274 (16)	0.0237 (15)	0.0055 (13)	-0.0045 (12)	-0.0050 (12)
O3	0.0419 (17)	0.0311 (16)	0.0277 (15)	0.0123 (14)	0.0084 (12)	0.0059 (13)
N1	0.0271 (17)	0.0232 (18)	0.0222 (16)	0.0013 (14)	0.0008 (13)	-0.0027 (14)
N2	0.0250 (16)	0.0260 (18)	0.0192 (16)	0.0003 (14)	0.0042 (13)	0.0003 (14)
C1	0.0199 (18)	0.029 (2)	0.0200 (19)	-0.0050 (16)	0.0023 (14)	0.0008 (17)
C2	0.0194 (18)	0.024 (2)	0.0195 (18)	-0.0024 (16)	0.0012 (14)	0.0012 (15)
C3	0.025 (2)	0.033 (2)	0.0200 (19)	-0.0033 (18)	-0.0027 (15)	-0.0012 (17)
C4	0.029 (2)	0.029 (2)	0.0241 (19)	0.0065 (18)	-0.0018 (15)	-0.0001 (17)
C5	0.026 (2)	0.025 (2)	0.0218 (19)	-0.0037 (17)	0.0009 (14)	-0.0006 (16)
C6	0.025 (2)	0.029 (2)	0.0204 (18)	-0.0059 (17)	-0.0005 (15)	-0.0040 (17)
C7	0.0220 (19)	0.028 (2)	0.0238 (19)	-0.0009 (17)	-0.0001 (14)	0.0034 (17)

C8	0.0236 (19)	0.029 (2)	0.0211 (19)	-0.0008 (18)	0.0019 (14)	0.0010 (17)
C9	0.044 (2)	0.029 (2)	0.020 (2)	0.002 (2)	0.0029 (17)	-0.0001 (18)
C10	0.030 (2)	0.027 (2)	0.0237 (19)	0.0031 (18)	0.0024 (15)	0.0033 (17)
C11	0.0230 (19)	0.027 (2)	0.026 (2)	-0.0023 (17)	0.0008 (15)	-0.0008 (17)
C12	0.0177 (18)	0.028 (2)	0.027 (2)	0.0000 (16)	-0.0033 (14)	-0.0035 (17)
C13	0.023 (2)	0.032 (2)	0.031 (2)	-0.0025 (18)	0.0005 (16)	-0.0011 (18)
C14	0.027 (2)	0.027 (2)	0.047 (3)	-0.0004 (19)	-0.0059 (18)	-0.001 (2)
C15	0.031 (2)	0.036 (3)	0.041 (3)	0.009 (2)	-0.0062 (18)	-0.011 (2)
C16	0.028 (2)	0.046 (3)	0.031 (2)	0.008 (2)	0.0005 (17)	-0.004 (2)
C17	0.026 (2)	0.034 (2)	0.028 (2)	0.0020 (19)	0.0032 (15)	0.0007 (19)
C18	0.043 (3)	0.040 (3)	0.031 (2)	-0.008 (2)	0.0029 (18)	0.003 (2)
C19	0.045 (3)	0.042 (3)	0.048 (3)	-0.011 (2)	0.006 (2)	-0.020 (2)

Geometric parameters (Å, °)

Sn—O1	2.156 (3)	C8—C9	1.506 (5)
Sn—O3	2.099 (3)	C9—H9A	0.9800
Sn—N2	2.148 (3)	C9—H9B	0.9800
Sn—C18	2.105 (4)	C9—H9C	0.9800
Sn—C19	2.112 (4)	C10—C11	1.377 (5)
O1—C1	1.317 (4)	C10—H10	0.9500
O2—C5	1.367 (5)	C11—C12	1.476 (5)
O2—H2O	0.8400	C12—C13	1.390 (6)
O3—C11	1.303 (5)	C12—C17	1.398 (5)
N1—C1	1.300 (5)	C13—C14	1.380 (6)
N1—N2	1.396 (4)	C13—H13	0.9500
N2—C8	1.313 (5)	C14—C15	1.393 (6)
C1—C2	1.479 (5)	C14—H14	0.9500
C2—C3	1.391 (5)	C15—C16	1.384 (6)
C2—C7	1.400 (5)	C15—H15	0.9500
C3—C4	1.384 (6)	C16—C17	1.388 (6)
C3—H3	0.9500	C16—H16	0.9500
C4—C5	1.383 (5)	C17—H17	0.9500
C4—H4	0.9500	C18—H18A	0.9800
C5—C6	1.404 (5)	C18—H18B	0.9800
C6—C7	1.381 (5)	C18—H18C	0.9800
C6—H6	0.9500	C19—H19A	0.9800
C7—H7	0.9500	C19—H19B	0.9800
C8—C10	1.417 (5)	C19—H19C	0.9800
O3—Sn—C18	90.09 (15)	C8—C9—H9B	109.5
O3—Sn—C19	98.26 (16)	H9A—C9—H9B	109.5
O1—Sn—O3	155.08 (10)	C8—C9—H9C	109.5
C18—Sn—C19	124.65 (18)	H9A—C9—H9C	109.5
O3—Sn—N2	83.82 (11)	H9B—C9—H9C	109.5
C18—Sn—N2	123.05 (15)	C11—C10—C8	126.7 (4)
C19—Sn—N2	112.24 (16)	C11—C10—H10	116.7
C18—Sn—O1	94.10 (15)	C8—C10—H10	116.7

C19—Sn—O1	99.46 (15)	O3—C11—C10	124.1 (4)
N2—Sn—O1	73.31 (11)	O3—C11—C12	114.8 (3)
C1—O1—Sn	113.6 (2)	C10—C11—C12	121.0 (4)
C5—O2—H2O	109.5	C13—C12—C17	118.8 (4)
C11—O3—Sn	125.1 (2)	C13—C12—C11	121.8 (3)
C1—N1—N2	112.4 (3)	C17—C12—C11	119.3 (4)
C8—N2—N1	116.8 (3)	C14—C13—C12	120.7 (4)
C8—N2—Sn	126.3 (3)	C14—C13—H13	119.7
N1—N2—Sn	116.5 (2)	C12—C13—H13	119.7
N1—C1—O1	123.6 (4)	C13—C14—C15	120.5 (4)
N1—C1—C2	118.4 (3)	C13—C14—H14	119.7
O1—C1—C2	118.0 (3)	C15—C14—H14	119.7
C3—C2—C7	118.5 (4)	C16—C15—C14	119.1 (4)
C3—C2—C1	121.0 (3)	C16—C15—H15	120.4
C7—C2—C1	120.5 (3)	C14—C15—H15	120.4
C4—C3—C2	121.3 (4)	C15—C16—C17	120.6 (4)
C4—C3—H3	119.3	C15—C16—H16	119.7
C2—C3—H3	119.3	C17—C16—H16	119.7
C5—C4—C3	119.9 (4)	C16—C17—C12	120.3 (4)
C5—C4—H4	120.1	C16—C17—H17	119.9
C3—C4—H4	120.1	C12—C17—H17	119.9
O2—C5—C4	117.8 (3)	Sn—C18—H18A	109.5
O2—C5—C6	122.6 (3)	Sn—C18—H18B	109.5
C4—C5—C6	119.6 (4)	H18A—C18—H18B	109.5
C7—C6—C5	120.0 (3)	Sn—C18—H18C	109.5
C7—C6—H6	120.0	H18A—C18—H18C	109.5
C5—C6—H6	120.0	H18B—C18—H18C	109.5
C6—C7—C2	120.7 (4)	Sn—C19—H19A	109.5
C6—C7—H7	119.7	Sn—C19—H19B	109.5
C2—C7—H7	119.7	H19A—C19—H19B	109.5
N2—C8—C10	123.3 (3)	Sn—C19—H19C	109.5
N2—C8—C9	119.0 (4)	H19A—C19—H19C	109.5
C10—C8—C9	117.7 (4)	H19B—C19—H19C	109.5
C8—C9—H9A	109.5		
O3—Sn—O1—C1	-17.4 (4)	C3—C4—C5—O2	179.6 (3)
C18—Sn—O1—C1	-116.5 (3)	C3—C4—C5—C6	-1.3 (6)
C19—Sn—O1—C1	117.4 (3)	O2—C5—C6—C7	-179.7 (3)
N2—Sn—O1—C1	6.8 (2)	C4—C5—C6—C7	1.2 (6)
C18—Sn—O3—C11	158.3 (3)	C5—C6—C7—C2	0.3 (6)
C19—Sn—O3—C11	-76.6 (3)	C3—C2—C7—C6	-1.7 (5)
N2—Sn—O3—C11	35.0 (3)	C1—C2—C7—C6	175.6 (3)
O1—Sn—O3—C11	58.3 (4)	N1—N2—C8—C10	179.9 (3)
C1—N1—N2—C8	-168.5 (3)	Sn—N2—C8—C10	8.0 (5)
C1—N1—N2—Sn	4.2 (4)	N1—N2—C8—C9	1.2 (5)
O3—Sn—N2—C8	-24.1 (3)	Sn—N2—C8—C9	-170.7 (3)
C18—Sn—N2—C8	-110.2 (3)	N2—C8—C10—C11	11.3 (6)
C19—Sn—N2—C8	72.3 (3)	C9—C8—C10—C11	-170.0 (4)

O1—Sn—N2—C8	165.9 (3)	Sn—O3—C11—C10	-30.7 (5)
O3—Sn—N2—N1	164.0 (3)	Sn—O3—C11—C12	151.4 (3)
C18—Sn—N2—N1	78.0 (3)	C8—C10—C11—O3	0.6 (6)
C19—Sn—N2—N1	-99.5 (3)	C8—C10—C11—C12	178.3 (4)
O1—Sn—N2—N1	-5.9 (2)	O3—C11—C12—C13	-150.4 (4)
N2—N1—C1—O1	2.4 (5)	C10—C11—C12—C13	31.7 (5)
N2—N1—C1—C2	-176.3 (3)	O3—C11—C12—C17	26.3 (5)
Sn—O1—C1—N1	-7.7 (5)	C10—C11—C12—C17	-151.6 (4)
Sn—O1—C1—C2	171.1 (2)	C17—C12—C13—C14	-0.5 (6)
N1—C1—C2—C3	-167.5 (3)	C11—C12—C13—C14	176.2 (4)
O1—C1—C2—C3	13.7 (5)	C12—C13—C14—C15	1.1 (6)
N1—C1—C2—C7	15.3 (5)	C13—C14—C15—C16	-0.6 (6)
O1—C1—C2—C7	-163.5 (3)	C14—C15—C16—C17	-0.6 (6)
C7—C2—C3—C4	1.7 (6)	C15—C16—C17—C12	1.3 (6)
C1—C2—C3—C4	-175.6 (3)	C13—C12—C17—C16	-0.7 (6)
C2—C3—C4—C5	-0.2 (6)	C11—C12—C17—C16	-177.5 (4)

Hydrogen-bond geometry (Å, °)

Cg1, Cg2 and Cg3 are the centroids of the C12—C17, Sn,O1,C1,N1,N2 and C2—C7 rings, respectively.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots O1 ⁱ	0.84	1.91	2.702 (3)	156
C4—H4 \cdots Cg1 ⁱⁱ	0.95	2.91	3.624 (4)	133
C9—H9 \cdots Cg2 ⁱⁱⁱ	0.98	2.88	3.777 (4)	152
C16—H16 \cdots Cg3 ^{iv}	0.95	2.89	3.668 (4)	140

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x, y+1/2, -z+1/2$; (iii) $-x, -y, -z+1$; (iv) $-x+1, y-1/2, -z+1/2$.