

6-Phenyl-6,7-dihydrodibenzo[*c,f*][1,5]-azabismocin-12(5*H*)-yl perchlorate

 Xiao-Wen Zhang^{a*} and Ting Fan^b

^aCollege of Environment Protection and Safety Engineering, University of South China, Hengyang 421001, People's Republic of China, and ^bKey Laboratory of Pollution Control and Resource Use of Hunan Province, University of South China, Hengyang 421001, People's Republic of China
Correspondence e-mail: shawn_zhang@sina.com


Received 4 May 2011; accepted 31 May 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.018$ Å; R factor = 0.065; wR factor = 0.163; data-to-parameter ratio = 13.4.

In the title compound, $[\text{Bi}(\text{C}_{20}\text{H}_{17}\text{N})(\text{ClO}_4)]$ or $\text{C}_{20}\text{H}_{17}\text{BiClNO}_4$, the Bi^{III} ion assumes a distorted ψ trigonal-bipyramidal geometry, with two C atoms and the electron lone pair of the Bi atom at the equatorial positions and an amine N atom and a perchlorate O atom at the apical positions. Weak intermolecular C—H...O hydrogen bonding is present in the crystal structure.

Related literature

For the synthesis of 12-chloro-6-phenyl-5,6,7,12-tetrahydrodibenzo[*c,f*][1,5]azabismocine, see: Zhang *et al.* (2009). For general background, see: Shimada *et al.* (2004); Yin *et al.* (2008); Zhang *et al.* (2010); Tan & Zhang (2011). For related structures, see: Ohkata *et al.* (1989); Minoura *et al.* (1999).



Experimental

Crystal data

$[\text{Bi}(\text{C}_{20}\text{H}_{17}\text{N})(\text{ClO}_4)]$
 $M_r = 579.78$
 Monoclinic, $P2_1/c$
 $a = 12.0635$ (10) Å
 $b = 14.0755$ (12) Å
 $c = 11.5121$ (10) Å
 $\beta = 107.590$ (2)°
 $V = 1863.4$ (3) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 9.63$ mm⁻¹

$T = 293$ K
 $0.32 \times 0.21 \times 0.20$ mm

Data collection

Bruker SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\text{min}} = 0.100$, $T_{\text{max}} = 0.145$

9267 measured reflections
 3279 independent reflections
 2585 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.166$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.163$
 $S = 1.02$
 3279 reflections

244 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 4.89$ e Å⁻³
 $\Delta\rho_{\text{min}} = -4.14$ e Å⁻³

Table 1

Selected bond lengths (Å).

Bi—N1	2.387 (10)	Bi—C1	2.245 (13)
Bi—O1	2.546 (10)	Bi—C8	2.204 (12)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4...O3 ⁱ	0.93	2.46	3.137 (17)	130
C14—H14B...O2 ⁱⁱ	0.97	2.55	3.398 (16)	146

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge the National Science Foundation of China (grant No. 51074093) for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5208).

References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Minoura, M., Kanamori, Y., Miyake, A. & Akiba, K. (1999). *Chem. Lett.* pp. 861–862.
 Ohkata, K., Takemoto, S., Ohnishi, M. & Akiba, K. Y. (1989). *Tetrahedron Lett.* **30**, 4841–4844.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Shimada, S., Yamazaki, O., Tanaka, T., Suzuki, Y. & Tanaka, M. (2004). *J. Organomet. Chem.* **689**, 3012–3023.
 Tan, N. & Zhang, X. (2011). *Acta Cryst.* **E67**, m252.
 Yin, S., Maruyama, J., Yamashita, T. & Shimada, S. (2008). *Angew. Chem. Int. Ed.* **47**, 6590–6593.
 Zhang, X.-W., Qiu, R.-H., Tan, N.-Y., Yin, S.-F., Xia, J., Au, C.-T. & Luo, S.-L. (2010). *Tetrahedron Lett.* **51**, 153–156.
 Zhang, X.-W., Xia, J., Yan, H.-W., Luo, S.-L., Yin, S.-F., Au, C.-T. & Wong, W.-Y. (2009). *J. Organomet. Chem.* **694**, 3019–3026.

supporting information

Acta Cryst. (2011). E67, m875 [doi:10.1107/S1600536811021039]

6-Phenyl-6,7-dihydrodibenzo[*c,f*][1,5]azabismocin-12(5*H*)-yl perchlorate**Xiao-Wen Zhang and Ting Fan****S1. Comment**

Bismuth is a nontoxic and noncarcinogenic element and many of its compounds are low in toxicity and can be safely used in areas such as medicine, catalysis, and synthesis (Shimada *et al.*, 2004; Yin *et al.*, 2008; Zhang, Qiu, Tan *et al.*, 2010). The 5,6,7,12-tetrahydrodibenzo[*c,f*][1,5]azabismocine framework is highly stable as a organobismuth Fragment because the weakly coordination exists between bismuth and nitrogen atoms on 1,5-azabismocine (Ohkata *et al.*, 1989; Minoura *et al.*, 1999), and therefore, is suitable for the study of organobismuth compounds bearing various groups on the bismuth and nitrogen atom for potential uses.

In the present paper, we report the crystal structure of the title compound (Fig. 1). The central bismuth-containing part of the complex exhibits a distorted pseudo trigonal-bipyramidal structure. The C (8), C (1) atoms and a lone electron pair of the Bi atom exist at the equatorial positions while the N (1) and O (1) atoms are located at the apical positions. The Bi–C (8) and Bi–C (1) distance is 2.250 (13) Å and 2.204 (12) Å, respectively. The C (8)–Bi–C (1) angle is 92.5 (5) ° while the N (1)–Bi–O (1) angle is 154.0 (3)° (rather than 180°). The Bi–N (1) distance (2.388 (10) Å) is shorter than 2.607 (5) Å of the precursor, C₆H₅N(CH₂C₆H₄)₂BiCl. The Bi–O (1) distance (2.546 (10) Å) is shorter than Bi–Cl(1) distance 2.597 (19) Å also (Zhang, Xia, Yan *et al.*, 2009).

S2. Experimental

The following procedures are recommended for synthesis of the title compound (I): 12-chloro-6-phenyl-5,6,7,12-tetrahydrodibenzo[*c,f*][1,5]azabismocine (0.516 g, 1.0 mmol) was dissolved in 15 ml THF, then a solution of AgClO₄ (0.207 g, 1.0 mmol) in 15.0 ml THF was added. After the mixture was stirred in the dark at room temperature for 2 h, it was filtered. The filtrate mixed with 1.0 ml hexane was refrigerated for 24 h, giving colorless crystals (0.539 g, 93.0%).

S3. Refinement

All H atoms were positioned geometrically and refined using a riding model, with C–H = 0.93 Å for aryl, 0.98 Å methine and 0.97 Å for methylene H atoms, respectively. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for all H atoms.




Figure 1

The molecular structure of the title compound with atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. All hydrogen atoms are omitted for clarity.

6-Phenyl-6,7-dihydrodibenzo[*c,f*][1,5]azabismocin-12(5*H*)-yl perchlorate

Crystal data

[Bi(C₂₀H₁₇N)(ClO₄)]

M_r = 579.78

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 12.0635 (10) Å

b = 14.0755 (12) Å

c = 11.5121 (10) Å

β = 107.590 (2)°

V = 1863.4 (3) Å³

Z = 4

F(000) = 1104

D_x = 2.067 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2936 reflections

θ = 2.7–24.2°

μ = 9.63 mm⁻¹

T = 293 K

Prismatic, colorless

0.32 × 0.21 × 0.20 mm

Data collection

Bruker SMART 1000 CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: empirical (using
intensity measurements)
(*SADABS*; Bruker, 2001)
 $T_{\min} = 0.100$, $T_{\max} = 0.145$

9267 measured reflections
3279 independent reflections
2585 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.166$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.8^\circ$
 $h = -10 \rightarrow 14$
 $k = -13 \rightarrow 16$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.163$
 $S = 1.02$
3279 reflections
244 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0816P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 4.89 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -4.14 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Bi	0.74646 (4)	0.77309 (3)	0.84604 (4)	0.0321 (2)
C11	0.7240 (3)	1.0213 (2)	0.8825 (3)	0.0378 (7)
N1	0.7225 (7)	0.6171 (8)	0.7606 (7)	0.033 (2)
O1	0.7940 (9)	0.9494 (7)	0.8480 (10)	0.063 (3)
O2	0.6291 (9)	0.9730 (7)	0.9075 (10)	0.069 (3)
O3	0.6836 (10)	1.0854 (8)	0.7850 (10)	0.071 (3)
O4	0.7937 (12)	1.0687 (8)	0.9874 (10)	0.086 (4)
C1	0.6226 (11)	0.7895 (8)	0.6574 (11)	0.037 (3)
C2	0.6002 (10)	0.8714 (8)	0.5878 (10)	0.034 (3)
H2	0.6373	0.9281	0.6181	0.041*
C3	0.5204 (11)	0.8671 (9)	0.4706 (11)	0.041 (3)
H3	0.5026	0.9222	0.4241	0.049*
C4	0.4688 (12)	0.7846 (8)	0.4240 (12)	0.042 (3)
H4	0.4163	0.7830	0.3459	0.051*
C5	0.4949 (11)	0.7005 (9)	0.4943 (11)	0.037 (3)

H5	0.4603	0.6435	0.4617	0.045*
C6	0.5695 (10)	0.7026 (9)	0.6080 (10)	0.033 (3)
C7	0.5996 (10)	0.6133 (8)	0.6827 (9)	0.031 (3)
H7A	0.5489	0.6068	0.7335	0.037*
H7B	0.5881	0.5586	0.6292	0.037*
C8	0.8978 (11)	0.7594 (8)	0.7776 (11)	0.030 (3)
C9	0.9896 (11)	0.8252 (10)	0.7942 (12)	0.047 (3)
H9	0.9914	0.8787	0.8421	0.056*
C10	1.0741 (11)	0.8122 (10)	0.7424 (14)	0.051 (4)
H10	1.1337	0.8565	0.7542	0.062*
C11	1.0723 (14)	0.7321 (10)	0.6709 (16)	0.055 (4)
H11	1.1291	0.7246	0.6321	0.066*
C12	0.9893 (10)	0.6651 (9)	0.6570 (11)	0.038 (3)
H12	0.9919	0.6101	0.6132	0.045*
C13	0.8986 (9)	0.6785 (8)	0.7091 (9)	0.030 (2)
C14	0.8015 (10)	0.6065 (8)	0.6812 (10)	0.033 (3)
H14A	0.8350	0.5433	0.6923	0.040*
H14B	0.7555	0.6128	0.5965	0.040*
C15	0.7472 (9)	0.5399 (9)	0.8534 (10)	0.033 (3)
C16	0.6852 (11)	0.4552 (9)	0.8367 (12)	0.041 (3)
H16	0.6240	0.4455	0.7659	0.049*
C17	0.7139 (13)	0.3865 (10)	0.9238 (14)	0.053 (4)
H17	0.6723	0.3298	0.9117	0.064*
C18	0.8062 (14)	0.4002 (11)	1.0325 (14)	0.058 (4)
H18	0.8251	0.3532	1.0919	0.069*
C19	0.8671 (14)	0.4831 (12)	1.0489 (13)	0.064 (4)
H19	0.9281	0.4920	1.1202	0.077*
C20	0.8408 (11)	0.5534 (11)	0.9634 (11)	0.050 (3)
H20	0.8833	0.6097	0.9763	0.060*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Bi	0.0313 (3)	0.0438 (3)	0.0201 (3)	0.00040 (18)	0.0062 (2)	-0.00470 (18)
Cl1	0.0407 (17)	0.0414 (16)	0.0248 (14)	0.0024 (13)	0.0001 (12)	0.0037 (13)
N1	0.018 (5)	0.070 (7)	0.007 (4)	0.002 (4)	-0.001 (4)	0.001 (4)
O1	0.047 (6)	0.064 (7)	0.079 (8)	0.005 (5)	0.020 (5)	-0.009 (6)
O2	0.054 (6)	0.076 (7)	0.086 (8)	0.001 (5)	0.036 (6)	0.026 (6)
O3	0.082 (8)	0.065 (7)	0.059 (7)	0.000 (6)	0.008 (6)	0.034 (6)
O4	0.122 (11)	0.076 (8)	0.035 (6)	-0.017 (7)	-0.015 (6)	-0.011 (6)
C1	0.038 (7)	0.045 (7)	0.033 (7)	0.005 (5)	0.016 (6)	-0.005 (6)
C2	0.041 (7)	0.038 (6)	0.025 (6)	0.005 (5)	0.012 (5)	0.003 (5)
C3	0.043 (7)	0.053 (8)	0.032 (7)	0.013 (6)	0.019 (6)	0.013 (6)
C4	0.044 (8)	0.049 (8)	0.030 (7)	0.006 (6)	0.006 (6)	-0.004 (6)
C5	0.034 (7)	0.050 (8)	0.022 (6)	0.002 (5)	-0.002 (5)	-0.008 (5)
C6	0.021 (6)	0.054 (7)	0.025 (6)	0.000 (5)	0.006 (5)	0.007 (5)
C7	0.033 (6)	0.036 (6)	0.020 (5)	-0.006 (5)	0.003 (5)	0.004 (5)
C8	0.030 (7)	0.038 (6)	0.027 (6)	0.001 (5)	0.015 (5)	0.003 (5)

C9	0.045 (8)	0.045 (8)	0.048 (8)	-0.013 (6)	0.010 (6)	-0.015 (7)
C10	0.031 (7)	0.058 (8)	0.066 (9)	-0.013 (6)	0.015 (7)	0.005 (8)
C11	0.051 (10)	0.066 (10)	0.058 (10)	0.007 (7)	0.030 (8)	0.012 (8)
C12	0.035 (7)	0.046 (7)	0.034 (6)	0.004 (5)	0.012 (5)	-0.003 (6)
C13	0.031 (6)	0.039 (6)	0.020 (5)	0.002 (5)	0.006 (5)	0.003 (5)
C14	0.034 (6)	0.042 (7)	0.026 (6)	-0.001 (5)	0.014 (5)	-0.003 (5)
C15	0.036 (7)	0.036 (7)	0.030 (6)	0.004 (5)	0.015 (5)	0.006 (5)
C16	0.038 (7)	0.045 (8)	0.039 (7)	0.005 (6)	0.010 (6)	0.010 (6)
C17	0.059 (9)	0.045 (8)	0.063 (9)	0.005 (7)	0.029 (8)	0.020 (7)
C18	0.066 (10)	0.063 (10)	0.050 (9)	0.020 (8)	0.026 (8)	0.024 (8)
C19	0.061 (10)	0.083 (12)	0.041 (8)	0.011 (9)	0.005 (7)	0.018 (8)
C20	0.042 (8)	0.069 (10)	0.027 (7)	0.000 (7)	-0.007 (6)	0.004 (7)

Geometric parameters (Å, °)

Bi—N1	2.387 (10)	C8—C13	1.387 (17)
Bi—O1	2.546 (10)	C8—C9	1.411 (17)
Bi—C1	2.245 (13)	C9—C10	1.340 (18)
Bi—C8	2.204 (12)	C9—H9	0.9300
C11—O3	1.407 (10)	C10—C11	1.39 (2)
C11—O4	1.413 (10)	C10—H10	0.9300
C11—O2	1.434 (10)	C11—C12	1.349 (19)
C11—O1	1.448 (10)	C11—H11	0.9300
N1—C7	1.484 (13)	C12—C13	1.410 (16)
N1—C15	1.490 (15)	C12—H12	0.9300
N1—C14	1.514 (12)	C13—C14	1.508 (15)
C1—C2	1.383 (16)	C14—H14A	0.9700
C1—C6	1.418 (17)	C14—H14B	0.9700
C2—C3	1.402 (17)	C15—C16	1.389 (18)
C2—H2	0.9300	C15—C20	1.432 (16)
C3—C4	1.351 (18)	C16—C17	1.360 (18)
C3—H3	0.9300	C16—H16	0.9300
C4—C5	1.414 (18)	C17—C18	1.41 (2)
C4—H4	0.9300	C17—H17	0.9300
C5—C6	1.347 (16)	C18—C19	1.36 (2)
C5—H5	0.9300	C18—H18	0.9300
C6—C7	1.504 (16)	C19—C20	1.365 (19)
C7—H7A	0.9700	C19—H19	0.9300
C7—H7B	0.9700	C20—H20	0.9300
C8—Bi—C1	92.5 (5)	C13—C8—C9	118.4 (11)
C8—Bi—N1	77.4 (4)	C13—C8—Bi	115.1 (8)
C1—Bi—N1	74.6 (4)	C9—C8—Bi	126.5 (9)
C8—Bi—O1	83.1 (4)	C10—C9—C8	121.5 (13)
C1—Bi—O1	89.4 (4)	C10—C9—H9	119.3
N1—Bi—O1	154.0 (3)	C8—C9—H9	119.3
O3—C11—O4	110.7 (7)	C9—C10—C11	119.7 (13)
O3—C11—O2	111.0 (7)	C9—C10—H10	120.2

O4—C11—O2	110.9 (7)	C11—C10—H10	120.2
O3—C11—O1	108.6 (7)	C12—C11—C10	120.9 (13)
O4—C11—O1	108.6 (8)	C12—C11—H11	119.5
O2—C11—O1	106.9 (6)	C10—C11—H11	119.5
C7—N1—C15	110.7 (9)	C11—C12—C13	120.0 (12)
C7—N1—C14	109.2 (8)	C11—C12—H12	120.0
C15—N1—C14	109.6 (8)	C13—C12—H12	120.0
C7—N1—Bi	104.8 (7)	C8—C13—C12	119.4 (11)
C15—N1—Bi	113.7 (6)	C8—C13—C14	122.3 (10)
C14—N1—Bi	108.6 (7)	C12—C13—C14	118.2 (10)
C11—O1—Bi	122.3 (6)	C13—C14—N1	113.3 (9)
C2—C1—C6	120.1 (12)	C13—C14—H14A	108.9
C2—C1—Bi	127.1 (9)	N1—C14—H14A	108.9
C6—C1—Bi	112.7 (8)	C13—C14—H14B	108.9
C1—C2—C3	118.6 (12)	N1—C14—H14B	108.9
C1—C2—H2	120.7	H14A—C14—H14B	107.7
C3—C2—H2	120.7	C16—C15—C20	119.0 (11)
C4—C3—C2	121.3 (12)	C16—C15—N1	123.0 (10)
C4—C3—H3	119.4	C20—C15—N1	118.0 (11)
C2—C3—H3	119.4	C17—C16—C15	120.1 (13)
C3—C4—C5	119.8 (13)	C17—C16—H16	119.9
C3—C4—H4	120.1	C15—C16—H16	119.9
C5—C4—H4	120.1	C16—C17—C18	120.8 (14)
C6—C5—C4	120.4 (12)	C16—C17—H17	119.6
C6—C5—H5	119.8	C18—C17—H17	119.6
C4—C5—H5	119.8	C19—C18—C17	119.2 (13)
C5—C6—C1	119.8 (12)	C19—C18—H18	120.4
C5—C6—C7	120.9 (11)	C17—C18—H18	120.4
C1—C6—C7	119.3 (11)	C18—C19—C20	121.6 (15)
N1—C7—C6	109.9 (9)	C18—C19—H19	119.2
N1—C7—H7A	109.7	C20—C19—H19	119.2
C6—C7—H7A	109.7	C19—C20—C15	119.4 (14)
N1—C7—H7B	109.7	C19—C20—H20	120.3
C6—C7—H7B	109.7	C15—C20—H20	120.3
H7A—C7—H7B	108.2		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4 \cdots O3 ⁱ	0.93	2.46	3.137 (17)	130
C14—H14B \cdots O2 ⁱⁱ	0.97	2.55	3.398 (16)	146

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x, -y+3/2, z-1/2$.