

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4,4'-Bipyridine–pyridine-3,5-dicarboxylic acid (3/4)

 Sheng-Bo Liu,^a Chao Xu,^a Taike Duan,^a Qun Chen^b and Qian-Feng Zhang^{a,b*}
^aInstitute of Molecular Engineering and Applied Chemistry, Anhui University of Technology, Ma'anshan, Anhui 243002, People's Republic of China, and

^bDepartment of Applied Chemistry, School of Petrochemical Engineering, Changzhou University, Jiangsu 213164, People's Republic of China

Correspondence e-mail: zhangqf@cczu.edu.cn


Received 20 June 2011; accepted 7 July 2011

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.118; data-to-parameter ratio = 15.6.

In the title compound, $3\text{C}_{10}\text{H}_8\text{N}_2 \cdot 4\text{C}_7\text{H}_5\text{NO}_4$, the asymmetric unit contains two molecules of pyridine-3,5-dicarboxylic acid and one molecule of 4,4'-bipyridine in general positions together with one molecule of 4,4'-bipyridine lying across a centre of inversion, thus giving a 4:3 molar ratio of pyridine-3,5-dicarboxylic acid to 4,4'-bipyridine. The dihedral angle between the bipyridine rings on general positions is $21.2(2)^\circ$. These molecular units are linked by $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds forming an extended two-dimensional framework in the crystal.

Related literature

For structures and properties of self-assembled supermolecular compounds, see: Lehn (1990). For hydrogen bonds and $\pi-\pi$ stacking interactions in supermolecular compounds, see: Roesky & Andruh (2003). For related structures, see: Soleimannejad *et al.* (2009); Jiang *et al.* (2010).



Experimental

Crystal data

 $3\text{C}_{10}\text{H}_8\text{N}_2 \cdot 4\text{C}_7\text{H}_5\text{NO}_4$
 $M_r = 1137.04$
 Monoclinic, $P2_1/n$
 $a = 13.8461(4)$ Å
 $b = 11.0564(3)$ Å
 $c = 18.1060(4)$ Å
 $\beta = 110.511(1)^\circ$
 $V = 2596.09(12)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 296$ K
 $0.14 \times 0.12 \times 0.05$ mm

Data collection

 Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1997)
 $T_{\min} = 0.985$, $T_{\max} = 0.995$
 24265 measured reflections
 5918 independent reflections
 4086 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.118$
 $S = 1.03$
 5918 reflections

 379 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O2}-\text{H2} \cdots \text{N5}^i$	0.82	1.80	2.5928 (17)	162
$\text{O4}-\text{H4} \cdots \text{N4}^{ii}$	0.82	1.79	2.6037 (18)	172
$\text{O5}-\text{H5} \cdots \text{N3}^{iii}$	0.82	1.78	2.5956 (17)	173
$\text{O7}-\text{H7} \cdots \text{N2}^{ii}$	0.82	1.83	2.5862 (16)	152

 Symmetry codes: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 2, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This project was supported by the Program for New Century Excellent Talents in Universities of China (NCET-08-0618).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2358).

References

- Bruker (2005). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Jiang, X.-R., Wang, X.-J. & Feng, Y.-L. (2010). *Acta Cryst.* E66, o3308.
 Lehn, M.-L. (1990). *Angew. Chem. Int. Ed.* 29, 1304–1319.
 Roesky, H. W. & Andruh, M. (2003). *Coord. Chem. Rev.* 236, 91–119.
 Sheldrick, G. M. (1997). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* A64, 112–122.
 Soleimannejad, J., Aghabozorg, H., Morsali, A., Hemmati, F. & Manteghi, F. (2009). *Acta Cryst.* E65, o153.

supporting information

Acta Cryst. (2011). E67, o2033 [doi:10.1107/S160053681102719X]

4,4'-Bipyridine–pyridine-3,5-dicarboxylic acid (3/4)

Sheng-Bo Liu, Chao Xu, Taike Duan, Qun Chen and Qian-Feng Zhang

S1. Comment

In the past decades, the field of self-assembled supermolecular chemistry attracted increasing concerns due to their interesting structures and unique properties (Lehn, 1990). Hydrogen-bonds and pi—pi stacking are the most two important interactions in supermolecular compounds (Roesky & Andruh, 2003). As well known, many organic molecules with N- and O-donors such as bipyridine and pyridine-carboxylate acid have been widely employed for constructing supermolecular complexes. In this paper, we report the hydrothermal synthesis and co-crystal structure of a two-dimensional framework of 4,4'-bipyridine (4,4'-bipy) and pyridine-3,5-dicarboxylic acid (3,5-pydcH₂) assembled by the intermolecular hydrogen-bonding interactions. In the title compound, the asymmetric unit is formed by two molecules of pyridine-3,5-dicarboxylic acid and one molecule of 4,4'-bipyridine lying in general positions, together with one molecule of 4,4'-bipyridine lying across a centre of inversion, thus giving a 4:3 molar ratio of pyridine-3,5-dicarboxylic acid to 4,4'-bipyridine. These molecular units are linked by O—H...N hydrogen-bonds forming an extended two-dimensional framework, Table 1, Fig.2. The geometric parameters are in normal range.

S2. Experimental

4, 4'-bipyridine (31.2 mg, 0.2 mmol) and 3, 5-pyridinedicarboxylic acid (50.1 mg, 0.3 mmol) were mixed in 2.5 ml water solution and the mixture was stirred for a while. The solution was transferred to a 23 ml Teflon-lined stainless autoclave. The vessel was sealed and heated to 110 C° for 2 days. The autoclave was cooled to ambient temperature. Colorless flake crystals of title co-crystal compound were obtained and air dried. Yield: 43.2 mg, *ca* 67%. Anal. Calcd. for C₂₉H₂₁O₈N₅: C, 61.4; H, 3.71; N, 12.3%. Found: C, 61.1; H, 3.65; N, 12.1%.

S3. Refinement

H atoms were positioned geometrically and refined using a riding model C—H = 0.93 Å and O—H = 0.82 Å with $U_{\text{iso}}(\text{H}) = 1.2$ and 1.5 time for $U_{\text{eq}}(\text{C})$ and hydroxyl groups, respectively.




Figure 1

The structure of the title co-crystal compound, showing the atom-numbering scheme and displacement ellipsoids at the 50% probability level.




Figure 2

Packing diagram of the title compound viewed along *b*-axis showing hydrogen bonds as dash lines.

4,4'-Bipyridine–pyridine-3,5-dicarboxylic acid (3/4)

Crystal data

$3C_{10}H_8N_2 \cdot 4C_7H_5NO_4$

$M_r = 1137.04$

Monoclinic, $P2_1/n$

$a = 13.8461 (4) \text{ \AA}$

$b = 11.0564 (3) \text{ \AA}$

$c = 18.1060 (4) \text{ \AA}$

$\beta = 110.511 (1)^\circ$
 $V = 2596.09 (12) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 1180$
 $D_x = 1.455 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6026 reflections
 $\theta = 2.3\text{--}27.4^\circ$
 $\mu = 0.11 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
 Bar, colourless
 $0.14 \times 0.12 \times 0.05 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 phi and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1997)
 $T_{\min} = 0.985, T_{\max} = 0.995$

24265 measured reflections
 5918 independent reflections
 4086 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\max} = 27.5^\circ, \theta_{\min} = 1.6^\circ$
 $h = -17 \rightarrow 17$
 $k = -14 \rightarrow 14$
 $l = -19 \rightarrow 23$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.118$
 $S = 1.03$
 5918 reflections
 379 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.058P)^2 + 0.3037P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.19 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.22 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.89232 (10)	0.36614 (11)	1.13048 (7)	0.0566 (4)
O2	0.91840 (11)	0.19752 (11)	1.20316 (7)	0.0556 (4)
H2	0.9475	0.2439	1.2395	0.083*
O3	0.74670 (12)	0.28184 (12)	0.82900 (7)	0.0623 (4)
O4	0.71741 (11)	0.08686 (12)	0.79668 (7)	0.0619 (4)
H4	0.7058	0.1124	0.7519	0.093*
O5	0.87147 (10)	0.21216 (11)	0.58878 (6)	0.0513 (3)
H5	0.8940	0.2100	0.6371	0.077*
O6	0.91991 (11)	0.01904 (12)	0.60066 (7)	0.0624 (4)
O7	0.79322 (9)	-0.10426 (10)	0.23327 (6)	0.0469 (3)

H7	0.7969	-0.1689	0.2123	0.070*
O8	0.84624 (11)	-0.21541 (11)	0.34327 (7)	0.0577 (4)
N1	0.75488 (11)	-0.01589 (13)	1.02110 (8)	0.0448 (4)
N2	0.76208 (10)	0.19983 (11)	0.34879 (7)	0.0357 (3)
N3	1.06273 (12)	0.77515 (15)	0.25843 (8)	0.0504 (4)
N4	0.83365 (12)	0.65049 (16)	-0.14945 (8)	0.0520 (4)
N5	0.51891 (10)	0.82688 (12)	0.16936 (7)	0.0391 (3)
C1	0.88290 (13)	0.25840 (16)	1.13706 (9)	0.0403 (4)
C2	0.82884 (12)	0.17824 (14)	1.06824 (9)	0.0360 (4)
C3	0.79903 (13)	0.06220 (15)	1.07946 (9)	0.0412 (4)
H3	0.8105	0.0371	1.1309	0.049*
C4	0.74003 (12)	0.02231 (15)	0.94810 (9)	0.0399 (4)
H4A	0.7118	-0.0317	0.9067	0.048*
C5	0.76425 (12)	0.13790 (15)	0.93021 (9)	0.0365 (4)
C6	0.80869 (12)	0.21694 (15)	0.99183 (9)	0.0387 (4)
H6	0.8249	0.2955	0.9819	0.046*
C7	0.74147 (13)	0.17741 (17)	0.84653 (9)	0.0435 (4)
C8	0.88248 (12)	0.10611 (15)	0.56092 (9)	0.0387 (4)
C9	0.84316 (11)	0.10225 (13)	0.47271 (8)	0.0326 (3)
C10	0.79689 (12)	0.20178 (14)	0.42774 (9)	0.0341 (3)
H10	0.7897	0.2724	0.4533	0.041*
C11	0.77338 (12)	0.09854 (14)	0.31231 (9)	0.0351 (4)
H11	0.7507	0.0980	0.2575	0.042*
C12	0.81732 (11)	-0.00551 (13)	0.35251 (8)	0.0333 (3)
C13	0.85226 (12)	-0.00289 (14)	0.43401 (9)	0.0353 (4)
H13	0.8818	-0.0715	0.4628	0.042*
C14	0.82183 (12)	-0.11939 (14)	0.30945 (9)	0.0365 (4)
C15	1.05291 (14)	0.66299 (18)	0.23018 (10)	0.0517 (5)
H15	1.0778	0.5997	0.2656	0.062*
C16	1.00790 (13)	0.63556 (17)	0.15141 (10)	0.0468 (4)
H16	1.0007	0.5554	0.1348	0.056*
C17	0.97328 (12)	0.72881 (15)	0.09689 (9)	0.0377 (4)
C18	0.92417 (12)	0.70293 (16)	0.01128 (9)	0.0380 (4)
C19	0.88180 (13)	0.58973 (16)	-0.01519 (9)	0.0444 (4)
H19	0.8830	0.5294	0.0209	0.053*
C20	0.83816 (14)	0.56738 (18)	-0.09498 (10)	0.0501 (5)
H20	0.8107	0.4911	-0.1115	0.060*
C21	0.87225 (15)	0.75957 (19)	-0.12476 (10)	0.0557 (5)
H21	0.8684	0.8187	-0.1622	0.067*
C22	0.91764 (14)	0.78877 (17)	-0.04623 (10)	0.0484 (4)
H22	0.9439	0.8660	-0.0317	0.058*
C23	0.98663 (14)	0.84544 (17)	0.12672 (10)	0.0505 (5)
H23	0.9666	0.9110	0.0926	0.061*
C24	1.02949 (15)	0.86454 (18)	0.20685 (11)	0.0553 (5)
H24	1.0354	0.9435	0.2256	0.066*
C25	0.48131 (13)	0.77880 (15)	0.09709 (9)	0.0407 (4)
H25	0.4599	0.6985	0.0919	0.049*
C26	0.47296 (12)	0.84300 (14)	0.02999 (9)	0.0376 (4)

H26	0.4469	0.8057	-0.0191	0.045*
C27	0.50350 (11)	0.96381 (14)	0.03536 (8)	0.0323 (3)
C28	0.54147 (13)	1.01290 (15)	0.11086 (9)	0.0438 (4)
H28	0.5624	1.0933	0.1180	0.053*
C29	0.54790 (13)	0.94206 (16)	0.17488 (9)	0.0444 (4)
H29	0.5740	0.9767	0.2248	0.053*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0855 (9)	0.0374 (7)	0.0412 (7)	-0.0112 (6)	0.0149 (6)	-0.0045 (6)
O2	0.0860 (9)	0.0441 (7)	0.0262 (6)	-0.0050 (6)	0.0065 (6)	-0.0053 (5)
O3	0.0973 (11)	0.0522 (9)	0.0348 (7)	-0.0113 (7)	0.0198 (7)	0.0069 (6)
O4	0.0973 (10)	0.0599 (9)	0.0235 (6)	-0.0126 (7)	0.0148 (6)	-0.0025 (6)
O5	0.0793 (9)	0.0441 (7)	0.0249 (6)	0.0074 (6)	0.0109 (6)	-0.0062 (5)
O6	0.1012 (10)	0.0483 (8)	0.0291 (6)	0.0213 (7)	0.0123 (6)	0.0055 (6)
O7	0.0818 (9)	0.0287 (6)	0.0277 (6)	-0.0007 (5)	0.0160 (6)	-0.0037 (5)
O8	0.0974 (10)	0.0319 (7)	0.0366 (7)	0.0155 (6)	0.0147 (7)	0.0028 (5)
N1	0.0630 (9)	0.0356 (8)	0.0315 (8)	-0.0053 (6)	0.0110 (7)	-0.0003 (6)
N2	0.0488 (8)	0.0287 (7)	0.0277 (7)	0.0007 (6)	0.0111 (6)	0.0008 (5)
N3	0.0596 (9)	0.0583 (10)	0.0293 (8)	0.0016 (8)	0.0105 (7)	-0.0065 (7)
N4	0.0595 (9)	0.0672 (11)	0.0277 (8)	0.0081 (8)	0.0129 (7)	-0.0009 (8)
N5	0.0472 (8)	0.0396 (8)	0.0284 (7)	0.0025 (6)	0.0104 (6)	0.0056 (6)
C1	0.0531 (10)	0.0393 (10)	0.0283 (8)	-0.0024 (8)	0.0141 (7)	-0.0041 (7)
C2	0.0427 (9)	0.0362 (9)	0.0284 (8)	0.0006 (7)	0.0114 (7)	-0.0014 (7)
C3	0.0583 (10)	0.0381 (9)	0.0255 (8)	0.0010 (8)	0.0125 (7)	0.0009 (7)
C4	0.0486 (9)	0.0390 (9)	0.0286 (8)	-0.0021 (7)	0.0092 (7)	-0.0044 (7)
C5	0.0425 (9)	0.0409 (9)	0.0250 (8)	-0.0014 (7)	0.0104 (7)	-0.0001 (7)
C6	0.0487 (9)	0.0363 (9)	0.0313 (9)	-0.0033 (7)	0.0141 (7)	0.0013 (7)
C7	0.0491 (10)	0.0525 (11)	0.0278 (9)	-0.0062 (8)	0.0119 (7)	0.0005 (8)
C8	0.0488 (9)	0.0392 (10)	0.0271 (8)	0.0020 (7)	0.0121 (7)	-0.0024 (7)
C9	0.0404 (8)	0.0315 (8)	0.0256 (8)	0.0001 (6)	0.0112 (6)	-0.0005 (6)
C10	0.0441 (9)	0.0297 (8)	0.0288 (8)	-0.0016 (7)	0.0131 (7)	-0.0032 (6)
C11	0.0473 (9)	0.0323 (8)	0.0241 (8)	-0.0019 (7)	0.0106 (6)	-0.0005 (6)
C12	0.0420 (8)	0.0290 (8)	0.0278 (8)	-0.0015 (6)	0.0110 (6)	-0.0020 (6)
C13	0.0464 (9)	0.0305 (8)	0.0267 (8)	0.0030 (7)	0.0100 (7)	0.0022 (6)
C14	0.0489 (9)	0.0304 (9)	0.0284 (8)	0.0010 (7)	0.0113 (7)	-0.0006 (6)
C15	0.0652 (12)	0.0530 (12)	0.0299 (9)	0.0009 (9)	0.0080 (8)	0.0027 (8)
C16	0.0588 (11)	0.0435 (10)	0.0340 (9)	-0.0013 (8)	0.0112 (8)	-0.0023 (8)
C17	0.0393 (8)	0.0444 (10)	0.0292 (8)	0.0030 (7)	0.0116 (7)	-0.0020 (7)
C18	0.0388 (8)	0.0477 (10)	0.0279 (8)	0.0066 (7)	0.0119 (7)	0.0006 (7)
C19	0.0513 (10)	0.0498 (11)	0.0298 (9)	0.0020 (8)	0.0114 (7)	0.0000 (7)
C20	0.0586 (11)	0.0560 (12)	0.0330 (9)	0.0013 (9)	0.0129 (8)	-0.0066 (8)
C21	0.0693 (13)	0.0634 (13)	0.0351 (10)	0.0069 (10)	0.0190 (9)	0.0098 (9)
C22	0.0583 (11)	0.0500 (11)	0.0370 (10)	0.0004 (8)	0.0166 (8)	0.0020 (8)
C23	0.0645 (11)	0.0435 (11)	0.0372 (10)	0.0082 (9)	0.0097 (8)	0.0006 (8)
C24	0.0693 (12)	0.0499 (11)	0.0409 (11)	0.0036 (9)	0.0119 (9)	-0.0121 (9)
C25	0.0529 (10)	0.0320 (9)	0.0353 (9)	-0.0008 (7)	0.0130 (8)	0.0026 (7)

C26	0.0496 (9)	0.0319 (9)	0.0276 (8)	-0.0005 (7)	0.0088 (7)	-0.0018 (7)
C27	0.0361 (8)	0.0317 (8)	0.0268 (8)	0.0015 (6)	0.0080 (6)	0.0016 (6)
C28	0.0607 (11)	0.0354 (9)	0.0306 (9)	-0.0111 (8)	0.0104 (8)	-0.0024 (7)
C29	0.0583 (10)	0.0442 (10)	0.0258 (8)	-0.0061 (8)	0.0086 (7)	-0.0013 (7)

Geometric parameters (Å, °)

O1—C1	1.209 (2)	C9—C13	1.386 (2)
O2—C1	1.310 (2)	C10—H10	0.9300
O2—H2	0.8200	C11—C12	1.383 (2)
O3—C7	1.206 (2)	C11—H11	0.9300
O4—C7	1.311 (2)	C12—C13	1.383 (2)
O4—H4	0.8200	C12—C14	1.494 (2)
O5—C8	1.3063 (19)	C13—H13	0.9300
O5—H5	0.8200	C15—C16	1.375 (2)
O6—C8	1.2049 (19)	C15—H15	0.9300
O7—C14	1.3056 (18)	C16—C17	1.391 (2)
O7—H7	0.8200	C16—H16	0.9300
O8—C14	1.2129 (19)	C17—C23	1.385 (2)
N1—C4	1.333 (2)	C17—C18	1.486 (2)
N1—C3	1.335 (2)	C18—C22	1.388 (2)
N2—C11	1.3370 (19)	C18—C19	1.394 (2)
N2—C10	1.3391 (19)	C19—C20	1.379 (2)
N3—C24	1.326 (2)	C19—H19	0.9300
N3—C15	1.330 (2)	C20—H20	0.9300
N4—C21	1.331 (2)	C21—C22	1.376 (2)
N4—C20	1.333 (2)	C21—H21	0.9300
N5—C29	1.328 (2)	C22—H22	0.9300
N5—C25	1.337 (2)	C23—C24	1.378 (2)
C1—C2	1.498 (2)	C23—H23	0.9300
C2—C6	1.380 (2)	C24—H24	0.9300
C2—C3	1.384 (2)	C25—C26	1.377 (2)
C3—H3	0.9300	C25—H25	0.9300
C4—C5	1.388 (2)	C26—C27	1.394 (2)
C4—H4A	0.9300	C26—H26	0.9300
C5—C6	1.380 (2)	C27—C28	1.391 (2)
C5—C7	1.500 (2)	C27—C27 ⁱ	1.484 (3)
C6—H6	0.9300	C28—C29	1.376 (2)
C8—C9	1.496 (2)	C28—H28	0.9300
C9—C10	1.385 (2)	C29—H29	0.9300
C1—O2—H2	109.5	O8—C14—O7	124.97 (15)
C7—O4—H4	109.5	O8—C14—C12	121.98 (14)
C8—O5—H5	109.5	O7—C14—C12	113.00 (13)
C14—O7—H7	109.5	N3—C15—C16	123.54 (17)
C4—N1—C3	116.65 (14)	N3—C15—H15	118.2
C11—N2—C10	118.70 (13)	C16—C15—H15	118.2
C24—N3—C15	117.42 (15)	C15—C16—C17	119.38 (17)

C21—N4—C20	117.81 (15)	C15—C16—H16	120.3
C29—N5—C25	117.28 (13)	C17—C16—H16	120.3
O1—C1—O2	125.29 (15)	C23—C17—C16	116.58 (15)
O1—C1—C2	122.74 (15)	C23—C17—C18	122.40 (15)
O2—C1—C2	111.97 (14)	C16—C17—C18	121.02 (15)
C6—C2—C3	117.94 (15)	C22—C18—C19	116.55 (15)
C6—C2—C1	121.12 (15)	C22—C18—C17	122.44 (16)
C3—C2—C1	120.93 (14)	C19—C18—C17	121.01 (15)
N1—C3—C2	124.09 (15)	C20—C19—C18	119.88 (17)
N1—C3—H3	118.0	C20—C19—H19	120.1
C2—C3—H3	118.0	C18—C19—H19	120.1
N1—C4—C5	123.85 (15)	N4—C20—C19	122.77 (18)
N1—C4—H4A	118.1	N4—C20—H20	118.6
C5—C4—H4A	118.1	C19—C20—H20	118.6
C6—C5—C4	118.04 (14)	N4—C21—C22	122.94 (17)
C6—C5—C7	120.54 (15)	N4—C21—H21	118.5
C4—C5—C7	121.41 (14)	C22—C21—H21	118.5
C2—C6—C5	119.35 (15)	C21—C22—C18	120.04 (18)
C2—C6—H6	120.3	C21—C22—H22	120.0
C5—C6—H6	120.3	C18—C22—H22	120.0
O3—C7—O4	125.04 (15)	C24—C23—C17	120.18 (17)
O3—C7—C5	122.25 (16)	C24—C23—H23	119.9
O4—C7—C5	112.70 (15)	C17—C23—H23	119.9
O6—C8—O5	124.79 (15)	N3—C24—C23	122.84 (18)
O6—C8—C9	122.28 (15)	N3—C24—H24	118.6
O5—C8—C9	112.93 (14)	C23—C24—H24	118.6
C10—C9—C13	118.32 (14)	N5—C25—C26	122.83 (15)
C10—C9—C8	121.67 (14)	N5—C25—H25	118.6
C13—C9—C8	120.00 (13)	C26—C25—H25	118.6
N2—C10—C9	122.28 (14)	C25—C26—C27	120.19 (14)
N2—C10—H10	118.9	C25—C26—H26	119.9
C9—C10—H10	118.9	C27—C26—H26	119.9
N2—C11—C12	122.87 (14)	C28—C27—C26	116.31 (14)
N2—C11—H11	118.6	C28—C27—C27 ⁱ	121.69 (17)
C12—C11—H11	118.6	C26—C27—C27 ⁱ	122.00 (17)
C13—C12—C11	117.97 (14)	C29—C28—C27	119.75 (15)
C13—C12—C14	120.78 (14)	C29—C28—H28	120.1
C11—C12—C14	121.17 (13)	C27—C28—H28	120.1
C12—C13—C9	119.84 (14)	N5—C29—C28	123.63 (15)
C12—C13—H13	120.1	N5—C29—H29	118.2
C9—C13—H13	120.1	C28—C29—H29	118.2

Symmetry code: (i) $-x+1, -y+2, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots N5 ⁱⁱ	0.82	1.80	2.5928 (17)	162

O4—H4···N4 ⁱⁱⁱ	0.82	1.79	2.6037 (18)	172
O5—H5···N3 ^{iv}	0.82	1.78	2.5956 (17)	173
O7—H7···N2 ⁱⁱⁱ	0.82	1.83	2.5862 (16)	152

Symmetry codes: (ii) $-x+3/2, y-1/2, -z+3/2$; (iii) $-x+3/2, y-1/2, -z+1/2$; (iv) $-x+2, -y+1, -z+1$.