

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N*-tert-Butyl-2-methylpropanamide. Corrigendum**

**Kelly A. Kluge, Diana Fridlyand, Cora E. MacBeth* and
Kenneth I. Hardcastle**

Department of Chemistry, Emory University, 1515 Dickey Drive, Atlanta, GA 30322,
USA

Correspondence e-mail: cora.macbeth@emory.edu

Received 16 November 2011; accepted 16 November 2011

The name of one of the authors in the paper by Kluge *et al.* [*Acta Cryst.* (2011), **E67**, o2143] is corrected.

In the paper by Kluge *et al.* (2011), the name of the second author is given incorrectly. The correct name is given above.

References

Kluge, K. A., Fridlyand, D., MacBeth, C. A. & Hardcastle, K. I. (2011). *Acta Cryst.* **E67**, o2143.

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N*-tert-Butyl-2-methylpropanamide**

Kelly A. Kluge, Diana Fridlyand, Cora E. MacBeth* and Kenneth I. Hardcastle

Department of Chemistry, Emory University, 1515 Dickey Drive, Atlanta, GA 30322, USA

Correspondence e-mail: cora.macbeth@emory.edu


Received 21 June 2011; accepted 18 July 2011

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.035; wR factor = 0.106; data-to-parameter ratio = 14.7.

The title compound, $\text{C}_8\text{H}_{17}\text{NO}$, crystallizes with two independent molecules in the asymmetric unit. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonding is observed between neighboring molecules, forming continuous molecular chains along the c -axis direction.

Related literature

For the synthesis of the title compound, see: De Kimpe *et al.* (1978); Christensen *et al.* (1989); Yasuhara *et al.* (2000); Li *et al.* (2003). For its use as a ligand in Zr and Ti complexes, see: Li *et al.* (2003). For background to the coordination modes of carboxamides, see: Lee & Schafer (2007).



Experimental

Crystal data

 $\text{C}_8\text{H}_{17}\text{NO}$ $M_r = 143.23$ Monoclinic, $P2_1$ $a = 9.0378$ (6) Å $b = 11.3939$ (8) Å $c = 9.5390$ (6) Å $\beta = 106.133$ (3)° $V = 943.60$ (11) Å³ $Z = 4$ Cu $K\alpha$ radiation $\mu = 0.51$ mm⁻¹ $T = 173$ K

0.31 × 0.20 × 0.12 mm

Data collection

Bruker APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2008) $T_{\min} = 0.858$, $T_{\max} = 0.941$

6358 measured reflections

2662 independent reflections

2624 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.014$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.035$ $wR(F^2) = 0.106$ $S = 1.00$

2662 reflections

181 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.16$ e Å⁻³

Absolute structure: Flack (1983),

812 Friedel pairs

Flack parameter: 0.3 (2)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}$	0.88	2.03	2.8880 (16)	166
$\text{N2}-\text{H2A}\cdots\text{O1}^i$	0.88	2.10	2.9735 (16)	169

Symmetry code: (i) $x, y, z + 1$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We acknowledge the Emory University Center for X-ray Crystallography for assistance with data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2438).

References

- Bruker (2008). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Christensen, D. & Jorgensen, K. A. (1989). *J. Org. Chem.* **54**, 126–131.
- De Kimpe, N., Verhe, R., De Buyck, L., Chys, J. & Schamp, N. (1978). *Org. Prep. Proc. Intl.* **10**, 149–156.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Lee, A. & Schafer, L. L. (2007). *Eur. J. Inorg. Chem.* pp. 2243–2255.
- Li, C., Thomson, R. K., Gillon, B., Patrick, B. O. & Schafer, L. L. (2003). *Chem. Commun.* pp. 2462–2463.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yasuhara, T., Nagaoka, Y. & Tomioka, K. (2000). *J. Chem. Soc. Perkin Trans. 1*, pp. 2901–2902.

supporting information

Acta Cryst. (2011). E67, o2143 [doi:10.1107/S1600536811028947]

***N*-tert-Butyl-2-methylpropanamide**

Kelly A. Kluge, Diana Fridyland, Cora E. MacBeth and Kenneth I. Hardcastle

S1. Comment


Carboxamides can be deprotonated to form monoanionic amidate ligands. These species can coordinate to transition metal ions through a variety of different coordination modes, including monodentate and bidentate coordination modes, and therefore are coordinatively versatile ligands (Lee & Schafer, 2007). The ease of synthesis of carboxamides make them attractive ligands for a variety of transition metal mediated catalytic reactions, see: Li *et al.* (2003) and Lee & Schafer (2007). Although the synthesis of this compound has been previously described, its solid-state structure has not been reported. The two molecules (A and B) of *N*-tert-butyl-2-methylpropanamide (Fig. 1) are stabilized by intermolecular N—H···O hydrogen bonds (Table 1, Fig. 2).

S2. Experimental


The title molecule was synthesized using a modified literature procedure (Li *et al.*, 2003). Under a nitrogen atmosphere, a 100 ml round bottom flask was charged with 50 ml of dichloromethane, 4.31 ml (41.0 mmol) *tert*-butylamine, 8.55 ml (61.5 mmol) of triethylamine and a stir bar. The solution was cooled to 0 °C and 5.20 ml (49.2 mmol) of isobutyryl chloride was added dropwise. The solution was slowly warmed to room temperature overnight. The resulting pink solution was extracted three times with 50 ml of 0.10 M HCl. The organic layer was dried over anhydrous magnesium sulfate, filtered, and concentrated to dryness to yield the desired product in 60% yield. X-ray quality crystals were obtained by slowly evaporating a chloroform solution of the product. The spectroscopic data (NMR, IR, and ESI-MS) match well with the reported values (Li *et al.*, 2003).

S3. Refinement

The structures were solved using Direct Methods and difference Fourier techniques (*SHELXTL*, V6.12) (Sheldrick, 2008). Hydrogen atoms were added with the HFIX command. These were included in the final cycles of least squares refinement, with isotropic U^{ij} s that were determined by the riding model. All non-hydrogen atoms in the main residues were refined anisotropically, but residual solvent molecules in the unit cells were refined isotropically. Structure solution, refinement, and generation of publication materials were performed by using *SHELX*, V6.12 software.

**Figure 1**

Molecule A and Molecule B of *N*-*tert*-butyl-2-methylpropanamide. H atoms except H1A and H2A were omitted for clarity. Thermal ellipsoids are drawn at 50% probability.

**Figure 2**

Molecular packing and hydrogen bonding (dashed lines) network of *N*-*tert*-butyl-2-methylpropanamide viewed down the *b* axis.

N*-tert-butyl-2-methylpropanamideCrystal data*C₈H₁₇NO $M_r = 143.23$ Monoclinic, $P2_1$

Hall symbol: P 2yb

 $a = 9.0378$ (6) Å $b = 11.3939$ (8) Å $c = 9.5390$ (6) Å $\beta = 106.133$ (3)° $V = 943.60$ (11) Å³ $Z = 4$ $F(000) = 320$ $D_x = 1.008$ Mg m⁻³

Melting point: 393 K

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 4549 reflections

 $\theta = 4.8$ – 69.1 ° $\mu = 0.51$ mm⁻¹ $T = 173$ K

Block, colourless

 $0.31 \times 0.20 \times 0.12$ mm*Data collection*

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2008)

 $T_{\min} = 0.858$, $T_{\max} = 0.941$

6358 measured reflections

2662 independent reflections

2624 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.014$ $\theta_{\max} = 69.1$ °, $\theta_{\min} = 4.8$ ° $h = -10 \rightarrow 10$ $k = -13 \rightarrow 12$ $l = -11 \rightarrow 11$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.035$ $wR(F^2) = 0.106$ $S = 1.00$

2662 reflections

181 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.088P)^2 + 0.0504P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.19$ e Å⁻³ $\Delta\rho_{\min} = -0.16$ e Å⁻³Absolute structure: Flack (1983), 812 Friedel
pairs

Absolute structure parameter: 0.3 (2)

Special details

Geometry. All e.s.d.s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.30615 (14)	0.07943 (12)	0.55844 (13)	0.0364 (3)
H1A	0.2841	0.0677	0.6416	0.044*
N2	0.17705 (14)	0.08317 (12)	1.04274 (12)	0.0355 (3)
H2A	0.1857	0.1191	1.1263	0.043*
O1	0.23057 (13)	0.17773 (11)	0.34325 (11)	0.0394 (3)
O2	0.27727 (15)	0.06614 (13)	0.85203 (12)	0.0498 (3)

C1	0.1354 (2)	0.35161 (16)	0.5354 (2)	0.0541 (5)
H1B	0.0593	0.3932	0.5728	0.081*
H1C	0.1322	0.3819	0.4385	0.081*
H1D	0.2385	0.3638	0.6018	0.081*
C2	0.09866 (19)	0.22154 (14)	0.52483 (17)	0.0383 (4)
H2B	0.1043	0.1912	0.6246	0.046*
C3	-0.0618 (2)	0.19845 (19)	0.4252 (3)	0.0582 (5)
H3A	-0.1379	0.2398	0.4627	0.087*
H3B	-0.0829	0.1140	0.4220	0.087*
H3C	-0.0680	0.2266	0.3267	0.087*
C4	0.21920 (17)	0.15640 (14)	0.46654 (15)	0.0342 (3)
C5	0.43590 (17)	0.01194 (14)	0.53402 (15)	0.0350 (3)
C6	0.56316 (19)	0.09591 (16)	0.5211 (2)	0.0447 (4)
H6A	0.5246	0.1463	0.4354	0.067*
H6B	0.6517	0.0507	0.5105	0.067*
H6C	0.5948	0.1446	0.6090	0.067*
C7	0.3832 (2)	-0.06553 (16)	0.39919 (17)	0.0428 (4)
H7A	0.3015	-0.1183	0.4102	0.064*
H7B	0.4704	-0.1120	0.3879	0.064*
H7C	0.3437	-0.0162	0.3127	0.064*
C8	0.4953 (2)	-0.06704 (17)	0.66758 (18)	0.0466 (4)
H8A	0.4130	-0.1203	0.6759	0.070*
H8B	0.5277	-0.0185	0.7556	0.070*
H8C	0.5832	-0.1129	0.6568	0.070*
C9	0.4281 (2)	0.28811 (19)	0.9455 (3)	0.0580 (5)
H9A	0.5106	0.3416	0.9961	0.087*
H9B	0.3314	0.3318	0.9116	0.087*
H9C	0.4542	0.2529	0.8617	0.087*
C10	0.40976 (18)	0.19187 (15)	1.04972 (17)	0.0380 (3)
H10A	0.3831	0.2280	1.1351	0.046*
C11	0.5592 (2)	0.12282 (19)	1.1036 (2)	0.0519 (4)
H11A	0.6423	0.1759	1.1538	0.078*
H11B	0.5851	0.0865	1.0204	0.078*
H11C	0.5462	0.0616	1.1714	0.078*
C12	0.28088 (17)	0.10794 (14)	0.97174 (15)	0.0343 (3)
C13	0.04849 (19)	-0.00018 (18)	0.99122 (18)	0.0442 (4)
C14	-0.0612 (2)	0.0436 (2)	0.8479 (2)	0.0677 (6)
H14A	-0.0073	0.0448	0.7718	0.102*
H14B	-0.0963	0.1231	0.8618	0.102*
H14C	-0.1503	-0.0090	0.8183	0.102*
C15	0.1083 (3)	-0.1214 (2)	0.9736 (3)	0.0666 (6)
H15A	0.1785	-0.1470	1.0665	0.100*
H15B	0.1635	-0.1196	0.8985	0.100*
H15C	0.0216	-0.1762	0.9442	0.100*
C16	-0.0360 (3)	-0.0012 (3)	1.1101 (3)	0.0796 (8)
H16B	-0.1251	-0.0539	1.0810	0.119*
H16C	-0.0710	0.0783	1.1234	0.119*
H16A	0.0341	-0.0289	1.2020	0.119*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0423 (7)	0.0407 (7)	0.0305 (5)	0.0064 (6)	0.0173 (5)	0.0040 (5)
N2	0.0331 (6)	0.0460 (8)	0.0282 (5)	-0.0053 (5)	0.0099 (4)	-0.0068 (5)
O1	0.0445 (5)	0.0478 (7)	0.0295 (5)	0.0060 (5)	0.0159 (4)	0.0053 (5)
O2	0.0592 (7)	0.0643 (8)	0.0309 (5)	-0.0154 (6)	0.0208 (5)	-0.0115 (5)
C1	0.0598 (11)	0.0418 (10)	0.0689 (12)	-0.0012 (8)	0.0318 (9)	-0.0113 (9)
C2	0.0438 (8)	0.0397 (9)	0.0361 (7)	0.0048 (7)	0.0192 (6)	0.0048 (6)
C3	0.0375 (9)	0.0540 (11)	0.0866 (14)	0.0021 (8)	0.0229 (9)	-0.0120 (10)
C4	0.0373 (7)	0.0372 (8)	0.0306 (6)	-0.0009 (6)	0.0135 (5)	-0.0003 (6)
C5	0.0390 (7)	0.0344 (8)	0.0327 (7)	0.0047 (6)	0.0115 (6)	0.0030 (6)
C6	0.0389 (8)	0.0390 (9)	0.0580 (9)	0.0021 (7)	0.0168 (7)	0.0003 (8)
C7	0.0493 (9)	0.0380 (8)	0.0429 (8)	0.0040 (7)	0.0158 (7)	-0.0035 (7)
C8	0.0488 (9)	0.0495 (10)	0.0419 (8)	0.0126 (8)	0.0134 (6)	0.0090 (8)
C9	0.0492 (10)	0.0450 (10)	0.0796 (13)	-0.0033 (8)	0.0175 (9)	0.0128 (10)
C10	0.0388 (7)	0.0406 (8)	0.0378 (7)	-0.0043 (7)	0.0159 (6)	-0.0073 (6)
C11	0.0429 (9)	0.0549 (11)	0.0519 (9)	-0.0034 (8)	0.0031 (7)	-0.0001 (8)
C12	0.0379 (7)	0.0401 (8)	0.0263 (6)	0.0010 (6)	0.0111 (5)	0.0001 (6)
C13	0.0363 (7)	0.0559 (10)	0.0412 (8)	-0.0106 (7)	0.0120 (6)	-0.0078 (8)
C14	0.0457 (10)	0.0811 (15)	0.0627 (12)	-0.0095 (10)	-0.0075 (8)	-0.0083 (11)
C15	0.0647 (13)	0.0491 (11)	0.0838 (15)	-0.0171 (10)	0.0172 (10)	-0.0049 (10)
C16	0.0562 (12)	0.120 (2)	0.0741 (13)	-0.0411 (14)	0.0373 (10)	-0.0240 (15)

Geometric parameters (Å, °)

N1—C4	1.331 (2)	C7—H7C	0.9800
N1—C5	1.4739 (19)	C8—H8A	0.9800
N1—H1A	0.8800	C8—H8B	0.9800
N2—C12	1.3312 (19)	C8—H8C	0.9800
N2—C13	1.475 (2)	C9—C10	1.520 (3)
N2—H2A	0.8800	C9—H9A	0.9800
O1—C4	1.2328 (18)	C9—H9B	0.9800
O2—C12	1.2293 (19)	C9—H9C	0.9800
C1—C2	1.516 (2)	C10—C11	1.523 (2)
C1—H1B	0.9800	C10—C12	1.531 (2)
C1—H1C	0.9800	C10—H10A	1.0000
C1—H1D	0.9800	C11—H11A	0.9800
C2—C3	1.519 (2)	C11—H11B	0.9800
C2—C4	1.544 (2)	C11—H11C	0.9800
C2—H2B	1.0000	C13—C15	1.509 (3)
C3—H3A	0.9800	C13—C14	1.533 (3)
C3—H3B	0.9800	C13—C16	1.533 (3)
C3—H3C	0.9800	C14—H14A	0.9800
C5—C7	1.523 (2)	C14—H14B	0.9800
C5—C6	1.527 (2)	C14—H14C	0.9800
C5—C8	1.530 (2)	C15—H15A	0.9800
C6—H6A	0.9800	C15—H15B	0.9800

C6—H6B	0.9800	C15—H15C	0.9800
C6—H6C	0.9800	C16—H16B	0.9800
C7—H7A	0.9800	C16—H16C	0.9800
C7—H7B	0.9800	C16—H16A	0.9800
C4—N1—C5	126.13 (11)	C5—C8—H8C	109.5
C4—N1—H1A	116.9	H8A—C8—H8C	109.5
C5—N1—H1A	116.9	H8B—C8—H8C	109.5
C12—N2—C13	124.63 (13)	C10—C9—H9A	109.5
C12—N2—H2A	117.7	C10—C9—H9B	109.5
C13—N2—H2A	117.7	H9A—C9—H9B	109.5
C2—C1—H1B	109.5	C10—C9—H9C	109.5
C2—C1—H1C	109.5	H9A—C9—H9C	109.5
H1B—C1—H1C	109.5	H9B—C9—H9C	109.5
C2—C1—H1D	109.5	C9—C10—C11	110.23 (15)
H1B—C1—H1D	109.5	C9—C10—C12	109.85 (13)
H1C—C1—H1D	109.5	C11—C10—C12	108.96 (14)
C1—C2—C3	111.37 (16)	C9—C10—H10A	109.3
C1—C2—C4	109.28 (14)	C11—C10—H10A	109.3
C3—C2—C4	109.74 (14)	C12—C10—H10A	109.3
C1—C2—H2B	108.8	C10—C11—H11A	109.5
C3—C2—H2B	108.8	C10—C11—H11B	109.5
C4—C2—H2B	108.8	H11A—C11—H11B	109.5
C2—C3—H3A	109.5	C10—C11—H11C	109.5
C2—C3—H3B	109.5	H11A—C11—H11C	109.5
H3A—C3—H3B	109.5	H11B—C11—H11C	109.5
C2—C3—H3C	109.5	O2—C12—N2	123.37 (15)
H3A—C3—H3C	109.5	O2—C12—C10	120.91 (14)
H3B—C3—H3C	109.5	N2—C12—C10	115.72 (13)
O1—C4—N1	124.59 (14)	N2—C13—C15	110.67 (15)
O1—C4—C2	120.20 (14)	N2—C13—C14	110.00 (16)
N1—C4—C2	115.22 (12)	C15—C13—C14	111.19 (18)
N1—C5—C7	111.09 (12)	N2—C13—C16	105.46 (15)
N1—C5—C6	109.64 (12)	C15—C13—C16	110.0 (2)
C7—C5—C6	111.15 (14)	C14—C13—C16	109.32 (18)
N1—C5—C8	106.60 (12)	C13—C14—H14A	109.5
C7—C5—C8	108.50 (14)	C13—C14—H14B	109.5
C6—C5—C8	109.75 (13)	H14A—C14—H14B	109.5
C5—C6—H6A	109.5	C13—C14—H14C	109.5
C5—C6—H6B	109.5	H14A—C14—H14C	109.5
H6A—C6—H6B	109.5	H14B—C14—H14C	109.5
C5—C6—H6C	109.5	C13—C15—H15A	109.5
H6A—C6—H6C	109.5	C13—C15—H15B	109.5
H6B—C6—H6C	109.5	H15A—C15—H15B	109.5
C5—C7—H7A	109.5	C13—C15—H15C	109.5
C5—C7—H7B	109.5	H15A—C15—H15C	109.5
H7A—C7—H7B	109.5	H15B—C15—H15C	109.5
C5—C7—H7C	109.5	C13—C16—H16B	109.5

H7A—C7—H7C	109.5	C13—C16—H16C	109.5
H7B—C7—H7C	109.5	H16B—C16—H16C	109.5
C5—C8—H8A	109.5	C13—C16—H16A	109.5
C5—C8—H8B	109.5	H16B—C16—H16A	109.5
H8A—C8—H8B	109.5	H16C—C16—H16A	109.5
C5—N1—C4—O1	3.9 (3)	C13—N2—C12—O2	2.5 (3)
C5—N1—C4—C2	-175.58 (14)	C13—N2—C12—C10	-176.63 (15)
C1—C2—C4—O1	-63.0 (2)	C9—C10—C12—O2	49.6 (2)
C3—C2—C4—O1	59.4 (2)	C11—C10—C12—O2	-71.2 (2)
C1—C2—C4—N1	116.50 (17)	C9—C10—C12—N2	-131.26 (16)
C3—C2—C4—N1	-121.10 (17)	C11—C10—C12—N2	107.89 (17)
C4—N1—C5—C7	-60.1 (2)	C12—N2—C13—C15	59.7 (2)
C4—N1—C5—C6	63.11 (19)	C12—N2—C13—C14	-63.5 (2)
C4—N1—C5—C8	-178.15 (16)	C12—N2—C13—C16	178.7 (2)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1A...O2	0.88	2.03	2.8880 (16)	166
N2—H2A...O1 ⁱ	0.88	2.10	2.9735 (16)	169

Symmetry code: (i) *x, y, z+1*.