

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(Quinolin-8-yl)ferrocene-1-carboxamide

Xia Li* and Ling-Zhi Du

 Department of Chemistry and Chemical Engineering, Henan University of Urban Construction, Pingdingshan, Henan 467044, People's Republic of China
 Correspondence e-mail: lixia@hncj.edu.cn


Received 29 June 2011; accepted 3 July 2011

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.034; wR factor = 0.097; data-to-parameter ratio = 17.7.

In the title compound, $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{15}\text{H}_{11}\text{N}_2\text{O})]$, the cyclopentadienyl rings are essentially eclipsed, and the dihedral angle between the cyclopentadienyl ring planes is $0.632(10)^\circ$. The Fe atom is slightly closer to the substituted cyclopentadienyl ring, with an Fe-centroid distance of $1.6374(3)$ Å [$1.6494(3)$ Å for the unsubstituted ring]. The amide group is essentially coplanar with the substituted cyclopentadienyl ring, with an N—C(O)—C—C torsion angle of $2.3(3)^\circ$.

Related literature

For background to the chemical, stereochemical and electrochemical properties of ferrocene, see: Togni & Hayashi (1995). Ferrocene has been extensively incorporated into larger compounds in order to take advantage of these properties, see: Abd-El-Aziz & Manners (2007); Fang *et al.* (2001); Mata *et al.* (2001). For our research on ferrocenyl derivatives and their metal complexes, see: Li *et al.* (2008, 2009).



Experimental

Crystal data

$[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{15}\text{H}_{11}\text{N}_2\text{O})]$	$V = 3217.4(10)$ Å ³
$M_r = 356.20$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 10.1680(17)$ Å	$\mu = 0.95$ mm ⁻¹
$b = 12.133(2)$ Å	$T = 296$ K
$c = 26.079(4)$ Å	$0.46 \times 0.37 \times 0.25$ mm

Data collection

Rigaku Mercury CCD diffractometer	18658 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2000)	3940 independent reflections
$T_{\min} = 0.668$, $T_{\max} = 0.795$	2996 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	222 parameters
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\text{max}} = 0.26$ e Å ⁻³
3940 reflections	$\Delta\rho_{\text{min}} = -0.37$ e Å ⁻³

Data collection: *CrystalClear* (Rigaku, 2000); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We gratefully acknowledge financial support from the Foundation of Henan Educational Committee (2011B150001) and the Foundation of Henan University of Urban Construction (2010JYB007).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2441).

References

- Abd-El-Aziz, A. S. & Manners, I. (2007). In *Frontiers in Transition Metal-Containing Polymers*. Hoboken, NJ: Wiley.
- Fang, C. J., Duan, C. Y., Guo, D., He, C., Meng, Q. J., Wang, Z. M. & Yan, C. H. (2001). *Chem. Commun.* pp. 2540–2541.
- Li, X., Liu, W., Zhang, H. Y. & Wu, B. L. (2008). *J. Organomet. Chem.* **693**, 3295–3302.
- Li, X., Wu, B. L., Liu, W., Niu, C. Y., Niu, Y. Y. & Zhang, H. Y. (2009). *J. Coord. Chem.* **62**, 3142–3156.
- Mata, J. A., Peris, E., Asselberghs, I., Van Boxel, R. & Persoons, A. (2001). *New J. Chem.* **25**, 1043–1046.
- Rigaku (2000). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Togni, A. & Hayashi, T. (1995). In *Ferrocenes: Homogeneous Catalysis, Organic Synthesis, Materials Science*. New York: VCH Publishers.

supporting information

Acta Cryst. (2011). E67, m1113 [doi:10.1107/S1600536811026341]

N-(Quinolin-8-yl)ferrocene-1-carboxamide**Xia Li and Ling-Zhi Du****S1. Comment**

Due to its special chemical, stereochemical and electrochemical properties (Togni *et al.*, 1995), ferrocene has been extensively incorporated into larger compounds in order to take advantage of these properties (Mata *et al.*, 2001; Abd-El-Aziz *et al.*, 2007; Fang *et al.*, 2001). As a continuation of our research related to ferrocenyl derivatives and their metal complexes (Li *et al.*, 2008; Li *et al.*, 2009), herein we report the crystal structure of 8-(Ferrocenoylamino)quinoline.


The molecular structure of the title compound is composed of a ferrocenyl group and a quinolyl group joined by an organic amide spacer. The Fe—C bond distances within the ferrocene group are in the range of 2.0259 (19)–2.048 (2) Å for the substituted cyclopentadienyl (Cp) ring [C1—C5] and 2.026 (2)–2.039 (2) Å for the unsubstituted Cp ring [C6—C10]. The planar cyclopentadienyl rings of the ferrocenyl unit are nearly parallel to each other [the interplanar angle is 0.632 (10) °]. The Cp rings are essentially eclipsed and the Fe-centroid distances are 1.6374 (3) Å (*Cg*1) and 1.6494 (3) Å (*Cg*2) with *Cg*1 and *Cg*2 are the centroids of the [C1—C5] and [C6—C10] rings. The [*Cg*1—Fe1—*Cg*2] angle is 179.324 (17) °. The carbamoyl group is essentially coplanar with the substituted cyclopentadienyl ring with a deviation of 4.9 (2) °. The angle formed by the carbamoyl group and the quinolyl group system is 9.4 (3) °.

S2. Experimental

A solution of Chlorocarbonyl ferrocene (0.248 g, 1 mmol) in CH₂Cl₂ (20 ml) was added dropwise to a vigorously stirred solution of the 8-Aminoquinoline (0.144 g, 1 mmol) in CH₂Cl₂ (20 ml) containing pyridine (0.5 ml). The stirred reaction mixture was maintained at room temperature for 4 h. Removal of the solvent afforded the crude amide and the residue was recrystallized in dichloromethane/ether to give orange crystals 0.325 g. Yield 91.3%.

S3. Refinement

H atom bonded to N atom was located from difference Fourier maps and refined with a *DFIX* restraint of 0.86 (2) Å. Aromatic H atoms were positioned geometrically with C—H = 0.95 Å and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of the title compound, showing the atomic numbering and 30% probability displacement ellipsoids.

***N*-(Quinolin-8-yl)ferrocene-1-carboxamide**

Crystal data

[Fe(C₅H₅)(C₁₅H₁₁N₂O)]

$M_r = 356.20$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 10.1680$ (17) Å

$b = 12.133$ (2) Å

$c = 26.079$ (4) Å

$V = 3217.4$ (10) Å³

$Z = 8$

$F(000) = 1472$

$D_x = 1.471$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

$\theta = 1.6$ – 28.3°

$\mu = 0.95$ mm⁻¹

$T = 296$ K

Block, orange

$0.46 \times 0.37 \times 0.25$ mm

Data collection

Rigaku Mercury CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scan

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2000)

$T_{\min} = 0.668$, $T_{\max} = 0.795$

18658 measured reflections

3940 independent reflections

2996 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -13 \rightarrow 13$

$k = -12 \rightarrow 15$

$l = -34 \rightarrow 29$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.097$

$S = 1.01$

3940 reflections

222 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.P)^2 + 1.0596P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.26$ e Å⁻³

$\Delta\rho_{\min} = -0.37$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.08194 (2)	0.61371 (2)	0.324077 (9)	0.03985 (12)
O1	-0.14938 (18)	0.85691 (16)	0.34733 (7)	0.0757 (5)
N1	0.02468 (18)	0.87266 (15)	0.40153 (7)	0.0471 (4)
N2	0.19232 (16)	0.94139 (13)	0.47326 (6)	0.0433 (4)
C1	-0.04130 (18)	0.69359 (18)	0.37255 (7)	0.0432 (4)
C2	-0.1135 (2)	0.6191 (2)	0.34083 (8)	0.0509 (5)
H2	-0.1811	0.6384	0.3186	0.061*
C3	-0.0656 (2)	0.5119 (2)	0.34881 (8)	0.0580 (6)
H3	-0.0959	0.4483	0.3328	0.070*
C4	0.0368 (2)	0.5175 (2)	0.38554 (8)	0.0585 (6)
H4	0.0856	0.4581	0.3977	0.070*
C5	0.0521 (2)	0.62850 (18)	0.40057 (7)	0.0484 (5)
H5	0.1122	0.6549	0.4245	0.058*
C6	0.1656 (3)	0.7278 (2)	0.27759 (10)	0.0756 (8)
H6	0.1460	0.8027	0.2772	0.091*
C7	0.1037 (2)	0.6449 (3)	0.24815 (9)	0.0741 (8)
H7	0.0355	0.6550	0.2249	0.089*
C8	0.1644 (3)	0.5444 (3)	0.26050 (10)	0.0736 (8)
H8	0.1437	0.4758	0.2468	0.088*
C9	0.2604 (3)	0.5657 (3)	0.29680 (10)	0.0767 (8)
H9	0.3153	0.5133	0.3117	0.092*
C10	0.2614 (2)	0.6765 (3)	0.30727 (11)	0.0775 (8)
H10	0.3171	0.7115	0.3304	0.093*
C11	-0.06124 (18)	0.81347 (19)	0.37209 (7)	0.0475 (5)
C12	0.02430 (18)	0.98542 (16)	0.41223 (7)	0.0419 (4)
C13	-0.0553 (2)	1.0624 (2)	0.38937 (8)	0.0525 (5)
H13	-0.1144	1.0406	0.3641	0.063*
C14	-0.0483 (2)	1.1737 (2)	0.40382 (9)	0.0602 (6)
H14	-0.1021	1.2247	0.3875	0.072*
C15	0.0351 (2)	1.20882 (19)	0.44113 (9)	0.0583 (5)
H15	0.0374	1.2829	0.4503	0.070*
C16	0.1185 (2)	1.13215 (16)	0.46605 (7)	0.0437 (4)
C17	0.11420 (17)	1.02001 (15)	0.45129 (6)	0.0381 (4)
C18	0.2047 (2)	1.16091 (17)	0.50584 (8)	0.0506 (5)
H18	0.2107	1.2338	0.5166	0.061*

C19	0.2793 (2)	1.08233 (18)	0.52850 (8)	0.0512 (5)
H19	0.3351	1.1000	0.5555	0.061*
C20	0.2707 (2)	0.97401 (17)	0.51045 (7)	0.0489 (5)
H20	0.3241	0.9213	0.5258	0.059*
H1	0.078 (2)	0.8400 (19)	0.4191 (8)	0.046 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.03308 (17)	0.0496 (2)	0.03683 (16)	-0.00552 (11)	0.00144 (10)	-0.00378 (11)
O1	0.0693 (11)	0.0826 (12)	0.0752 (11)	0.0229 (10)	-0.0350 (9)	-0.0182 (9)
N1	0.0441 (9)	0.0508 (10)	0.0464 (9)	0.0082 (8)	-0.0110 (7)	-0.0072 (7)
N2	0.0460 (9)	0.0388 (9)	0.0451 (8)	0.0007 (7)	-0.0054 (7)	-0.0007 (7)
C1	0.0344 (9)	0.0619 (13)	0.0334 (8)	-0.0037 (9)	0.0001 (7)	-0.0077 (8)
C2	0.0341 (10)	0.0756 (16)	0.0431 (10)	-0.0137 (10)	0.0019 (8)	-0.0091 (10)
C3	0.0572 (13)	0.0613 (15)	0.0554 (12)	-0.0254 (11)	0.0076 (10)	-0.0071 (11)
C4	0.0674 (14)	0.0551 (13)	0.0530 (12)	-0.0080 (11)	0.0008 (10)	0.0106 (10)
C5	0.0501 (11)	0.0597 (13)	0.0356 (9)	-0.0065 (10)	-0.0038 (8)	0.0002 (9)
C6	0.0774 (18)	0.0681 (17)	0.0813 (17)	-0.0078 (14)	0.0356 (15)	0.0080 (14)
C7	0.0482 (13)	0.133 (3)	0.0411 (11)	-0.0043 (15)	0.0088 (9)	0.0110 (14)
C8	0.0768 (18)	0.0833 (19)	0.0608 (14)	-0.0106 (15)	0.0243 (13)	-0.0214 (13)
C9	0.0494 (14)	0.106 (2)	0.0749 (16)	0.0155 (15)	0.0118 (12)	-0.0030 (16)
C10	0.0445 (13)	0.113 (3)	0.0748 (16)	-0.0260 (15)	0.0118 (12)	-0.0170 (16)
C11	0.0408 (10)	0.0643 (14)	0.0374 (9)	0.0060 (9)	-0.0037 (8)	-0.0098 (9)
C12	0.0406 (10)	0.0467 (11)	0.0385 (9)	0.0054 (8)	0.0056 (7)	-0.0018 (8)
C13	0.0492 (12)	0.0625 (14)	0.0457 (10)	0.0110 (10)	-0.0004 (9)	0.0057 (10)
C14	0.0623 (14)	0.0544 (14)	0.0640 (13)	0.0192 (11)	0.0068 (11)	0.0163 (11)
C15	0.0645 (14)	0.0443 (12)	0.0660 (13)	0.0077 (11)	0.0078 (11)	0.0087 (10)
C16	0.0476 (10)	0.0360 (10)	0.0475 (10)	0.0009 (8)	0.0121 (8)	0.0050 (8)
C17	0.0390 (9)	0.0388 (10)	0.0366 (8)	0.0017 (8)	0.0064 (7)	0.0027 (7)
C18	0.0603 (13)	0.0368 (10)	0.0548 (11)	-0.0077 (9)	0.0087 (10)	-0.0059 (9)
C19	0.0599 (13)	0.0455 (11)	0.0483 (10)	-0.0104 (10)	-0.0063 (9)	-0.0018 (9)
C20	0.0562 (12)	0.0390 (10)	0.0515 (10)	-0.0009 (9)	-0.0102 (9)	0.0011 (8)

Geometric parameters (Å, °)

Fe1—C10	2.026 (2)	C6—C10	1.391 (4)
Fe1—C5	2.0258 (19)	C6—C7	1.413 (4)
Fe1—C1	2.0268 (19)	C6—H6	0.9300
Fe1—C6	2.028 (2)	C7—C8	1.404 (4)
Fe1—C7	2.028 (2)	C7—H7	0.9300
Fe1—C9	2.034 (2)	C8—C9	1.384 (4)
Fe1—C4	2.036 (2)	C8—H8	0.9300
Fe1—C2	2.036 (2)	C9—C10	1.372 (4)
Fe1—C8	2.039 (2)	C9—H9	0.9300
Fe1—C3	2.048 (2)	C10—H10	0.9300
O1—C11	1.224 (2)	C12—C13	1.372 (3)
N1—C11	1.367 (3)	C12—C17	1.432 (3)

N1—C12	1.396 (3)	C13—C14	1.405 (4)
N1—H1	0.81 (2)	C13—H13	0.9300
N2—C20	1.316 (2)	C14—C15	1.359 (3)
N2—C17	1.367 (2)	C14—H14	0.9300
C1—C2	1.429 (3)	C15—C16	1.417 (3)
C1—C5	1.435 (3)	C15—H15	0.9300
C1—C11	1.469 (3)	C16—C18	1.402 (3)
C2—C3	1.404 (3)	C16—C17	1.415 (3)
C2—H2	0.9300	C18—C19	1.354 (3)
C3—C4	1.417 (3)	C18—H18	0.9300
C3—H3	0.9300	C19—C20	1.399 (3)
C4—C5	1.412 (3)	C19—H19	0.9300
C4—H4	0.9300	C20—H20	0.9300
C5—H5	0.9300		
C10—Fe1—C5	108.35 (10)	C5—C4—H4	125.8
C10—Fe1—C1	120.82 (10)	C3—C4—H4	125.8
C5—Fe1—C1	41.47 (8)	Fe1—C4—H4	126.3
C10—Fe1—C6	40.16 (12)	C4—C5—C1	108.12 (18)
C5—Fe1—C6	126.24 (11)	C4—C5—Fe1	70.04 (12)
C1—Fe1—C6	107.81 (10)	C1—C5—Fe1	69.30 (10)
C10—Fe1—C7	67.67 (11)	C4—C5—H5	125.9
C5—Fe1—C7	164.01 (13)	C1—C5—H5	125.9
C1—Fe1—C7	125.97 (11)	Fe1—C5—H5	126.3
C6—Fe1—C7	40.79 (11)	C10—C6—C7	107.2 (3)
C10—Fe1—C9	39.50 (13)	C10—C6—Fe1	69.85 (15)
C5—Fe1—C9	120.23 (10)	C7—C6—Fe1	69.62 (15)
C1—Fe1—C9	155.04 (10)	C10—C6—H6	126.4
C6—Fe1—C9	67.18 (12)	C7—C6—H6	126.4
C7—Fe1—C9	67.32 (11)	Fe1—C6—H6	125.7
C10—Fe1—C4	126.11 (12)	C8—C7—C6	107.4 (2)
C5—Fe1—C4	40.68 (9)	C8—C7—Fe1	70.25 (14)
C1—Fe1—C4	69.13 (10)	C6—C7—Fe1	69.59 (14)
C6—Fe1—C4	163.18 (11)	C8—C7—H7	126.3
C7—Fe1—C4	154.22 (12)	C6—C7—H7	126.3
C9—Fe1—C4	108.19 (11)	Fe1—C7—H7	125.4
C10—Fe1—C2	156.06 (12)	C9—C8—C7	107.7 (3)
C5—Fe1—C2	68.87 (8)	C9—C8—Fe1	69.94 (14)
C1—Fe1—C2	41.18 (8)	C7—C8—Fe1	69.38 (14)
C6—Fe1—C2	121.07 (11)	C9—C8—H8	126.1
C7—Fe1—C2	108.08 (10)	C7—C8—H8	126.1
C9—Fe1—C2	162.73 (11)	Fe1—C8—H8	126.1
C4—Fe1—C2	68.23 (10)	C10—C9—C8	108.9 (3)
C10—Fe1—C8	66.96 (11)	C10—C9—Fe1	69.91 (14)
C5—Fe1—C8	153.99 (11)	C8—C9—Fe1	70.33 (14)
C1—Fe1—C8	163.43 (10)	C10—C9—H9	125.5
C6—Fe1—C8	67.85 (12)	C8—C9—H9	125.5
C7—Fe1—C8	40.38 (12)	Fe1—C9—H9	125.8

C9—Fe1—C8	39.72 (11)	C9—C10—C6	108.8 (2)
C4—Fe1—C8	119.76 (12)	C9—C10—Fe1	70.60 (15)
C2—Fe1—C8	126.08 (10)	C6—C10—Fe1	70.00 (14)
C10—Fe1—C3	162.78 (13)	C9—C10—H10	125.6
C5—Fe1—C3	68.49 (9)	C6—C10—H10	125.6
C1—Fe1—C3	68.83 (9)	Fe1—C10—H10	125.4
C6—Fe1—C3	155.20 (12)	O1—C11—N1	122.5 (2)
C7—Fe1—C3	120.02 (11)	O1—C11—C1	122.14 (19)
C9—Fe1—C3	126.23 (12)	N1—C11—C1	115.31 (17)
C4—Fe1—C3	40.59 (9)	C13—C12—N1	125.55 (19)
C2—Fe1—C3	40.22 (9)	C13—C12—C17	119.11 (19)
C8—Fe1—C3	107.98 (10)	N1—C12—C17	115.32 (16)
C11—N1—C12	128.71 (18)	C12—C13—C14	120.5 (2)
C11—N1—H1	119.1 (16)	C12—C13—H13	119.7
C12—N1—H1	111.5 (16)	C14—C13—H13	119.7
C20—N2—C17	116.78 (17)	C15—C14—C13	121.7 (2)
C2—C1—C5	106.7 (2)	C15—C14—H14	119.2
C2—C1—C11	123.43 (18)	C13—C14—H14	119.2
C5—C1—C11	129.84 (17)	C14—C15—C16	119.8 (2)
C2—C1—Fe1	69.76 (11)	C14—C15—H15	120.1
C5—C1—Fe1	69.23 (11)	C16—C15—H15	120.1
C11—C1—Fe1	123.63 (14)	C18—C16—C17	117.39 (18)
C3—C2—C1	108.78 (19)	C18—C16—C15	123.4 (2)
C3—C2—Fe1	70.34 (12)	C17—C16—C15	119.2 (2)
C1—C2—Fe1	69.06 (11)	N2—C17—C16	122.62 (17)
C3—C2—H2	125.6	N2—C17—C12	117.68 (17)
C1—C2—H2	125.6	C16—C17—C12	119.70 (17)
Fe1—C2—H2	126.6	C19—C18—C16	119.8 (2)
C2—C3—C4	108.1 (2)	C19—C18—H18	120.1
C2—C3—Fe1	69.44 (12)	C16—C18—H18	120.1
C4—C3—Fe1	69.24 (12)	C18—C19—C20	118.7 (2)
C2—C3—H3	125.9	C18—C19—H19	120.7
C4—C3—H3	125.9	C20—C19—H19	120.7
Fe1—C3—H3	127.0	N2—C20—C19	124.65 (19)
C5—C4—C3	108.3 (2)	N2—C20—H20	117.7
C5—C4—Fe1	69.29 (12)	C19—C20—H20	117.7
C3—C4—Fe1	70.16 (12)		
C10—Fe1—C1—C2	159.14 (15)	C5—Fe1—C6—C7	-167.14 (16)
C5—Fe1—C1—C2	-117.82 (19)	C1—Fe1—C6—C7	-124.89 (16)
C6—Fe1—C1—C2	117.14 (16)	C9—Fe1—C6—C7	81.22 (18)
C7—Fe1—C1—C2	75.68 (18)	C4—Fe1—C6—C7	158.4 (3)
C9—Fe1—C1—C2	-168.8 (2)	C2—Fe1—C6—C7	-81.73 (18)
C4—Fe1—C1—C2	-80.40 (15)	C8—Fe1—C6—C7	38.08 (16)
C8—Fe1—C1—C2	45.1 (4)	C3—Fe1—C6—C7	-47.0 (3)
C3—Fe1—C1—C2	-36.77 (13)	C10—C6—C7—C8	-0.3 (3)
C10—Fe1—C1—C5	-83.03 (17)	Fe1—C6—C7—C8	-60.39 (16)
C6—Fe1—C1—C5	-125.04 (14)	C10—C6—C7—Fe1	60.05 (17)

C7—Fe1—C1—C5	-166.50 (15)	C10—Fe1—C7—C8	80.21 (18)
C9—Fe1—C1—C5	-51.0 (3)	C5—Fe1—C7—C8	158.8 (3)
C4—Fe1—C1—C5	37.42 (12)	C1—Fe1—C7—C8	-167.07 (14)
C2—Fe1—C1—C5	117.82 (19)	C6—Fe1—C7—C8	118.2 (2)
C8—Fe1—C1—C5	163.0 (3)	C9—Fe1—C7—C8	37.31 (16)
C3—Fe1—C1—C5	81.05 (14)	C4—Fe1—C7—C8	-47.7 (3)
C10—Fe1—C1—C11	41.8 (2)	C2—Fe1—C7—C8	-124.92 (16)
C5—Fe1—C1—C11	124.8 (2)	C3—Fe1—C7—C8	-82.58 (17)
C6—Fe1—C1—C11	-0.25 (19)	C10—Fe1—C7—C6	-37.94 (17)
C7—Fe1—C1—C11	-41.7 (2)	C5—Fe1—C7—C6	40.7 (4)
C9—Fe1—C1—C11	73.8 (3)	C1—Fe1—C7—C6	74.78 (18)
C4—Fe1—C1—C11	162.22 (18)	C9—Fe1—C7—C6	-80.84 (18)
C2—Fe1—C1—C11	-117.4 (2)	C4—Fe1—C7—C6	-165.9 (2)
C8—Fe1—C1—C11	-72.3 (4)	C2—Fe1—C7—C6	116.93 (17)
C3—Fe1—C1—C11	-154.15 (18)	C8—Fe1—C7—C6	-118.2 (2)
C5—C1—C2—C3	-0.4 (2)	C3—Fe1—C7—C6	159.27 (16)
C11—C1—C2—C3	176.95 (18)	C6—C7—C8—C9	0.3 (3)
Fe1—C1—C2—C3	59.31 (14)	Fe1—C7—C8—C9	-59.66 (17)
C5—C1—C2—Fe1	-59.67 (13)	C6—C7—C8—Fe1	59.97 (16)
C11—C1—C2—Fe1	117.64 (18)	C10—Fe1—C8—C9	36.84 (19)
C10—Fe1—C2—C3	-169.1 (2)	C5—Fe1—C8—C9	-47.9 (3)
C5—Fe1—C2—C3	-81.27 (14)	C1—Fe1—C8—C9	158.4 (3)
C1—Fe1—C2—C3	-120.17 (18)	C6—Fe1—C8—C9	80.49 (19)
C6—Fe1—C2—C3	158.25 (14)	C7—Fe1—C8—C9	118.9 (2)
C7—Fe1—C2—C3	115.40 (16)	C4—Fe1—C8—C9	-82.8 (2)
C9—Fe1—C2—C3	43.8 (4)	C2—Fe1—C8—C9	-166.38 (17)
C4—Fe1—C2—C3	-37.43 (13)	C3—Fe1—C8—C9	-125.57 (18)
C8—Fe1—C2—C3	74.31 (18)	C10—Fe1—C8—C7	-82.11 (19)
C10—Fe1—C2—C1	-48.9 (3)	C5—Fe1—C8—C7	-166.9 (2)
C5—Fe1—C2—C1	38.90 (13)	C1—Fe1—C8—C7	39.4 (4)
C6—Fe1—C2—C1	-81.57 (16)	C6—Fe1—C8—C7	-38.46 (16)
C7—Fe1—C2—C1	-124.43 (16)	C9—Fe1—C8—C7	-118.9 (2)
C9—Fe1—C2—C1	164.0 (3)	C4—Fe1—C8—C7	158.25 (16)
C4—Fe1—C2—C1	82.75 (14)	C2—Fe1—C8—C7	74.68 (19)
C8—Fe1—C2—C1	-165.52 (15)	C3—Fe1—C8—C7	115.49 (17)
C3—Fe1—C2—C1	120.17 (18)	C7—C8—C9—C10	-0.2 (3)
C1—C2—C3—C4	0.1 (2)	Fe1—C8—C9—C10	-59.46 (18)
Fe1—C2—C3—C4	58.59 (15)	C7—C8—C9—Fe1	59.30 (16)
C1—C2—C3—Fe1	-58.52 (14)	C5—Fe1—C9—C10	-82.30 (19)
C10—Fe1—C3—C2	164.9 (3)	C1—Fe1—C9—C10	-45.7 (3)
C5—Fe1—C3—C2	82.30 (13)	C6—Fe1—C9—C10	37.51 (17)
C1—Fe1—C3—C2	37.62 (12)	C7—Fe1—C9—C10	81.92 (19)
C6—Fe1—C3—C2	-49.2 (3)	C4—Fe1—C9—C10	-125.21 (17)
C7—Fe1—C3—C2	-82.63 (16)	C2—Fe1—C9—C10	159.7 (3)
C9—Fe1—C3—C2	-165.24 (14)	C8—Fe1—C9—C10	119.8 (3)
C4—Fe1—C3—C2	119.84 (19)	C3—Fe1—C9—C10	-166.61 (16)
C8—Fe1—C3—C2	-125.11 (15)	C10—Fe1—C9—C8	-119.8 (3)
C10—Fe1—C3—C4	45.1 (4)	C5—Fe1—C9—C8	157.87 (17)

C5—Fe1—C3—C4	-37.54 (14)	C1—Fe1—C9—C8	-165.6 (2)
C1—Fe1—C3—C4	-82.22 (15)	C6—Fe1—C9—C8	-82.32 (19)
C6—Fe1—C3—C4	-169.0 (2)	C7—Fe1—C9—C8	-37.91 (18)
C7—Fe1—C3—C4	157.53 (16)	C4—Fe1—C9—C8	114.96 (18)
C9—Fe1—C3—C4	74.93 (18)	C2—Fe1—C9—C8	39.9 (4)
C2—Fe1—C3—C4	-119.84 (19)	C3—Fe1—C9—C8	73.6 (2)
C8—Fe1—C3—C4	115.05 (16)	C8—C9—C10—C6	-0.1 (3)
C2—C3—C4—C5	0.3 (2)	Fe1—C9—C10—C6	-59.78 (17)
Fe1—C3—C4—C5	58.97 (15)	C8—C9—C10—Fe1	59.72 (18)
C2—C3—C4—Fe1	-58.72 (15)	C7—C6—C10—C9	0.3 (3)
C10—Fe1—C4—C5	75.47 (18)	Fe1—C6—C10—C9	60.15 (18)
C1—Fe1—C4—C5	-38.14 (12)	C7—C6—C10—Fe1	-59.90 (16)
C6—Fe1—C4—C5	44.4 (4)	C5—Fe1—C10—C9	115.56 (16)
C7—Fe1—C4—C5	-169.1 (2)	C1—Fe1—C10—C9	159.40 (15)
C9—Fe1—C4—C5	115.51 (15)	C6—Fe1—C10—C9	-119.5 (2)
C2—Fe1—C4—C5	-82.48 (14)	C7—Fe1—C10—C9	-80.97 (18)
C8—Fe1—C4—C5	157.37 (14)	C4—Fe1—C10—C9	73.90 (19)
C3—Fe1—C4—C5	-119.6 (2)	C2—Fe1—C10—C9	-165.3 (2)
C10—Fe1—C4—C3	-164.95 (16)	C8—Fe1—C10—C9	-37.05 (16)
C5—Fe1—C4—C3	119.6 (2)	C3—Fe1—C10—C9	39.1 (4)
C1—Fe1—C4—C3	81.44 (15)	C5—Fe1—C10—C6	-124.94 (17)
C6—Fe1—C4—C3	164.0 (3)	C1—Fe1—C10—C6	-81.10 (18)
C7—Fe1—C4—C3	-49.6 (3)	C7—Fe1—C10—C6	38.53 (17)
C9—Fe1—C4—C3	-124.92 (16)	C9—Fe1—C10—C6	119.5 (2)
C2—Fe1—C4—C3	37.09 (13)	C4—Fe1—C10—C6	-166.60 (16)
C8—Fe1—C4—C3	-83.05 (17)	C2—Fe1—C10—C6	-45.8 (3)
C3—C4—C5—C1	-0.5 (2)	C8—Fe1—C10—C6	82.45 (19)
Fe1—C4—C5—C1	59.03 (14)	C3—Fe1—C10—C6	158.6 (3)
C3—C4—C5—Fe1	-59.52 (15)	C12—N1—C11—O1	5.3 (3)
C2—C1—C5—C4	0.5 (2)	C12—N1—C11—C1	-174.16 (18)
C11—C1—C5—C4	-176.56 (19)	C2—C1—C11—O1	6.2 (3)
Fe1—C1—C5—C4	-59.49 (15)	C5—C1—C11—O1	-177.1 (2)
C2—C1—C5—Fe1	60.01 (13)	Fe1—C1—C11—O1	92.8 (2)
C11—C1—C5—Fe1	-117.1 (2)	C2—C1—C11—N1	-174.32 (18)
C10—Fe1—C5—C4	-124.51 (17)	C5—C1—C11—N1	2.3 (3)
C1—Fe1—C5—C4	119.40 (18)	Fe1—C1—C11—N1	-87.69 (19)
C6—Fe1—C5—C4	-165.47 (16)	C11—N1—C12—C13	-8.9 (3)
C7—Fe1—C5—C4	162.7 (3)	C11—N1—C12—C17	169.84 (19)
C9—Fe1—C5—C4	-82.90 (17)	N1—C12—C13—C14	179.05 (19)
C2—Fe1—C5—C4	80.77 (15)	C17—C12—C13—C14	0.4 (3)
C8—Fe1—C5—C4	-49.6 (3)	C12—C13—C14—C15	-1.0 (3)
C3—Fe1—C5—C4	37.46 (14)	C13—C14—C15—C16	0.6 (3)
C10—Fe1—C5—C1	116.08 (15)	C14—C15—C16—C18	-178.2 (2)
C6—Fe1—C5—C1	75.13 (16)	C14—C15—C16—C17	0.5 (3)
C7—Fe1—C5—C1	43.3 (4)	C20—N2—C17—C16	1.9 (3)
C9—Fe1—C5—C1	157.69 (15)	C20—N2—C17—C12	-177.45 (17)
C4—Fe1—C5—C1	-119.40 (18)	C18—C16—C17—N2	-1.8 (3)
C2—Fe1—C5—C1	-38.63 (13)	C15—C16—C17—N2	179.49 (18)

C8—Fe1—C5—C1	-169.0 (2)	C18—C16—C17—C12	177.61 (17)
C3—Fe1—C5—C1	-81.94 (14)	C15—C16—C17—C12	-1.1 (3)
C5—Fe1—C6—C10	74.7 (2)	C13—C12—C17—N2	-179.88 (17)
C1—Fe1—C6—C10	116.98 (17)	N1—C12—C17—N2	1.3 (2)
C7—Fe1—C6—C10	-118.1 (2)	C13—C12—C17—C16	0.7 (3)
C9—Fe1—C6—C10	-36.91 (16)	N1—C12—C17—C16	-178.11 (16)
C4—Fe1—C6—C10	40.3 (4)	C17—C16—C18—C19	-0.1 (3)
C2—Fe1—C6—C10	160.14 (15)	C15—C16—C18—C19	178.6 (2)
C8—Fe1—C6—C10	-80.05 (18)	C16—C18—C19—C20	1.7 (3)
C3—Fe1—C6—C10	-165.1 (2)	C17—N2—C20—C19	-0.2 (3)
C10—Fe1—C6—C7	118.1 (2)	C18—C19—C20—N2	-1.6 (3)
