

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Methyl-*N*-[(*Z*)-3-(4-methylphenyl-sulfonyl)-1,3-thiazolidin-2-ylidene]-benzenesulfonamide

 Hui-Ling Hu,^a Geng-Ren Yang^a and Chun-Wei Yeh^{b*}

^aDepartment of Chemical Engineering and Material Engineering, Graduate School of Materials Applied Technology, Nanya Institute of Technology, Chung-Li, Taiwan, and ^bDepartment of Chemistry, Chung-Yuan Christian University, Chung-Li, Taiwan
Correspondence e-mail: cwye@cycu.org.tw


Received 9 July 2011; accepted 17 July 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.107; data-to-parameter ratio = 19.7.

In the crystal structure of the title compound, $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_4\text{S}_3$, molecules are connected into centrosymmetric dimers *via* weak intermolecular $\text{C}-\text{H}\cdots\pi$ interactions. These dimers are further connected through a series of weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, while further $\text{C}-\text{H}\cdots\pi$ interactions involving the phenyl and thiazoline rings are also observed. The thiazolidine ring is twisted from the benzene rings rings by dihedral angles of 79.1 (1) and 85.0 (1)°, while the dihedral angle between two benzene rings is 76.0 (1)°.

Related literature

For background to *N*-heterocyclic sulfanilamide derivatives, see: Kuz'mina *et al.* (1962); Jensen & Thorsteinsson (1941); Hunter & Kolloff (1943); Hultquist *et al.* (1951). For a related synthesis, see: Razvodovskaya *et al.* (1990).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_4\text{S}_3$
 $M_r = 410.51$
Monoclinic, $P2_1/n$

$a = 9.3825$ (2) Å
 $b = 14.4047$ (2) Å
 $c = 14.2279$ (3) Å

$\beta = 102.666$ (1)°
 $V = 1876.14$ (6) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.42$ mm⁻¹
 $T = 296$ K
 $0.40 \times 0.40 \times 0.40$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\min} = 0.845$, $T_{\max} = 0.845$

17749 measured reflections
4652 independent reflections
3756 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.107$
 $S = 1.04$
4652 reflections

236 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and Cg2 are the centroids of the $\text{C4}-\text{C9}$ and $\text{C11}-\text{C16}$ benzene rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C10}-\text{H10B}\cdots\text{Cg1}^{\text{i}}$	0.97	2.91	3.567 (1)	127
$\text{C2}-\text{H2B}\cdots\text{Cg2}^{\text{ii}}$	0.97	3.09	3.821 (1)	134
$\text{C1}-\text{H1B}\cdots\text{O1}^{\text{iii}}$	0.97	2.59	3.394 (3)	141
$\text{C12}-\text{H12A}\cdots\text{O3}^{\text{iii}}$	0.93	2.47	3.318 (2)	151

Symmetry codes: (i) $-x, -y, -z$; (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $-x, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2010); cell refinement: SAINT (Bruker, 2010); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

We are grateful to the National Science Council of the Republic of China and the Nanya Institute of Technology for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2445).

References

- Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2010). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Hultquist, M. E., Germann, R. P., Webb, J. S., Wright, W. B. Jr, Roth, B., Smith, J. M. Jr & SubbaRow, Y. (1951). *J. Am. Chem. Soc.* **73**, 2558–2566.
Hunter, J. H. & Kolloff, H. G. (1943). *J. Am. Chem. Soc.* **65**, 156–159.
Jensen, K. A. & Thorsteinsson, T. (1941). *Dansk Tidsskrift Farmaci*, **15**, 41–77.
Kuz'mina, K. K., Ostroumova, N. G., Markova, Yu. V. & Shchukina, M. N. (1962). *Zhurnal Obshchei Khimii*, **32**, 3390–3393.
Razvodovskaya, L. V., Vorob'eva, N. N., Grapov, A. F. & Mel'nikov, N. N. (1990). *Zhurnal Obshchei Khimii*, **60**, 1518–1525.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, o2103 [doi:10.1107/S1600536811028807]

4-Methyl-N-[(Z)-3-(4-methylphenylsulfonyl)-1,3-thiazolidin-2-ylidene]benzenesulfonamide

Hui-Ling Hu, Geng-Ren Yang and Chun-Wei Yeh

S1. Comment

In a series of N-heterocyclic sulfanilamide derivatives which prepared and are investigating biologically one of the compounds, 2-sulfanilyl-aminothiazoline, proved to be of particular interest, both chemically and therapeutically. (Kuz'mina *et al.*, 1962; Jensen *et al.*, 1941; Hunter *et al.*, 1943; Hultquist *et al.*, 1951). The synthesis and character the 3-substituted 2-(thiophosphorylimino)thiazolidine compounds are also reported (Razvodovskaya *et al.*, 1990). Within this project the crystal structure of the title compound was determined. The crystal structure features inversion-related dimers linked by the weak intermolecular C—H \cdots π i interactions in the solid state, while *Cg1* and *Cg2* are the centers of C4—C9 and C11—C16 and these carbon atoms of mean deviation from plane are 0.0008 and 0.0043 Å. Weak C—H \cdots O hydrogen bonds among the molecules are also observed in the solid state. The thiazolidine and the phenyl rings are not coplanar but twisted with each other by an interplanar angles of 79.1 (1) and 85.0 (1)°, respectively, while the dihedral angle between two phenyl groups is 76.0 (1)°.

S2. Experimental

The title compound was prepared according to a published procedure (Razvodovskaya *et al.*, 1990). Block like crystals suitable for X-ray crystallography were obtained by slow evaporation of the solvent from a solution of the title compound in methanol.

S3. Refinement

All the hydrogen atoms were discernible in the difference Fourier maps. However, they were situated into the idealized positions and constrained by the riding atom approximation: C—H_{methyl} = 0.96 Å and C—H_{methylene} = 0.97 Å while the methyls and methylenes were allowed to rotate about their respective axes; C—H_{aryl} = 0.93 Å; $U_{\text{iso}}(\text{H}_{\text{methyl}}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$; $U_{\text{iso}}(\text{H}_{\text{aryl}} \text{ or methylene}) = 1.2U_{\text{eq}}(\text{C}_{\text{aryl}} \text{ or methylene})$.




Figure 1

Crystal structure of the title compound with atom labeling and displacement ellipsoids drawn at the 30% probability level.

4-Methyl-N-[(Z)-3-(4-methylphenylsulfonyl)-1,3-thiazolidin-2-ylidene]benzenesulfonamide

Crystal data

$C_{17}H_{18}N_2O_4S_3$

$M_r = 410.51$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 9.3825\ (2)\ \text{\AA}$

$b = 14.4047\ (2)\ \text{\AA}$

$c = 14.2279\ (3)\ \text{\AA}$

$\beta = 102.666\ (1)^\circ$

$V = 1876.14\ (6)\ \text{\AA}^3$

$Z = 4$

$F(000) = 856$

$D_x = 1.453\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 8638 reflections

$\theta = 2.4\text{--}28.2^\circ$

$\mu = 0.42\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Block, colourless

$0.40 \times 0.40 \times 0.40\ \text{mm}$

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.845$, $T_{\max} = 0.845$

17749 measured reflections

4652 independent reflections

3756 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.0^\circ$

$h = -12 \rightarrow 12$

$k = -15 \rightarrow 19$

$l = -17 \rightarrow 18$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.107$ $S = 1.04$

4652 reflections

236 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0497P)^2 + 0.6237P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$ Extinction correction: *SHELXL*, $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0091 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.16502 (15)	0.17067 (10)	0.38640 (10)	0.0446 (3)
N2	0.08736 (15)	0.26664 (10)	0.25505 (10)	0.0448 (3)
S1	0.34585 (5)	0.17353 (4)	0.27357 (4)	0.06158 (16)
S2	0.10549 (5)	0.31474 (3)	0.15432 (3)	0.04983 (13)
S3	0.01808 (5)	0.19186 (3)	0.43270 (3)	0.04805 (13)
O1	-0.00727 (19)	0.38361 (9)	0.13422 (10)	0.0693 (4)
O2	0.25206 (18)	0.34381 (12)	0.15775 (11)	0.0739 (4)
O3	0.04496 (18)	0.13807 (11)	0.51854 (10)	0.0702 (4)
O4	0.00217 (15)	0.28956 (9)	0.43817 (10)	0.0580 (3)
C1	0.2585 (2)	0.08847 (14)	0.41762 (15)	0.0590 (5)
H1A	0.2091	0.0321	0.3909	0.071*
H1B	0.2818	0.0836	0.4873	0.071*
C2	0.3947 (2)	0.10280 (16)	0.38077 (16)	0.0664 (6)
H2A	0.4335	0.0436	0.3656	0.080*
H2B	0.4684	0.1338	0.4290	0.080*
C3	0.18420 (17)	0.21042 (11)	0.30207 (11)	0.0407 (3)
C4	0.06041 (19)	0.22754 (11)	0.06650 (12)	0.0431 (4)
C5	-0.08309 (19)	0.19554 (12)	0.04084 (13)	0.0481 (4)
H5A	-0.1533	0.2177	0.0723	0.058*
C6	-0.1198 (2)	0.13072 (13)	-0.03153 (14)	0.0534 (4)
H6A	-0.2156	0.1094	-0.0486	0.064*
C7	-0.0173 (2)	0.09647 (13)	-0.07954 (13)	0.0542 (4)
C8	0.1244 (2)	0.12907 (15)	-0.05287 (14)	0.0607 (5)

H8A	0.1943	0.1068	-0.0844	0.073*
C9	0.1647 (2)	0.19422 (14)	0.01986 (14)	0.0538 (4)
H9A	0.2606	0.2152	0.0370	0.065*
C10	-0.0606 (3)	0.02670 (16)	-0.15964 (16)	0.0787 (7)
H10A	-0.1623	0.0119	-0.1675	0.118*
H10B	-0.0034	-0.0287	-0.1439	0.118*
H10C	-0.0438	0.0524	-0.2185	0.118*
C11	-0.12963 (18)	0.14496 (12)	0.34887 (12)	0.0460 (4)
C12	-0.1544 (2)	0.04982 (13)	0.35063 (15)	0.0578 (5)
H12A	-0.0965	0.0124	0.3971	0.069*
C13	-0.2660 (2)	0.01228 (15)	0.28240 (17)	0.0652 (5)
H13A	-0.2837	-0.0512	0.2836	0.078*
C14	-0.3526 (2)	0.06605 (15)	0.21206 (15)	0.0579 (5)
C15	-0.3267 (2)	0.16119 (15)	0.21243 (15)	0.0557 (5)
H15A	-0.3850	0.1986	0.1661	0.067*
C16	-0.21635 (19)	0.20075 (13)	0.28026 (14)	0.0507 (4)
H16A	-0.2002	0.2644	0.2800	0.061*
C17	-0.4719 (3)	0.0231 (2)	0.13607 (19)	0.0864 (8)
H17A	-0.4738	-0.0427	0.1461	0.130*
H17B	-0.4536	0.0354	0.0735	0.130*
H17C	-0.5644	0.0494	0.1403	0.130*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0461 (7)	0.0435 (7)	0.0418 (7)	0.0049 (6)	0.0045 (6)	-0.0002 (6)
N2	0.0453 (7)	0.0459 (8)	0.0429 (7)	0.0017 (6)	0.0095 (6)	0.0013 (6)
S1	0.0508 (3)	0.0734 (3)	0.0623 (3)	0.0140 (2)	0.0163 (2)	-0.0073 (2)
S2	0.0627 (3)	0.0401 (2)	0.0470 (2)	-0.00536 (18)	0.0128 (2)	0.00129 (17)
S3	0.0555 (3)	0.0491 (3)	0.0406 (2)	-0.00057 (18)	0.01281 (18)	-0.00071 (17)
O1	0.1039 (12)	0.0425 (7)	0.0589 (8)	0.0189 (7)	0.0122 (8)	0.0043 (6)
O2	0.0802 (10)	0.0758 (10)	0.0684 (9)	-0.0367 (8)	0.0217 (8)	-0.0044 (8)
O3	0.0878 (11)	0.0805 (10)	0.0427 (7)	0.0010 (8)	0.0149 (7)	0.0108 (7)
O4	0.0645 (8)	0.0504 (7)	0.0619 (8)	0.0006 (6)	0.0197 (7)	-0.0130 (6)
C1	0.0684 (12)	0.0496 (10)	0.0521 (10)	0.0146 (9)	-0.0017 (9)	0.0011 (8)
C2	0.0637 (12)	0.0655 (13)	0.0631 (12)	0.0251 (10)	-0.0009 (10)	-0.0143 (10)
C3	0.0399 (8)	0.0402 (8)	0.0402 (8)	-0.0027 (6)	0.0052 (6)	-0.0082 (6)
C4	0.0484 (9)	0.0396 (8)	0.0416 (8)	0.0018 (7)	0.0110 (7)	0.0055 (7)
C5	0.0472 (9)	0.0468 (9)	0.0520 (10)	0.0050 (7)	0.0144 (7)	0.0031 (8)
C6	0.0529 (10)	0.0479 (10)	0.0546 (10)	-0.0025 (8)	0.0014 (8)	0.0027 (8)
C7	0.0726 (12)	0.0458 (10)	0.0403 (9)	0.0106 (9)	0.0041 (8)	0.0038 (7)
C8	0.0659 (12)	0.0680 (13)	0.0519 (11)	0.0153 (10)	0.0210 (9)	-0.0008 (9)
C9	0.0484 (9)	0.0636 (11)	0.0512 (10)	0.0007 (8)	0.0153 (8)	0.0020 (8)
C10	0.1124 (19)	0.0631 (13)	0.0513 (12)	0.0143 (13)	-0.0025 (12)	-0.0083 (10)
C11	0.0473 (9)	0.0452 (9)	0.0478 (9)	-0.0052 (7)	0.0155 (7)	0.0038 (7)
C12	0.0651 (12)	0.0470 (10)	0.0626 (12)	-0.0066 (9)	0.0171 (9)	0.0100 (9)
C13	0.0710 (13)	0.0470 (11)	0.0826 (15)	-0.0152 (9)	0.0279 (11)	-0.0052 (10)
C14	0.0491 (10)	0.0687 (12)	0.0607 (11)	-0.0102 (9)	0.0226 (9)	-0.0147 (10)

C15	0.0441 (9)	0.0665 (12)	0.0578 (11)	-0.0001 (8)	0.0139 (8)	0.0053 (9)
C16	0.0467 (9)	0.0451 (9)	0.0622 (11)	-0.0022 (7)	0.0157 (8)	0.0064 (8)
C17	0.0689 (14)	0.0996 (19)	0.0894 (18)	-0.0144 (13)	0.0148 (13)	-0.0413 (15)

Geometric parameters (Å, °)

N1—C3	1.376 (2)	C6—H6A	0.9300
N1—C1	1.483 (2)	C7—C8	1.382 (3)
N1—S3	1.6811 (15)	C7—C10	1.507 (3)
N2—C3	1.289 (2)	C8—C9	1.387 (3)
N2—S2	1.6340 (15)	C8—H8A	0.9300
S1—C3	1.7368 (16)	C9—H9A	0.9300
S1—C2	1.808 (2)	C10—H10A	0.9600
S2—O2	1.4282 (15)	C10—H10B	0.9600
S2—O1	1.4327 (15)	C10—H10C	0.9600
S2—C4	1.7566 (17)	C11—C16	1.383 (3)
S3—O4	1.4192 (14)	C11—C12	1.391 (3)
S3—O3	1.4215 (14)	C12—C13	1.373 (3)
S3—C11	1.7536 (18)	C12—H12A	0.9300
C1—C2	1.498 (3)	C13—C14	1.380 (3)
C1—H1A	0.9700	C13—H13A	0.9300
C1—H1B	0.9700	C14—C15	1.392 (3)
C2—H2A	0.9700	C14—C17	1.508 (3)
C2—H2B	0.9700	C15—C16	1.375 (3)
C4—C9	1.383 (2)	C15—H15A	0.9300
C4—C5	1.394 (2)	C16—H16A	0.9300
C5—C6	1.377 (3)	C17—H17A	0.9600
C5—H5A	0.9300	C17—H17B	0.9600
C6—C7	1.386 (3)	C17—H17C	0.9600
C3—N1—C1	114.30 (15)	C7—C6—H6A	119.2
C3—N1—S3	122.77 (11)	C8—C7—C6	118.23 (17)
C1—N1—S3	120.66 (13)	C8—C7—C10	121.2 (2)
C3—N2—S2	121.62 (12)	C6—C7—C10	120.6 (2)
C3—S1—C2	92.76 (9)	C7—C8—C9	121.55 (18)
O2—S2—O1	117.85 (10)	C7—C8—H8A	119.2
O2—S2—N2	112.27 (9)	C9—C8—H8A	119.2
O1—S2—N2	104.73 (8)	C4—C9—C8	119.15 (18)
O2—S2—C4	108.29 (9)	C4—C9—H9A	120.4
O1—S2—C4	107.52 (9)	C8—C9—H9A	120.4
N2—S2—C4	105.40 (8)	C7—C10—H10A	109.5
O4—S3—O3	119.63 (9)	C7—C10—H10B	109.5
O4—S3—N1	107.86 (8)	H10A—C10—H10B	109.5
O3—S3—N1	103.39 (8)	C7—C10—H10C	109.5
O4—S3—C11	110.02 (9)	H10A—C10—H10C	109.5
O3—S3—C11	109.83 (9)	H10B—C10—H10C	109.5
N1—S3—C11	104.90 (8)	C16—C11—C12	120.69 (18)
N1—C1—C2	106.21 (17)	C16—C11—S3	120.74 (14)

N1—C1—H1A	110.5	C12—C11—S3	118.52 (15)
C2—C1—H1A	110.5	C13—C12—C11	118.67 (19)
N1—C1—H1B	110.5	C13—C12—H12A	120.7
C2—C1—H1B	110.5	C11—C12—H12A	120.7
H1A—C1—H1B	108.7	C12—C13—C14	121.87 (19)
C1—C2—S1	107.17 (13)	C12—C13—H13A	119.1
C1—C2—H2A	110.3	C14—C13—H13A	119.1
S1—C2—H2A	110.3	C13—C14—C15	118.41 (19)
C1—C2—H2B	110.3	C13—C14—C17	121.1 (2)
S1—C2—H2B	110.3	C15—C14—C17	120.5 (2)
H2A—C2—H2B	108.5	C16—C15—C14	120.96 (19)
N2—C3—N1	120.10 (15)	C16—C15—H15A	119.5
N2—C3—S1	128.55 (13)	C14—C15—H15A	119.5
N1—C3—S1	111.35 (12)	C15—C16—C11	119.39 (17)
C9—C4—C5	120.25 (17)	C15—C16—H16A	120.3
C9—C4—S2	120.28 (14)	C11—C16—H16A	120.3
C5—C4—S2	119.40 (13)	C14—C17—H17A	109.5
C6—C5—C4	119.28 (17)	C14—C17—H17B	109.5
C6—C5—H5A	120.4	H17A—C17—H17B	109.5
C4—C5—H5A	120.4	C14—C17—H17C	109.5
C5—C6—C7	121.53 (18)	H17A—C17—H17C	109.5
C5—C6—H6A	119.2	H17B—C17—H17C	109.5

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C4—C9 and C11—C16 benzene rings, respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C10—H10 <i>B</i> ...Cg1 ⁱ	0.97	2.91	3.567 (1)	127
C2—H2 <i>B</i> ...Cg2 ⁱⁱ	0.97	3.09	3.821 (1)	134
C1—H1 <i>B</i> ...O1 ⁱⁱ	0.97	2.59	3.394 (3)	141
C12—H12 <i>A</i> ...O3 ⁱⁱⁱ	0.93	2.47	3.318 (2)	151

Symmetry codes: (i) $-x, -y, -z$; (ii) $x+1/2, -y+1/2, z+1/2$; (iii) $-x, -y, -z+1$.