

[2-(1*H*-Benzimidazol-2-yl- κ N³)aniline- κ N]dichloridozinc

Naser Eltaher Eltayeb,^{a,b} Siang Guan Teoh,^a Suchada Chantrapromma^c‡ and Hoong-Kun Fun^{d,*§}

^aSchool of Chemical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Chemistry, Faculty of Pure and Applied Sciences, International University of Africa, Sudan, ^cCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, and ^dX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia
Correspondence e-mail: hkfun@usm.my


Received 1 July 2011; accepted 4 July 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.027; wR factor = 0.081; data-to-parameter ratio = 42.7.

In the title benzimidazole mononuclear complex, $[ZnCl_2(C_{13}H_{11}N_3)]$, the Zn^{II} ion is four-coordinated in a distorted tetrahedral geometry by an imidazole N, an amino N and two Cl atoms. The dihedral angle between the benzimidazole and benzene rings is 9.57 (1)°. In the crystal, molecules are linked by weak N—H···Cl hydrogen bonds into layers parallel to the *bc* plane. π – π interactions with centroid–centroid distances in the range 3.4452 (8)–3.8074 (8) Å are also observed.

Related literature

For bond-length data, see: Allen *et al.* (1987). For background to benzimidazoles and their applications, see: Chassaing *et al.* (2008); Podunavac-Kuzmonovic *et al.* (1999); Xue *et al.* (2011). For related structures, see: Eltayeb *et al.* (2007; 2009; 2011); Maldonado-Rogado *et al.* (2007). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-5085-2009.
§ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$[ZnCl_2(C_{13}H_{11}N_3)]$	$V = 2782.32$ (18) Å ³
$M_r = 345.54$	$Z = 8$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 22.0252$ (7) Å	$\mu = 2.14$ mm ^{−1}
$b = 10.0651$ (3) Å	$T = 100$ K
$c = 15.3676$ (6) Å	$0.48 \times 0.31 \times 0.31$ mm
$\beta = 125.244$ (2)°	

Data collection

Bruker APEX DUO CCD area-detector diffractometer	50659 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	7344 independent reflections
$T_{\min} = 0.428$, $T_{\max} = 0.560$	5887 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$	172 parameters
$wR(F^2) = 0.081$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\max} = 0.54$ e Å ^{−3}
7344 reflections	$\Delta\rho_{\min} = -0.55$ e Å ^{−3}

Table 1
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N2—H1N2···Cl1 ⁱ	0.82	2.56	3.3503 (9)	164
N3—H1N3···Cl1 ⁱⁱ	0.89	2.49	3.3753 (9)	174
N3—H2N3···Cl2 ⁱⁱⁱ	0.94	2.44	3.3015 (12)	153

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x, -y + 2, -z$; (iii) $-x, y, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors thank the Malaysian Government and Universiti Sains Malaysia for the RU research grant (1001/PKIMIA/815067). NEE thanks Universiti Sains Malaysia for a post-doctoral fellowship and the International University of Africa (Sudan) for providing study leave. The authors also thank Universiti Sains Malaysia for the Research University grant No. 1001/PFIZIK/811160.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2622).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2009). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chassaing, C., Berger, M., Heckereth, A., Ilg, T., Jaeger, M., Kern, C., Schmid, K. & Uphoff, M. (2008). *J. Med. Chem.* **51**, 1111–1114.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Eltayeb, N. E., Teoh, S. G., Chantrapromma, S. & Fun, H.-K. (2007). *Acta Cryst. E63*, o4141–o4142.
- Eltayeb, N. E., Teoh, S. G., Quah, C. K. Fun, H.-K. & Adnan, R. (2009). *Acta Cryst. E65*, o1613–o1614.

- Eltayeb, N. E., Teoh, S. G., Yeap, C. S. & Fun, H.-K. (2011). *Acta Cryst. E*67, o1721–o1722.
- Maldonado-Rogado, M. A., Viñuelas-Zahinos, E., Luna-Giles, F. & Bernalte-García, A. (2007). *Polyhedron*, **26**, 3112–3120.
- Podunavac-Kuzmonovic, S. O., Leovac, L. M., Perisic-Janjic, N. U., Rogan, J. & Balaz, J. (1999). *J. Serb. Chem. Soc.* **64**, 381–388.
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D*65, 148–155.
- Xue, F., Luo, X., Ye, C., Ye, W. & Wang, Y. (2011). *Bioorg. Med. Chem.* **19**, 2641–2649.

supporting information

Acta Cryst. (2011). E67, m1062–m1063 [doi:10.1107/S1600536811026572]

[2-(1*H*-Benzimidazol-2-yl- κ N³)aniline- κ N]dichloridozinc

Naser Eltaher Eltayeb, Siang Guan Teoh, Suchada Chantrapromma and Hoong-Kun Fun

S1. Comment

Benzimidazole compounds have a wide range of biological properties such as antibacterial (Chassaing *et al.*, 2008) and inhibitory activity against enteroviruses (Xue *et al.*, 2011). The complexes of transition metal salts with benzimidazole derivatives have been extensively studied as models of some important biological molecules (Podunavac-Kuzmonovic *et al.*, 1999). As part of our ongoing structural studies of benzimidazoles (Eltayeb *et al.*, 2007; 2009; 2011) and as an extension of investigation on their complexes, the title zinc(II) complex, (I), is reported here.

Complex (I) is a mononuclear zinc(II) complex (Fig. 1) in which the coordination geometry around the zinc ion is distorted tetrahedral, with the metal four-coordinated by an imidazole N, an amino N and two Cl atoms. In the complex, the 2-(2-aminophenyl)-1*H*-benzimidazole acts as a bidentate ligand. The bond angles around the central metal zinc(II) show some deviations from ideal tetrahedral geometry [N3-Zn1-C11 = 109.27 (3) $^{\circ}$, N1-Zn1-Cl2 = 114.83 (3) $^{\circ}$, C11-Zn1-Cl2 = 117.844 (13) $^{\circ}$ and the bite angle N1-Zn1-N3 = 89.36 (3) $^{\circ}$]. The Zn-N [2.0068 (8) and 2.0471 (9) Å] and Zn-Cl [2.2041 (3) and 2.2589 (3) Å] bond lengths are comparable to those found in a similar zinc(II) benzimidazole complex (Maldonado-Rogado *et al.*, 2007). The benzimidazole ring system (C1–C7/N1–N2) is planar (*r.m.s.* of 0.0097 (1) Å), the larger deviation being observed for atom C1 (0.019 (1) Å). The dihedral angle between the benzimidazole and phenyl rings is 9.57 (6) $^{\circ}$. The bond lengths of ligand are within normal ranges (Allen *et al.*, 1987).


In the crystal structure (Fig. 2), the molecules are linked through N—H \cdots Cl hydrogen bonds into two dimensional layers parallel to the *bc* plane. $\pi\cdots\pi$ interactions are observed with centroid \cdots centroid distances: $Cg_1\cdots Cg_1^{iii} = 3.4452$ (8) Å, $Cg_1\cdots Cg_2^{iii} = 3.6879$ (9) Å and $Cg_2\cdots Cg_3^i = 3.8074$ (8) Å (Cg_1 , Cg_2 and Cg_3 are the centroids of the C1/C6–C7/N1–N2, C1–C6 and C8–C13 rings, respectively; symmetry codes: (i) -*x*, 1-*y*, -*z*; (iii) -*x*, *y*, 1/2-*z*).

S2. Experimental

The title compound was synthesized by adding 2-hydroxy-3-methylbenzaldehyde (0.136 g, 1.0 mmol) to a solution of 2-(2-aminophenyl)-1*H*-benzimidazole (0.209 g, 1.0 mmol) in ethanol (30 mL). The colour of the resulting solution was pale-yellow. Then upon addition of zinc chloride (0.136 g, 1.0 mmol), the colour of the solution became golden-yellow. The mixture was refluxed with stirring for 3 h. The resultant solution was filtered and the filtrate was evaporated to give a yellow solid product. Yellow block-shaped single crystals of the title compound suitable for *X*-ray structure determination were obtained by slow evaporation of an ethanol solution at room temperature after several days.

S3. Refinement

All H atoms were positioned geometrically and allowed to ride on their parent atoms, with d(N—H) = 0.82 Å for NH; 0.88 and 0.94 Å for NH₂, d(C—H) = 0.93 Å for aromatic. The U_{iso} values were constrained to be 1.2 U_{eq} of the carrier atoms. The highest residual electron density peak is located at 0.64 Å from C11 and the deepest hole is located at 0.49 Å from C11. An outlier reflection (2 4 3) was omitted.

**Figure 1**

The molecular structure of the title compound, with 50% probability displacement ellipsoids.





Figure 2

The crystal packing of the title compound viewed down the *b* axis. Hydrogen bonds are shown as dashed lines.

[2-(1*H*-Benzimidazol-2-yl-*kN*³)aniline- *kN*]dichloridozinc*Crystal data*

$M_r = 345.54$

Monoclinic, $C2/c$

Hall symbol: -C 2yc

$a = 22.0252 (7) \text{ \AA}$

$b = 10.0651 (3) \text{ \AA}$

$c = 15.3676 (6) \text{ \AA}$

$\beta = 125.244 (2)^\circ$

$V = 2782.32 (18) \text{ \AA}^3$

$Z = 8$

$F(000) = 1392$

$D_x = 1.650 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 7344 reflections

$\theta = 2.3\text{--}37.7^\circ$

$\mu = 2.14 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Block, yellow

$0.48 \times 0.31 \times 0.31 \text{ mm}$

Data collection

Bruker APEX DUO CCD area-detector
diffractometer

Radiation source: sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2009)

$T_{\min} = 0.428$, $T_{\max} = 0.560$

50659 measured reflections

7344 independent reflections

5887 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 37.7^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -37 \rightarrow 36$

$k = -17 \rightarrow 17$

$l = -25 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.081$

$S = 1.02$

7344 reflections

172 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0419P)^2 + 0.7801P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.54 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.55 \text{ e \AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.057453 (6)	0.825871 (11)	0.151068 (9)	0.03470 (4)
Cl1	0.094347 (18)	0.82859 (2)	0.04109 (2)	0.04377 (6)
Cl2	0.121812 (17)	0.94430 (3)	0.29852 (2)	0.04885 (7)
N1	0.03795 (5)	0.63755 (8)	0.17124 (6)	0.03277 (14)
N2	-0.01309 (5)	0.43948 (8)	0.11361 (7)	0.03566 (16)
H1N2	-0.0400	0.3776	0.0772	0.043*
N3	-0.05430 (5)	0.86317 (8)	0.06474 (7)	0.03630 (16)
H1N3	-0.0613	0.9453	0.0395	0.044*
H2N3	-0.0606	0.8661	0.1203	0.044*
C1	0.08602 (5)	0.54368 (9)	0.24599 (8)	0.03401 (16)
C2	0.15487 (6)	0.55968 (12)	0.34379 (9)	0.0441 (2)
H2A	0.1765	0.6429	0.3686	0.053*
C3	0.18917 (8)	0.44533 (15)	0.40179 (11)	0.0548 (3)
H3A	0.2351	0.4519	0.4673	0.066*
C4	0.15673 (9)	0.32000 (14)	0.36457 (13)	0.0578 (3)
H4A	0.1816	0.2457	0.4063	0.069*
C5	0.08893 (8)	0.30296 (12)	0.26782 (11)	0.0487 (3)
H5A	0.0677	0.2195	0.2428	0.058*
C6	0.05428 (6)	0.41848 (9)	0.20999 (8)	0.03638 (18)
C7	-0.02133 (5)	0.57180 (8)	0.09356 (7)	0.03092 (15)
C8	-0.08750 (5)	0.62967 (9)	-0.00210 (7)	0.03133 (15)
C9	-0.13949 (6)	0.54428 (11)	-0.08355 (9)	0.0409 (2)
H9A	-0.1313	0.4531	-0.0751	0.049*
C10	-0.20257 (7)	0.59227 (14)	-0.17600 (10)	0.0501 (3)
H10A	-0.2366	0.5336	-0.2286	0.060*
C11	-0.21502 (7)	0.72704 (15)	-0.19038 (10)	0.0556 (3)
H11A	-0.2567	0.7598	-0.2535	0.067*
C12	-0.16525 (7)	0.81339 (12)	-0.11062 (10)	0.0478 (3)
H12A	-0.1741	0.9043	-0.1203	0.057*
C13	-0.10222 (5)	0.76671 (9)	-0.01632 (7)	0.03314 (16)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.03589 (6)	0.02529 (5)	0.03528 (6)	-0.00291 (3)	0.01612 (5)	-0.00044 (3)
Cl1	0.05837 (16)	0.03021 (11)	0.04711 (13)	-0.00299 (9)	0.03297 (13)	0.00024 (8)
Cl2	0.05149 (15)	0.04321 (13)	0.04471 (13)	-0.01328 (11)	0.02364 (11)	-0.01315 (10)
N1	0.0341 (3)	0.0252 (3)	0.0324 (3)	-0.0009 (3)	0.0154 (3)	0.0028 (2)
N2	0.0392 (4)	0.0235 (3)	0.0404 (4)	-0.0010 (3)	0.0207 (3)	0.0010 (3)
N3	0.0372 (4)	0.0252 (3)	0.0366 (4)	0.0020 (3)	0.0156 (3)	-0.0006 (3)
C1	0.0358 (4)	0.0293 (4)	0.0345 (4)	0.0030 (3)	0.0188 (3)	0.0054 (3)
C2	0.0390 (5)	0.0421 (5)	0.0387 (5)	0.0026 (4)	0.0153 (4)	0.0045 (4)
C3	0.0460 (6)	0.0552 (7)	0.0454 (6)	0.0125 (5)	0.0162 (5)	0.0139 (5)
C4	0.0568 (7)	0.0461 (7)	0.0588 (7)	0.0189 (5)	0.0266 (6)	0.0219 (5)
C5	0.0541 (6)	0.0299 (4)	0.0586 (7)	0.0101 (4)	0.0306 (6)	0.0126 (4)

C6	0.0405 (4)	0.0275 (4)	0.0413 (4)	0.0039 (3)	0.0237 (4)	0.0055 (3)
C7	0.0345 (4)	0.0238 (3)	0.0331 (4)	-0.0006 (3)	0.0187 (3)	0.0011 (3)
C8	0.0316 (4)	0.0280 (3)	0.0313 (3)	-0.0009 (3)	0.0163 (3)	-0.0002 (3)
C9	0.0391 (5)	0.0345 (4)	0.0406 (5)	-0.0050 (4)	0.0180 (4)	-0.0065 (4)
C10	0.0374 (5)	0.0505 (6)	0.0421 (5)	-0.0056 (4)	0.0112 (4)	-0.0100 (5)
C11	0.0369 (5)	0.0533 (7)	0.0440 (6)	0.0034 (5)	0.0045 (4)	-0.0008 (5)
C12	0.0372 (5)	0.0385 (5)	0.0438 (5)	0.0061 (4)	0.0095 (4)	0.0038 (4)
C13	0.0307 (4)	0.0288 (4)	0.0334 (4)	0.0015 (3)	0.0147 (3)	0.0007 (3)

Geometric parameters (\AA , $\text{^{\circ}}$)

Zn1—N1	2.0068 (8)	C3—H3A	0.9300
Zn1—N3	2.0471 (9)	C4—C5	1.381 (2)
Zn1—Cl2	2.2041 (3)	C4—H4A	0.9300
Zn1—Cl1	2.2589 (3)	C5—C6	1.3933 (14)
N1—C7	1.3300 (12)	C5—H5A	0.9300
N1—C1	1.3890 (12)	C7—C8	1.4668 (12)
N2—C7	1.3554 (12)	C8—C9	1.4009 (13)
N2—C6	1.3798 (13)	C8—C13	1.4049 (13)
N2—H1N2	0.8188	C9—C10	1.3797 (16)
N3—C13	1.4460 (12)	C9—H9A	0.9300
N3—H1N3	0.8877	C10—C11	1.376 (2)
N3—H2N3	0.9418	C10—H10A	0.9300
C1—C6	1.3916 (14)	C11—C12	1.3816 (18)
C1—C2	1.3978 (15)	C11—H11A	0.9300
C2—C3	1.3830 (17)	C12—C13	1.3878 (14)
C2—H2A	0.9300	C12—H12A	0.9300
C3—C4	1.399 (2)		
N1—Zn1—N3	89.36 (3)	C3—C4—H4A	118.9
N1—Zn1—Cl2	114.83 (3)	C4—C5—C6	115.92 (12)
N3—Zn1—Cl2	112.74 (3)	C4—C5—H5A	122.0
N1—Zn1—Cl1	109.18 (3)	C6—C5—H5A	122.0
N3—Zn1—Cl1	109.27 (3)	N2—C6—C1	105.68 (8)
Cl2—Zn1—Cl1	117.844 (13)	N2—C6—C5	131.80 (10)
C7—N1—C1	106.79 (8)	C1—C6—C5	122.52 (10)
C7—N1—Zn1	121.20 (6)	N1—C7—N2	110.58 (8)
C1—N1—Zn1	129.89 (7)	N1—C7—C8	126.51 (8)
C7—N2—C6	108.33 (8)	N2—C7—C8	122.91 (8)
C7—N2—H1N2	130.1	C9—C8—C13	117.84 (9)
C6—N2—H1N2	121.5	C9—C8—C7	118.62 (9)
C13—N3—Zn1	116.24 (6)	C13—C8—C7	123.53 (8)
C13—N3—H1N3	113.1	C10—C9—C8	121.57 (10)
Zn1—N3—H1N3	107.2	C10—C9—H9A	119.2
C13—N3—H2N3	112.7	C8—C9—H9A	119.2
Zn1—N3—H2N3	99.8	C11—C10—C9	119.95 (10)
H1N3—N3—H2N3	106.7	C11—C10—H10A	120.0
N1—C1—C6	108.60 (8)	C9—C10—H10A	120.0

N1—C1—C2	130.32 (9)	C10—C11—C12	119.67 (11)
C6—C1—C2	121.07 (9)	C10—C11—H11A	120.2
C3—C2—C1	116.57 (11)	C12—C11—H11A	120.2
C3—C2—H2A	121.7	C11—C12—C13	121.13 (11)
C1—C2—H2A	121.7	C11—C12—H12A	119.4
C2—C3—C4	121.78 (12)	C13—C12—H12A	119.4
C2—C3—H3A	119.1	C12—C13—C8	119.79 (9)
C4—C3—H3A	119.1	C12—C13—N3	117.73 (9)
C5—C4—C3	122.13 (11)	C8—C13—N3	122.48 (8)
C5—C4—H4A	118.9		
N3—Zn1—N1—C7	37.88 (8)	C4—C5—C6—C1	0.90 (19)
Cl2—Zn1—N1—C7	152.76 (7)	C1—N1—C7—N2	-1.27 (11)
Cl1—Zn1—N1—C7	-72.34 (8)	Zn1—N1—C7—N2	163.70 (7)
N3—Zn1—N1—C1	-160.99 (9)	C1—N1—C7—C8	178.98 (9)
Cl2—Zn1—N1—C1	-46.11 (10)	Zn1—N1—C7—C8	-16.04 (14)
Cl1—Zn1—N1—C1	88.79 (9)	C6—N2—C7—N1	1.11 (12)
N1—Zn1—N3—C13	-47.06 (7)	C6—N2—C7—C8	-179.14 (9)
Cl2—Zn1—N3—C13	-163.85 (6)	N1—C7—C8—C9	170.17 (10)
Cl1—Zn1—N3—C13	63.07 (7)	N2—C7—C8—C9	-9.55 (15)
C7—N1—C1—C6	0.96 (11)	N1—C7—C8—C13	-10.56 (16)
Zn1—N1—C1—C6	-162.24 (7)	N2—C7—C8—C13	169.73 (10)
C7—N1—C1—C2	-177.97 (12)	C13—C8—C9—C10	1.27 (17)
Zn1—N1—C1—C2	18.82 (17)	C7—C8—C9—C10	-179.41 (11)
N1—C1—C2—C3	178.91 (12)	C8—C9—C10—C11	0.9 (2)
C6—C1—C2—C3	0.09 (18)	C9—C10—C11—C12	-1.9 (2)
C1—C2—C3—C4	0.0 (2)	C10—C11—C12—C13	0.7 (2)
C2—C3—C4—C5	0.3 (3)	C11—C12—C13—C8	1.5 (2)
C3—C4—C5—C6	-0.8 (2)	C11—C12—C13—N3	-177.82 (13)
C7—N2—C6—C1	-0.46 (11)	C9—C8—C13—C12	-2.43 (16)
C7—N2—C6—C5	178.79 (13)	C7—C8—C13—C12	178.29 (11)
N1—C1—C6—N2	-0.30 (11)	C9—C8—C13—N3	176.84 (10)
C2—C1—C6—N2	178.75 (10)	C7—C8—C13—N3	-2.44 (15)
N1—C1—C6—C5	-179.64 (11)	Zn1—N3—C13—C12	-143.37 (10)
C2—C1—C6—C5	-0.59 (18)	Zn1—N3—C13—C8	37.35 (12)
C4—C5—C6—N2	-178.24 (13)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H1N2···Cl1 ⁱ	0.82	2.56	3.3503 (9)	164
N3—H1N3···Cl1 ⁱⁱ	0.89	2.49	3.3753 (9)	174
N3—H2N3···Cl2 ⁱⁱⁱ	0.94	2.44	3.3015 (12)	153

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x, -y+2, -z$; (iii) $-x, y, -z+1/2$.