

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2*E*,4*E*)-1-(2-Hydroxyphenyl)-5-phenyl-penta-2,4-dien-1-oneW. A. Silva,^{a*} C. C. Gatto^a and G. R. Oliveira^b

^aInstituto de Química – IQ, Universidade de Brasília – UnB, Campus Universitário Darcy Ribeiro, Caixa postal 04478, Asa Norte Brasília DF, CEP 70904-970, Brazil, and ^bInstituto de Química – IQ, Universidade Federal de Goiás, CP 131, Campus Samambaia, Goiânia GO, CEP 74001-970, Brazil
Correspondence e-mail: wender@unb.br


Received 28 June 2011; accepted 26 July 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.050; wR factor = 0.134; data-to-parameter ratio = 15.5.

In the structure of the title chalcone, $\text{C}_{17}\text{H}_{14}\text{O}_2$, derived from cinnamaldehyde, the olefine group has a *trans* configuration. The molecular conformation is stabilized by an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen-bond interaction with graph-set motif $S(6)$.

Related literature

For the preparation, see: Lawrence *et al.* (2001). For related structures, see: Patil *et al.* (2007); Zhao *et al.* (2007). For standard bond lengths, see: Allen (2002). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related activity and structures, see: Dyrager *et al.* (2011); Jasinski *et al.* (2009); Ruan *et al.* (2011); Vencato *et al.* (2006).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{14}\text{O}_2$	$V = 2631.32$ (12) Å ³
$M_r = 250.28$	$Z = 8$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
$a = 10.9068$ (3) Å	$\mu = 0.08$ mm ⁻¹
$b = 7.9851$ (2) Å	$T = 296$ K
$c = 30.2131$ (7) Å	$0.39 \times 0.17 \times 0.09$ mm

Data collection

Bruker X8 SMART APEXII diffractometer	18214 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2008)	2691 independent reflections
$T_{\min} = 0.92$, $T_{\max} = 0.99$	1479 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	174 parameters
$wR(F^2) = 0.134$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\max} = 0.19$ e Å ⁻³
2691 reflections	$\Delta\rho_{\min} = -0.15$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{O2}$	0.82	1.81	2.5317 (19)	146

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors gratefully acknowledge support from the CNPq, FINEP-CT INFRA No. 0970/01 and the Instituto de Química - UnB.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2361).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bruker (2008). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Dyrager, C., Wickström, M., Fridén-Saxin, M., Friberg, A., Dahlén, K., Wallén, E. A. A., Gullbo, J., Grotli, M. & Luthman, K. (2011). *Bioorg. Med. Chem.* **19**, 2659–2665.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Jasinski, J. P., Butcher, R. J., Mayekar, A. N., Yathirajan, H. S. & Narayana, B. (2009). *J. Chem. Crystallogr.* **39**, 157–162.
 Lawrence, N. J., Rennison, D., McGown, A. T., Ducki, S., Gul, L. A., Hadfield, J. A. & Khan, N. (2001). *J. Comb. Chem.* **3**, 421–426.
 Patil, P. S., Teh, J. B.-J., Fun, H.-K., Razak, I. A. & Dharmaprakash, S. M. (2007). *Acta Cryst.* **E63**, o2122–o2123.
 Ruan, B., Lu, X., Tang, J., Wei, Y., Wang, X., Zhang, Y., Wang, L. & Zhu, H. (2011). *Bioorg. Med. Chem.* **19**, 2688–2695.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Vencato, I., Andrade, C. K. Z., Silva, W. A. & Lariucci, C. (2006). *Acta Cryst.* **E62**, o1033–o1035.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
 Zhao, B., Rong, Y.-Z. & Huang, W. (2007). *Acta Cryst.* **E63**, o2971.

supporting information

Acta Cryst. (2011). E67, o2210 [doi:10.1107/S160053681103025X]

(2E,4E)-1-(2-Hydroxyphenyl)-5-phenylpenta-2,4-dien-1-one

W. A. Silva, C. C. Gatto and G. R. Oliveira

S1. Comment

A large number of chalcones showed significant activity against many diseases. Some results from our laboratory showed that the synthesis of analogues of chalcones derived from cinnamaldehyde was often accompanied by increased in vitro bioactivity. However, the synthesis and analysis cytotoxic of these analogs of chalcones appears to be a largely unexplored field. With that, needs to make an accurate study of this new class of compounds in order to improve their structure-activity relationships.

The configuration of the olefinic group is trans, this was characterized for ¹H NMR where was showed the coupling J trans is around 15 Hz [C8—C9—C10—C11]. The presence of α - β -unsaturated ketone is indicated by the short O2—C11 and C9—C10 bond lengths of 1.245 (3) and 1.334 (3) Å, respectively, and the O2—C11—C10 and C9—C10—C11 bond angles of 118.8 (2)° and 121.8 (2)°, respectively. The bond distances are of normal values and are comparable with those found in related structures [Zhao *et al.* (2007); Patil *et al.* (2007)]. The molecular conformation is stabilized by one intramolecular O—H...O hydrogen-bond interaction with set graph motif S(6), (Bernstein *et al.*, 1995) [O1...O2 2.5285 (19) Å, O1—H1...O2 146.4°] (Fig. 2).

S2. Experimental

Compound was obtained by the aldol condensation of cinnamaldehyde, and 2-hydroxyacetophenone, using a method described previously [Lawrence *et al.* 2001]. Crystals were obtained from an EtOH solution (m.p. 428 K). Yield = 80% ¹H NMR: δ 6.92 (ddd, J = 7.8, 7.6 and 1.1 Hz, 1 H, 16-H), 7.01 (dd, J = 8.1 and 1.1 Hz, 1 H, 14-H), 7.05–7.07 (m, 2 H, 8, 7-H), 7.22 (d, J = 14.7 Hz, 1 H, 10-H), 7.34–7.42 (m, 3 H, 3,4,5-H), 7.49 (ddd, J = 8.1, 7.6 and 1.4 Hz, 1 H, 15-H), 7.52 (dd, J = 7.8 and 1.7 Hz, 2 H, 2,6-H), 7.67–7.76 (m, 1 H, 9-H), 7.85 (dd, J = 7.8 and 1.4 Hz, 1 H, 17-H), 12.88 (s, 1 H, OH). ¹³C NMR: δ = 118.6 (C-14), 118.8 (C-16), 120.0 (C-12), 123.5 (C-10), 126.7 (C-8), 127.4 (C-2,6), 128.9 (C-3,5), 129.4 (C-4), 129.5 (C-17), 135.9 (C-1), 136.2 (C-15), 142.9 (C-7), 145.4 (C-9), 163.5 (C-13), 194.0 (C11).

S3. Refinement

All H atoms were positioned geometrically and allowed to ride on their parent atoms, with d(C—H) = 0.93 Å and $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of showing the atomic labelling scheme. The anisotropic displacement parameters are at the 50% level. The dashed line indicates an intramolecular hydrogen bond, O1—H1···O2.

(2*E*,4*E*)-1-(2-Hydroxyphenyl)-5-phenylpenta-2,4-dien-1-one

Crystal data

$C_{17}H_{14}O_2$

$M_r = 250.28$

Orthorhombic, *Pbca*

$a = 10.9068 (3) \text{ \AA}$

$b = 7.9851 (2) \text{ \AA}$

$c = 30.2131 (7) \text{ \AA}$

$V = 2631.32 (12) \text{ \AA}^3$

$Z = 8$

$F(000) = 1056$

$D_x = 1.264 \text{ Mg m}^{-3}$

Melting point: 428 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1573 reflections

$\theta = 2.7\text{--}19.5^\circ$

$\mu = 0.08 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, yellow

$0.39 \times 0.17 \times 0.09 \text{ mm}$

Data collection

Bruker X8 SMART APEXII

diffractometer

Graphite monochromator

Detector resolution: $8.3333 \text{ pixels mm}^{-1}$

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.92$, $T_{\max} = 0.99$

18214 measured reflections

2691 independent reflections

1479 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.059$

$\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 1.4^\circ$

$h = -13 \rightarrow 11$

$k = -9 \rightarrow 9$

$l = -36 \rightarrow 37$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.134$

$S = 1.01$

2691 reflections

174 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0634P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.19 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.15 \text{ e \AA}^{-3}$

Extinction correction: *SHELXL*

Extinction coefficient: 0.0109 (11)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O2	0.09900 (13)	0.62840 (18)	1.00369 (4)	0.0625 (4)
O1	0.07689 (13)	0.4619 (2)	1.07455 (5)	0.0652 (5)
H1	0.0543	0.5068	1.0515	0.098*
C12	0.27179 (18)	0.5570 (2)	1.04643 (6)	0.0443 (5)
C13	0.19982 (19)	0.4740 (2)	1.07803 (6)	0.0488 (5)
C11	0.21235 (18)	0.6346 (2)	1.00794 (6)	0.0472 (5)
C9	0.23958 (18)	0.7440 (2)	0.93309 (6)	0.0511 (5)
H9	0.1581	0.7152	0.9281	0.061*
C1	0.33154 (19)	0.8908 (3)	0.81740 (6)	0.0532 (5)
C10	0.28508 (18)	0.7155 (2)	0.97335 (6)	0.0515 (5)
H10	0.3651	0.748	0.9796	0.062*
C17	0.39835 (19)	0.5612 (3)	1.05313 (7)	0.0567 (6)
H17	0.4476	0.6164	1.0327	0.068*
C8	0.30670 (19)	0.8155 (2)	0.89703 (6)	0.0535 (5)
H8	0.3844	0.8579	0.903	0.064*
C7	0.26596 (19)	0.8255 (2)	0.85581 (7)	0.0554 (6)
H7	0.1868	0.7866	0.8508	0.066*
C6	0.4439 (2)	0.9703 (3)	0.82097 (7)	0.0620 (6)
H6	0.4796	0.9844	0.8487	0.074*
C5	0.5037 (2)	1.0290 (3)	0.78378 (8)	0.0762 (7)
H5	0.5794	1.0815	0.7866	0.091*
C15	0.3789 (2)	0.4036 (3)	1.11921 (7)	0.0686 (6)
H15	0.4151	0.3511	1.1434	0.082*
C16	0.4517 (2)	0.4863 (3)	1.08888 (7)	0.0671 (6)
H16	0.5362	0.491	1.0928	0.081*
C14	0.2542 (2)	0.3976 (3)	1.11417 (7)	0.0627 (6)
H14	0.2062	0.3423	1.135	0.075*
C2	0.2808 (2)	0.8734 (3)	0.77533 (7)	0.0663 (7)
H2	0.2052	0.8209	0.7721	0.08*
C3	0.3414 (3)	0.9331 (3)	0.73832 (7)	0.0791 (8)
H3	0.3065	0.9205	0.7104	0.095*
C4	0.4522 (3)	1.0105 (4)	0.74271 (8)	0.0822 (8)
H4	0.4929	1.0507	0.7178	0.099*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O2	0.0432 (9)	0.0774 (10)	0.0669 (10)	-0.0019 (7)	-0.0019 (7)	0.0144 (7)
O1	0.0552 (10)	0.0791 (11)	0.0615 (10)	-0.0017 (8)	0.0075 (7)	0.0119 (8)
C12	0.0438 (13)	0.0442 (11)	0.0450 (11)	0.0031 (9)	0.0010 (9)	-0.0063 (9)
C13	0.0491 (13)	0.0487 (11)	0.0485 (11)	0.0033 (10)	0.0014 (10)	-0.0078 (9)
C11	0.0418 (12)	0.0468 (11)	0.0529 (12)	-0.0010 (9)	0.0037 (10)	-0.0051 (9)
C9	0.0454 (13)	0.0494 (12)	0.0585 (13)	0.0008 (10)	0.0042 (10)	-0.0005 (10)
C1	0.0542 (15)	0.0540 (12)	0.0515 (13)	0.0060 (11)	0.0009 (10)	0.0018 (9)
C10	0.0447 (13)	0.0557 (12)	0.0540 (12)	-0.0048 (10)	-0.0008 (10)	-0.0004 (10)
C17	0.0505 (14)	0.0619 (14)	0.0577 (13)	0.0023 (10)	-0.0012 (11)	-0.0059 (10)
C8	0.0468 (12)	0.0561 (12)	0.0577 (13)	-0.0039 (10)	0.0033 (10)	0.0043 (10)
C7	0.0499 (13)	0.0580 (13)	0.0583 (13)	-0.0028 (10)	-0.0025 (11)	0.0024 (10)
C6	0.0575 (15)	0.0729 (15)	0.0555 (14)	-0.0001 (12)	0.0010 (11)	0.0080 (11)
C5	0.0616 (16)	0.0920 (18)	0.0748 (17)	0.0004 (13)	0.0064 (13)	0.0201 (14)
C15	0.0813 (19)	0.0689 (14)	0.0555 (14)	0.0150 (13)	-0.0163 (13)	-0.0059 (12)
C16	0.0564 (15)	0.0764 (16)	0.0684 (16)	0.0061 (12)	-0.0148 (12)	-0.0087 (13)
C14	0.0792 (17)	0.0614 (14)	0.0476 (12)	0.0033 (12)	0.0012 (12)	-0.0001 (10)
C2	0.0660 (16)	0.0763 (16)	0.0566 (14)	0.0051 (12)	-0.0074 (12)	0.0013 (11)
C3	0.086 (2)	0.101 (2)	0.0504 (14)	0.0173 (16)	-0.0046 (13)	0.0067 (13)
C4	0.0790 (19)	0.107 (2)	0.0610 (17)	0.0177 (16)	0.0127 (14)	0.0238 (14)

Geometric parameters (Å, °)

O2—C11	1.244 (2)	C8—C7	1.324 (3)
O1—C13	1.348 (2)	C8—H8	0.93
O1—H1	0.82	C7—H7	0.93
C12—C17	1.395 (3)	C6—C5	1.381 (3)
C12—C13	1.402 (3)	C6—H6	0.93
C12—C11	1.469 (3)	C5—C4	1.370 (3)
C13—C14	1.384 (3)	C5—H5	0.93
C11—C10	1.463 (3)	C15—C14	1.369 (3)
C9—C10	1.333 (2)	C15—C16	1.381 (3)
C9—C8	1.431 (3)	C15—H15	0.93
C9—H9	0.93	C16—H16	0.93
C1—C6	1.385 (3)	C14—H14	0.93
C1—C2	1.393 (3)	C2—C3	1.383 (3)
C1—C7	1.460 (3)	C2—H2	0.93
C10—H10	0.93	C3—C4	1.364 (3)
C17—C16	1.365 (3)	C3—H3	0.93
C17—H17	0.93	C4—H4	0.93
C13—O1—H1	109.5	C8—C7—H7	116.4
C17—C12—C13	117.80 (18)	C1—C7—H7	116.4
C17—C12—C11	122.77 (18)	C5—C6—C1	120.6 (2)
C13—C12—C11	119.44 (18)	C5—C6—H6	119.7
O1—C13—C14	117.13 (19)	C1—C6—H6	119.7

O1—C13—C12	122.51 (18)	C4—C5—C6	120.5 (2)
C14—C13—C12	120.4 (2)	C4—C5—H5	119.7
O2—C11—C10	118.87 (17)	C6—C5—H5	119.7
O2—C11—C12	120.23 (17)	C14—C15—C16	120.9 (2)
C10—C11—C12	120.87 (17)	C14—C15—H15	119.5
C10—C9—C8	124.9 (2)	C16—C15—H15	119.5
C10—C9—H9	117.6	C17—C16—C15	119.3 (2)
C8—C9—H9	117.6	C17—C16—H16	120.3
C6—C1—C2	117.93 (19)	C15—C16—H16	120.3
C6—C1—C7	122.38 (18)	C15—C14—C13	119.9 (2)
C2—C1—C7	119.7 (2)	C15—C14—H14	120.1
C9—C10—C11	121.70 (19)	C13—C14—H14	120.1
C9—C10—H10	119.2	C3—C2—C1	120.9 (2)
C11—C10—H10	119.2	C3—C2—H2	119.5
C16—C17—C12	121.7 (2)	C1—C2—H2	119.5
C16—C17—H17	119.1	C4—C3—C2	120.0 (2)
C12—C17—H17	119.1	C4—C3—H3	120.0
C7—C8—C9	124.6 (2)	C2—C3—H3	120.0
C7—C8—H8	117.7	C3—C4—C5	120.0 (2)
C9—C8—H8	117.7	C3—C4—H4	120.0
C8—C7—C1	127.2 (2)	C5—C4—H4	120.0
C17—C12—C13—O1	-179.60 (17)	C6—C1—C7—C8	-7.7 (3)
C11—C12—C13—O1	0.2 (3)	C2—C1—C7—C8	172.1 (2)
C17—C12—C13—C14	-0.8 (3)	C2—C1—C6—C5	-0.5 (3)
C11—C12—C13—C14	178.95 (17)	C7—C1—C6—C5	179.4 (2)
C17—C12—C11—O2	-179.47 (18)	C1—C6—C5—C4	0.5 (4)
C13—C12—C11—O2	0.8 (3)	C12—C17—C16—C15	0.3 (3)
C17—C12—C11—C10	2.5 (3)	C14—C15—C16—C17	-0.9 (3)
C13—C12—C11—C10	-177.20 (17)	C16—C15—C14—C13	0.6 (3)
C8—C9—C10—C11	-177.06 (18)	O1—C13—C14—C15	179.10 (18)
O2—C11—C10—C9	-17.4 (3)	C12—C13—C14—C15	0.2 (3)
C12—C11—C10—C9	160.63 (18)	C6—C1—C2—C3	0.3 (3)
C13—C12—C17—C16	0.5 (3)	C7—C1—C2—C3	-179.6 (2)
C11—C12—C17—C16	-179.20 (18)	C1—C2—C3—C4	-0.1 (4)
C10—C9—C8—C7	171.7 (2)	C2—C3—C4—C5	0.0 (4)
C9—C8—C7—C1	-177.45 (19)	C6—C5—C4—C3	-0.2 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...O2	0.82	1.81	2.5317 (19)	146