

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Amino-1-(4-bromophenyl)-9,10-dihydrophenanthrene-2,4-dicarbonitrile


 Abdullah M. Asiri,^{a,b} Hassan M. Faidallah,^a
 Abdulrahman O. Al-Youbi^a and Seik Weng Ng^{c,*}
^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, ^bCenter of Excellence for Advanced Materials Research, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
 Correspondence e-mail: seikweng@um.edu.my

Received 18 September 2011; accepted 20 September 2011

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.021; wR factor = 0.056; data-to-parameter ratio = 9.0.

In the title compound, $\text{C}_{22}\text{H}_{14}\text{BrN}_3$, the fused-ring system is buckled owing to the ethylene linkage in the central ring; the two flanking aromatic rings are twisted by $25.9(1)^\circ$ with respect to each other. The phenyl ring is twisted by $77.0(1)^\circ$ relative to the amino- and cyano-bearing aromatic ring. In the crystal, adjacent molecules are linked by two $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds, generating a zigzag chain along $[101]$.

Related literature

 For two related compounds, see: Asiri *et al.* (2011a,b).


Experimental

Crystal data

$\text{C}_{22}\text{H}_{14}\text{BrN}_3$	$b = 16.2557(3)$ Å
$M_r = 400.27$	$c = 9.7945(4)$ Å
Monoclinic, Cc	$\beta = 127.546(6)^\circ$
$a = 13.7683(5)$ Å	$V = 1738.07(17)$ Å ³

$Z = 4$	$T = 100$ K
Cu $K\alpha$ radiation	$0.20 \times 0.20 \times 0.20$ mm
$\mu = 3.29$ mm ⁻¹	

Data collection

Agilent SuperNova Dual diffractometer with Atlas detector	2976 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2010)	2195 independent reflections
$T_{\min} = 0.559$, $T_{\max} = 0.559$	2187 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.012$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.056$	$\Delta\rho_{\text{max}} = 0.22$ e Å ⁻³
$S = 1.08$	$\Delta\rho_{\text{min}} = -0.61$ e Å ⁻³
2195 reflections	Absolute structure: Flack (Flack, 1983), 482 Friedel pairs
243 parameters	Flack parameter: $-0.024(14)$
2 restraints	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}1\cdots\text{N}1^{\text{i}}$	0.93 (3)	2.23 (3)	3.097 (3)	155 (3)
$\text{N}2-\text{H}2\cdots\text{N}3^{\text{ii}}$	0.88 (4)	2.54 (4)	3.307 (3)	147 (3)

 Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdulaziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5646).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Asiri, A. M., Al-Youbi, A. O., Faidallah, H. M., Ng, S. W. & Tiekink, E. R. T. (2011a). *Acta Cryst.* **E67**, o2438.
 Asiri, A. M., Al-Youbi, A. O., Faidallah, H. M., Ng, S. W. & Tiekink, E. R. T. (2011b). *Acta Cryst.* **E67**, o2449.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, o2745 [https://doi.org/10.1107/S1600536811038517]

3-Amino-1-(4-bromophenyl)-9,10-dihydrophenanthrene-2,4-dicarbonitrile

Abdullah M. Asiri, Hassan M. Faidallah, Abdulrahman O. Al-Youbi and Seik Weng Ng

S1. Comment

2-Amino-4-aryl-5,6-dihydrobenzoquinoline-3-carbonitrile is synthesized from the reaction of the α -substituted cinnamitrile, $C_6H_5CH=C(CN)_2$, with α -tetralone in a reaction that is catalyzed by ammonium acetate. The synthesis when conducted under microwave irradiation leads to an improved yield. In previous studies, we obtained instead di-carbonitrile substituted dihydrophenanthrenes (3-amino-1-(4-methoxyphenyl)-9,10-dihydrophenanthrene-2,4-dicarbonitrile and 3-amino-1-(2*H*-1,3-benzodioxol-5-yl)-9,10-dihydrophenanthrene-2,4-dicarbonitrile) with 4-methoxybenzaldehyde and piperonaldehyde in syntheses that differed slightly from the reported ones as we used substituted benzaldehydes, α -tetralone and ethyl cyanoacetate along with a molar excess of ammonium acetate (Asiri *et al.*, 2011*a*; 2011*b*). The use of 4-bromobenzaldehyde furnishes the corresponding 4-bromophenyl analog (Scheme I, Fig. 1). The fused-ring system is buckled owing to the ethylene linkage in the central ring; the two flanking aromatic rings are twisted by 25.9 (1) °. Relative to the amino- and cyano-bearing aromatic ring, the phenyl ring is twisted by 77.0 (1) °. Adjacent molecules are linked by two N–H···N hydrogen bonds to generate a chain along [1 0 1] (Table 1).

S2. Experimental

4-Bromobenzaldehyde (1.85 g, 10 mmol), 1-tetralone (1.46 g, 10 mmol), malononitrile (0.66 g, 10 mmol) and ammonium acetate (6.2 g, 80 mmol) in absolute ethanol (50 ml) were heated for 6 h. The mixture was allowed to cool, and the precipitate was collected, washed with water, dried and then recrystallized from ethanol; m.p. 517–518.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [$C-H$ 0.95 to 0.99 Å, $U_{iso}(H)$ 1.2–1.5 $U_{eq}(C)$] and were included in the refinement in the riding model approximation. The amino H atoms were located in a difference Fourier map and were refined freely.

The Flack parameter was refined from 482 Friedel pairs; although the Friedel coverage is low (27%), the Flack parameter was reliably refined owing to the heavy atom.




Figure 1

Anisotropic displacement ellipsoid plot (Barbour, 2001) of $C_{22}H_{14}N_3Br$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

3-Amino-1-(4-bromophenyl)-9,10-dihydrophenanthrene-2,4-dicarbonitrile

Crystal data

$C_{22}H_{14}BrN_3$

$M_r = 400.27$

Monoclinic, Cc

Hall symbol: $C -2yc$

$a = 13.7683 (5) \text{ \AA}$

$b = 16.2557 (3) \text{ \AA}$

$c = 9.7945 (4) \text{ \AA}$

$\beta = 127.546 (6)^\circ$

$V = 1738.07 (17) \text{ \AA}^3$

$Z = 4$

$F(000) = 808$

$D_x = 1.530 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$

Cell parameters from 2539 reflections

$\theta = 4.9\text{--}74.2^\circ$

$\mu = 3.29 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Prism, orange

$0.20 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Agilent SuperNova Dual

diffractometer with Atlas detector

Radiation source: SuperNova (Cu) X-ray

Source

Mirror monochromator

Detector resolution: $10.4041 \text{ pixels mm}^{-1}$

ω scan

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2010)

$T_{\min} = 0.559, T_{\max} = 0.559$

2976 measured reflections

2195 independent reflections

2187 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$

$\theta_{\max} = 74.4^\circ, \theta_{\min} = 4.9^\circ$

$h = -17 \rightarrow 16$

$k = -11 \rightarrow 20$

$l = -11 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.021$

$wR(F^2) = 0.056$

$S = 1.08$

2195 reflections

243 parameters

2 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0393P)^2 + 0.2403P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.61 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (Flack, 1983), 482
 Friedel pairs
 Absolute structure parameter: -0.024 (14)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.00008 (3)	0.303704 (13)	0.00088 (3)	0.02570 (9)
N1	0.8004 (2)	0.69430 (12)	1.0721 (3)	0.0189 (4)
N2	0.5320 (2)	0.70911 (13)	0.6833 (3)	0.0173 (4)
H1	0.460 (3)	0.739 (2)	0.618 (4)	0.018 (7)*
H2	0.597 (3)	0.738 (2)	0.761 (5)	0.024 (8)*
N3	0.2436 (2)	0.64780 (13)	0.3201 (3)	0.0245 (5)
C1	0.5286 (2)	0.36690 (15)	0.8140 (3)	0.0172 (5)
H1A	0.4753	0.3368	0.7036	0.021*
H1B	0.4964	0.3581	0.8797	0.021*
C2	0.6591 (2)	0.33350 (15)	0.9166 (3)	0.0202 (5)
H2A	0.6604	0.2748	0.9447	0.024*
H2B	0.6891	0.3377	0.8472	0.024*
C3	0.7407 (2)	0.38245 (14)	1.0797 (3)	0.0168 (5)
C4	0.8329 (2)	0.34500 (16)	1.2344 (4)	0.0217 (5)
H4	0.8462	0.2875	1.2364	0.026*
C5	0.9058 (2)	0.38981 (17)	1.3858 (3)	0.0221 (5)
H5	0.9700	0.3635	1.4897	0.026*
C6	0.8845 (2)	0.47317 (16)	1.3845 (3)	0.0209 (5)
H6	0.9324	0.5040	1.4883	0.025*
C7	0.7926 (2)	0.51166 (15)	1.2307 (3)	0.0167 (5)
H7	0.7780	0.5687	1.2309	0.020*
C8	0.7217 (2)	0.46789 (13)	1.0763 (3)	0.0144 (4)
C9	0.6244 (2)	0.50717 (14)	0.9095 (3)	0.0130 (4)
C10	0.6259 (2)	0.59116 (14)	0.8755 (3)	0.0131 (4)
C11	0.5282 (2)	0.62902 (15)	0.7197 (3)	0.0133 (4)
C12	0.4277 (2)	0.57892 (14)	0.5984 (3)	0.0142 (4)
C13	0.4282 (2)	0.49398 (15)	0.6262 (3)	0.0162 (5)
C14	0.5264 (2)	0.45766 (14)	0.7787 (3)	0.0166 (4)
C15	0.7262 (2)	0.64517 (14)	0.9918 (3)	0.0137 (4)
C16	0.3254 (2)	0.61565 (15)	0.4420 (3)	0.0174 (5)
C17	0.3244 (2)	0.44469 (14)	0.4820 (3)	0.0143 (4)
C18	0.3423 (2)	0.39889 (15)	0.3792 (3)	0.0183 (5)
H18	0.4216	0.3967	0.4075	0.022*
C19	0.2471 (2)	0.35638 (14)	0.2367 (3)	0.0185 (5)
H19	0.2601	0.3253	0.1670	0.022*
C20	0.1321 (2)	0.36031 (14)	0.1981 (3)	0.0166 (5)
C21	0.1114 (2)	0.40498 (17)	0.2974 (3)	0.0238 (5)

H21	0.0318	0.4071	0.2682	0.029*
C22	0.2085 (2)	0.44712 (17)	0.4412 (3)	0.0212 (5)
H22	0.1953	0.4776	0.5114	0.025*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.02085 (13)	0.02220 (13)	0.01646 (13)	-0.00648 (12)	0.00229 (10)	-0.00716 (12)
N1	0.0175 (11)	0.0161 (11)	0.0163 (11)	0.0000 (8)	0.0068 (10)	0.0001 (8)
N2	0.0141 (10)	0.0118 (9)	0.0155 (11)	0.0002 (8)	0.0036 (9)	0.0002 (8)
N3	0.0210 (11)	0.0174 (10)	0.0201 (11)	0.0007 (9)	0.0048 (10)	0.0001 (9)
C1	0.0180 (12)	0.0132 (11)	0.0159 (11)	-0.0025 (9)	0.0079 (10)	-0.0014 (9)
C2	0.0217 (12)	0.0143 (11)	0.0213 (13)	-0.0016 (10)	0.0114 (11)	-0.0011 (10)
C3	0.0180 (12)	0.0132 (11)	0.0187 (12)	-0.0004 (9)	0.0110 (11)	0.0018 (9)
C4	0.0171 (11)	0.0202 (12)	0.0250 (13)	0.0042 (10)	0.0115 (11)	0.0090 (11)
C5	0.0191 (12)	0.0238 (13)	0.0197 (12)	0.0009 (10)	0.0100 (11)	0.0105 (10)
C6	0.0184 (11)	0.0261 (13)	0.0142 (11)	-0.0025 (10)	0.0079 (10)	0.0019 (10)
C7	0.0159 (10)	0.0162 (11)	0.0165 (11)	-0.0016 (9)	0.0091 (10)	0.0009 (9)
C8	0.0127 (10)	0.0134 (11)	0.0160 (11)	-0.0005 (9)	0.0081 (9)	0.0024 (9)
C9	0.0121 (11)	0.0131 (10)	0.0138 (11)	0.0014 (9)	0.0079 (10)	-0.0001 (9)
C10	0.0126 (10)	0.0133 (10)	0.0129 (10)	-0.0014 (9)	0.0075 (9)	-0.0030 (9)
C11	0.0130 (11)	0.0132 (10)	0.0140 (11)	-0.0003 (9)	0.0084 (10)	-0.0019 (9)
C12	0.0136 (10)	0.0140 (11)	0.0114 (11)	0.0009 (8)	0.0057 (9)	0.0006 (8)
C13	0.0146 (11)	0.0131 (11)	0.0157 (11)	-0.0013 (9)	0.0066 (10)	-0.0029 (10)
C14	0.0173 (10)	0.0139 (11)	0.0165 (11)	-0.0023 (9)	0.0093 (10)	-0.0007 (9)
C15	0.0140 (11)	0.0116 (10)	0.0123 (11)	0.0025 (9)	0.0063 (10)	0.0007 (9)
C16	0.0173 (11)	0.0133 (10)	0.0176 (11)	-0.0026 (9)	0.0085 (10)	-0.0045 (10)
C17	0.0149 (10)	0.0104 (10)	0.0121 (10)	-0.0007 (9)	0.0055 (9)	0.0000 (9)
C18	0.0143 (11)	0.0174 (11)	0.0193 (12)	-0.0003 (9)	0.0082 (10)	-0.0023 (10)
C19	0.0208 (11)	0.0168 (12)	0.0167 (11)	-0.0005 (10)	0.0108 (10)	-0.0031 (10)
C20	0.0165 (11)	0.0115 (10)	0.0115 (11)	-0.0049 (9)	0.0032 (9)	-0.0006 (9)
C21	0.0161 (11)	0.0328 (14)	0.0193 (12)	-0.0061 (11)	0.0091 (10)	-0.0056 (11)
C22	0.0187 (12)	0.0270 (13)	0.0180 (12)	-0.0043 (10)	0.0111 (10)	-0.0071 (10)

Geometric parameters (Å, °)

Br1—C20	1.898 (2)	C7—H7	0.9500
N1—C15	1.149 (3)	C8—C9	1.485 (3)
N2—C11	1.359 (3)	C9—C10	1.408 (3)
N2—H1	0.93 (3)	C9—C14	1.415 (3)
N2—H2	0.88 (4)	C10—C11	1.420 (3)
N3—C16	1.152 (4)	C10—C15	1.435 (3)
C1—C14	1.511 (3)	C11—C12	1.410 (3)
C1—C2	1.528 (4)	C12—C13	1.407 (3)
C1—H1A	0.9900	C12—C16	1.434 (3)
C1—H1B	0.9900	C13—C14	1.395 (3)
C2—C3	1.503 (3)	C13—C17	1.489 (3)
C2—H2A	0.9900	C17—C22	1.389 (3)

C2—H2B	0.9900	C17—C18	1.390 (3)
C3—C4	1.390 (4)	C18—C19	1.384 (3)
C3—C8	1.410 (3)	C18—H18	0.9500
C4—C5	1.387 (4)	C19—C20	1.387 (3)
C4—H4	0.9500	C19—H19	0.9500
C5—C6	1.385 (4)	C20—C21	1.376 (4)
C5—H5	0.9500	C21—C22	1.395 (4)
C6—C7	1.392 (4)	C21—H21	0.9500
C6—H6	0.9500	C22—H22	0.9500
C7—C8	1.395 (3)		
C11—N2—H1	119 (2)	C9—C10—C11	122.2 (2)
C11—N2—H2	117 (2)	C9—C10—C15	123.5 (2)
H1—N2—H2	115 (3)	C11—C10—C15	114.3 (2)
C14—C1—C2	110.4 (2)	N2—C11—C12	120.5 (2)
C14—C1—H1A	109.6	N2—C11—C10	122.3 (2)
C2—C1—H1A	109.6	C12—C11—C10	117.1 (2)
C14—C1—H1B	109.6	C11—C12—C13	121.2 (2)
C2—C1—H1B	109.6	C11—C12—C16	118.7 (2)
H1A—C1—H1B	108.1	C13—C12—C16	120.1 (2)
C3—C2—C1	109.2 (2)	C14—C13—C12	120.6 (2)
C3—C2—H2A	109.8	C14—C13—C17	122.1 (2)
C1—C2—H2A	109.8	C12—C13—C17	117.1 (2)
C3—C2—H2B	109.8	C13—C14—C9	119.7 (2)
C1—C2—H2B	109.8	C13—C14—C1	121.9 (2)
H2A—C2—H2B	108.3	C9—C14—C1	118.2 (2)
C4—C3—C8	119.3 (2)	N1—C15—C10	173.1 (2)
C4—C3—C2	121.4 (2)	N3—C16—C12	177.3 (3)
C8—C3—C2	119.3 (2)	C22—C17—C18	119.1 (2)
C3—C4—C5	121.4 (2)	C22—C17—C13	121.9 (2)
C3—C4—H4	119.3	C18—C17—C13	118.9 (2)
C5—C4—H4	119.3	C19—C18—C17	121.4 (2)
C6—C5—C4	119.5 (2)	C19—C18—H18	119.3
C6—C5—H5	120.2	C17—C18—H18	119.3
C4—C5—H5	120.2	C18—C19—C20	118.3 (2)
C5—C6—C7	119.8 (2)	C18—C19—H19	120.9
C5—C6—H6	120.1	C20—C19—H19	120.9
C7—C6—H6	120.1	C21—C20—C19	121.7 (2)
C6—C7—C8	121.2 (2)	C21—C20—Br1	119.50 (19)
C6—C7—H7	119.4	C19—C20—Br1	118.77 (18)
C8—C7—H7	119.4	C20—C21—C22	119.3 (2)
C7—C8—C3	118.6 (2)	C20—C21—H21	120.4
C7—C8—C9	122.7 (2)	C22—C21—H21	120.4
C3—C8—C9	118.7 (2)	C17—C22—C21	120.2 (2)
C10—C9—C14	118.7 (2)	C17—C22—H22	119.9
C10—C9—C8	122.9 (2)	C21—C22—H22	119.9
C14—C9—C8	118.3 (2)		

C14—C1—C2—C3	56.3 (3)	C10—C11—C12—C16	177.9 (2)
C1—C2—C3—C4	141.2 (2)	C11—C12—C13—C14	2.9 (3)
C1—C2—C3—C8	-37.6 (3)	C16—C12—C13—C14	-179.2 (2)
C8—C3—C4—C5	0.6 (4)	C11—C12—C13—C17	-173.1 (2)
C2—C3—C4—C5	-178.1 (2)	C16—C12—C13—C17	4.9 (3)
C3—C4—C5—C6	1.8 (4)	C12—C13—C14—C9	2.4 (3)
C4—C5—C6—C7	-1.9 (4)	C17—C13—C14—C9	178.1 (2)
C5—C6—C7—C8	-0.6 (4)	C12—C13—C14—C1	178.8 (2)
C6—C7—C8—C3	3.0 (3)	C17—C13—C14—C1	-5.5 (4)
C6—C7—C8—C9	-178.9 (2)	C10—C9—C14—C13	-6.0 (3)
C4—C3—C8—C7	-3.0 (3)	C8—C9—C14—C13	174.4 (2)
C2—C3—C8—C7	175.8 (2)	C10—C9—C14—C1	177.4 (2)
C4—C3—C8—C9	178.9 (2)	C8—C9—C14—C1	-2.2 (3)
C2—C3—C8—C9	-2.3 (3)	C2—C1—C14—C13	145.4 (2)
C7—C8—C9—C10	26.4 (3)	C2—C1—C14—C9	-38.2 (3)
C3—C8—C9—C10	-155.6 (2)	C14—C13—C17—C22	108.9 (3)
C7—C8—C9—C14	-154.1 (2)	C12—C13—C17—C22	-75.2 (3)
C3—C8—C9—C14	24.0 (3)	C14—C13—C17—C18	-75.0 (3)
C14—C9—C10—C11	4.7 (3)	C12—C13—C17—C18	100.9 (3)
C8—C9—C10—C11	-175.7 (2)	C22—C17—C18—C19	0.6 (4)
C14—C9—C10—C15	-173.7 (2)	C13—C17—C18—C19	-175.6 (2)
C8—C9—C10—C15	5.8 (3)	C17—C18—C19—C20	-0.1 (4)
C9—C10—C11—N2	-176.9 (2)	C18—C19—C20—C21	0.0 (4)
C15—C10—C11—N2	1.7 (3)	C18—C19—C20—Br1	179.03 (17)
C9—C10—C11—C12	0.3 (3)	C19—C20—C21—C22	-0.4 (4)
C15—C10—C11—C12	178.9 (2)	Br1—C20—C21—C22	-179.4 (2)
N2—C11—C12—C13	173.1 (2)	C18—C17—C22—C21	-0.9 (4)
C10—C11—C12—C13	-4.2 (3)	C13—C17—C22—C21	175.1 (2)
N2—C11—C12—C16	-4.8 (3)	C20—C21—C22—C17	0.8 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H1...N1 ⁱ	0.93 (3)	2.23 (3)	3.097 (3)	155 (3)
N2—H2...N3 ⁱⁱ	0.88 (4)	2.54 (4)	3.307 (3)	147 (3)

Symmetry codes: (i) $x-1/2, -y+3/2, z-1/2$; (ii) $x+1/2, -y+3/2, z+1/2$.