

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 1,4-Bis(1,1-dimethylpropyl)-2,5-dimethoxybenzene

Kiichi Amimoto

Department of Science Education, Graduate School of Education, Hiroshima University, 1-1-1 Kagamiyama, Higashi-Hiroshima, Hiroshima, Japan  
Correspondence e-mail: kamimo@hiroshima-u.ac.jp


Received 6 September 2011; accepted 15 September 2011

Key indicators: single-crystal X-ray study;  $T = 90$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.114; data-to-parameter ratio = 12.5.

The title compound,  $\text{C}_{18}\text{H}_{30}\text{O}_2$ , was prepared by Friedel-Crafts alkylation of 1,4-dimethoxybenzene with 2-methyl-2-butanol. The complete molecule is generated by the application of a crystallographic centre of inversion. The two methoxy groups are oriented in the same plane of the aromatic ring [ $\text{C}-\text{C}-\text{O}-\text{C}$  torsion angle =  $9.14$  ( $16$ )°]. While one methyl group of the *tert*-pentyl substituent is coplanar with the benzene ring [ $\text{C}-\text{C}-\text{C}-\text{C} = 0.45$  ( $15$ )°] and lies towards the less-hindered H atom, the other methyl and ethyl groups are directed to either side of the benzene ring [ $\text{C}-\text{C}-\text{C}-\text{C}$  torsion angles =  $118.78$  ( $12$ ) and  $59.11$  ( $14$ )°, respectively]. In the crystal, the hydrophobic molecules pack to form a brick-wall-like architecture.

## Related literature

For the synthesis of the title compound, see: Polito *et al.* (2010) and for the synthesis of the analogous compound, 1,4-di-*tert*-butyl-2,5-dimethoxybenzene, see: Williamson *et al.* (2006). For the unique crystal growth of 1,4-di-*tert*-butyl-2,5-dimethoxybenzene, see: Blatchly & Hartshorne (1966). For the crystal structure of 1,4-di-*tert*-butyl-2,5-dimethoxybenzene, see: Rosokha & Kochi (2007).



## Experimental

## Crystal data

 $\text{C}_{18}\text{H}_{30}\text{O}_2$  $M_r = 278.42$ 

Triclinic,  $P\bar{1}$   
 $a = 6.456$  (5) Å  
 $b = 6.551$  (5) Å  
 $c = 10.800$  (5) Å  
 $\alpha = 93.120$  (5)°  
 $\beta = 105.950$  (5)°  
 $\gamma = 108.460$  (5)°

$V = 411.5$  (5) Å<sup>3</sup>  
 $Z = 1$   
Mo  $K\alpha$  radiation  
 $\mu = 0.07$  mm<sup>-1</sup>  
 $T = 90$  K  
 $0.4 \times 0.2 \times 0.1$  mm

## Data collection

Bruker APEXII CCD area-detector diffractometer  
Absorption correction: empirical (using intensity measurements) (*SADABS*; Bruker, 2009)  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.993$   
2473 measured reflections  
1888 independent reflections  
1698 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.013$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.114$   
 $S = 1.10$   
1888 reflections

151 parameters  
All H-atom parameters refined  
 $\Delta\rho_{\text{max}} = 0.40$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.22$  e Å<sup>-3</sup>

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *publCIF* (Westrip, 2010).

This work was partially supported by a Grant-in-Aid for Young Scientists (B) (grant No. 23700956) and Grant-in-Aid for Scientific Research (C) (grant No. 22300272) from the Japan Society for the Promotion of Science (JSPS). The measurements of X-ray crystallographic, <sup>1</sup>H NMR and APCI-MS data were performed at the Natural Science Center for Basic Research and Development (N-BARD), Hiroshima University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2787).

## References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.  
Blatchly, J. M. & Hartshorne, N. H. (1966). *Trans. Faraday Soc.* **62**, 512–518.  
Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.  
Polito, V., Hamann, C. S. & Rhile, I. J. (2010). *J. Chem. Educ.* **87**, 969–967.  
Rosokha, S. V. & Kochi, J. K. (2007). *J. Am. Chem. Soc.* **129**, 3683–3697.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.  
Williamson, K. L., Minard, R. & Masters, K. M. (2006). *Macroscale and Microscale Organic Experiments*, 5th ed., ch. 29, pp. 436–450. Boston: Houghton Mifflin.

## supporting information

*Acta Cryst.* (2011). E67, o2708 [https://doi.org/10.1107/S160053681103772X]

**1,4-Bis(1,1-dimethylpropyl)-2,5-dimethoxybenzene****Kiichi Amimoto****S1. Comment**

Electrophilic aromatic substitution is one of the key reactions in organic chemistry. The Friedel-Crafts alkylation of 1,4-dimethoxybenzene with *tert*-butyl alcohol is a popular introductory organic laboratory experiment to illustrate the features of electrophilic aromatic substitution (Williamson *et al.*, 2006). The product, 1,4-di-*tert*-butyl-2,5-dimethoxybenzene, (I), has also been reported to exhibit a dramatic change of shape during crystal growth (Blatchly and Hartshorne, 1966). Recently, a synthesis involving electrophilic aromatic substitution coupled with a Wagner-Meerwein rearrangement using 1,4-dimethoxybenzene with 2-methyl-2-butanol was reported (Polito *et al.*, 2010). However, the product, 1,4-bis(1,1-dimethylpropyl)-2,5-dimethoxybenzene (II), has no dramatic crystal growth behaviour. To obtain the perspectives on both development of sophisticated laboratory activity in organic chemistry and the relationship between crystal growth and molecular structure, the X-ray diffraction analysis of the title compound (II) was performed, and the structural features of (I) and (II) discussed in this article. The crystal structure of (I) has already been reported (Rosokha and Kochi, 2007).

Compound (I) crystallizes in the monoclinic space group  $P2_1/c$ . The methoxy groups are inclined at  $33.9(1)^\circ$  to the benzene plane. The methyl C—H of the methoxy group points to the adjacent benzene ring in a face-on-edge manner, in which the distance between the methyl carbon and the  $\pi$ -plane is  $3.47(1) \text{ \AA}$ . The crystal packing exhibits a herringbone structure. Compound (II) crystallizes in the triclinic space group  $P\bar{1}$  (Fig. 1). The asymmetric unit of (II) contains half a molecule with the complete molecule being generated by a centre of inversion located in the benzene ring. In contrast to (I), the methoxy groups lie on almost the same plane as the benzene ring, where the dihedral angle is  $4.9(1)^\circ$ . The ethyl chains of two *tert*-pentyl groups stand to one side of the benzene plane in an *anti*-orientation, where the dihedral angle is  $83.8(2)^\circ$ . Since the ethyl chain and methyl group of *tert*-pentyl group and methoxy group are aggregated by hydrophobic interactions, each molecule is closely arranged in the brick-wall fashion (Fig. 2). Compared to (II) the packing mode of (I) seems to be relatively loose, resulting the expression of dynamic crystal growth of (I).


**S2. Experimental**

In a 50-ml Erlenmeyer flask were placed 690 mg (5 mmol) of 1,4-dimethoxybenzene, 1.3 g (1.5 mmol) of 2-methyl-2-butanol, and 1.5 ml of glacial acetic acid. The Erlenmeyer flask was immersed in an ice-water bath to cool the solution below 278 K. Concentrated sulfuric acid (5 ml) was added dropwise into the vigorously stirred reaction mixture so as not to exceed the temperature of 283 K. When all the sulfuric acid was added, the mixture was stirred at room temperature for 10 minutes. The Erlenmeyer flask was immersed again in an ice-water bath, and then a sufficient amount of ice-cold water was added into the reaction mixture to quench the reaction and isolate the product. The white solid was filtered off on a small Büchner funnel and washed with a small amount of ice-cold ethanol. Single crystals were obtained by recrystallization from its ethanol solution; *M.pt.* 382–383 K.  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.72 (s, 2 H), 3.76 (s, 6 H), 1.77 (q, 4 H), 1.29 (s, 12 H), 0.62 (q, 6 H) p.p.m. IR (KBr): 2962, 2933, 2868, 1506, 1481, 1392, 1369, 1209, 1128, 1041,

865  $\text{cm}^{-1}$ . APCI-MS:  $m/z$   $[\text{M}-\text{H}]^+ = 277.21621$ .


### S3. Refinement

All H atoms were found in a difference Fourier map and refined isotropically.



**Figure 1**

The molecular structure of the title compound, showing the atom-numbering scheme and displacement ellipsoids at the 50% probability level for non-H atoms. Atoms marked with <sup>i</sup> are at the symmetry positions 1 - x, 1 - y, 1 - z.



**Figure 2**

The crystal packing of the title compound showing the brick-wall packing arrangement. All H atoms are omitted for clarity.

### 1,4-Bis(1,1-dimethylpropyl)-2,5-dimethoxybenzene

#### Crystal data

$C_{18}H_{30}O_2$

$M_r = 278.42$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 6.456\ (5)\ \text{\AA}$

$b = 6.551\ (5)\ \text{\AA}$

$c = 10.800\ (5)\ \text{\AA}$

$\alpha = 93.120\ (5)^\circ$

$\beta = 105.950\ (5)^\circ$

$\gamma = 108.460\ (5)^\circ$

$V = 411.5\ (5)\ \text{\AA}^3$

$Z = 1$

$F(000) = 154$

$D_x = 1.123\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71069\ \text{\AA}$

Cell parameters from 1654 reflections

$\theta = 3.3\text{--}28.8^\circ$

$\mu = 0.07\ \text{mm}^{-1}$

$T = 90\ \text{K}$

Plate, colourless

$0.4 \times 0.2 \times 0.1\ \text{mm}$

#### Data collection

Bruker APEXII CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution:  $8.333\ \text{pixels mm}^{-1}$

$\varphi$  and  $\omega$  scan

Absorption correction: empirical (using  
intensity measurements)

(*SADABS*; Bruker, 2009)

$T_{\min} = 0.972$ ,  $T_{\max} = 0.993$

2473 measured reflections

1888 independent reflections

1698 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.013$

$\theta_{\max} = 29.0^\circ$ ,  $\theta_{\min} = 2.0^\circ$

$h = -8 \rightarrow 5$

$k = -8 \rightarrow 8$

$l = -12 \rightarrow 14$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.114$   
 $S = 1.10$   
 1888 reflections  
 151 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 All H-atom parameters refined  
 $w = 1/[\sigma^2(F_o^2) + (0.0566P)^2 + 0.1363P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.66045 (17)	0.40800 (16)	0.56010 (10)	0.0142 (2)
C2	0.54020 (18)	0.34722 (16)	0.42749 (10)	0.0146 (2)
C3	0.37620 (17)	0.43475 (16)	0.36311 (10)	0.0134 (2)
C4	0.24450 (18)	0.36052 (16)	0.21713 (10)	0.0149 (2)
C5	0.28316 (19)	0.55446 (18)	0.13993 (10)	0.0186 (2)
C6	0.5309 (2)	0.7013 (2)	0.16805 (13)	0.0273 (3)
C7	0.3162 (2)	0.18507 (18)	0.15790 (11)	0.0199 (3)
C8	-0.01503 (19)	0.25829 (19)	0.19763 (11)	0.0203 (3)
C9	0.8811 (2)	0.17879 (18)	0.54390 (11)	0.0191 (2)
O1	0.81796 (14)	0.31572 (13)	0.62220 (7)	0.0203 (2)
H2	0.574 (3)	0.240 (2)	0.3790 (14)	0.025 (4)*
H5A	0.215 (3)	0.492 (2)	0.0462 (15)	0.023 (4)*
H5B	0.196 (2)	0.646 (2)	0.1572 (14)	0.021 (3)*
H6A	0.599 (3)	0.782 (3)	0.2639 (17)	0.036 (4)*
H6B	0.635 (3)	0.615 (3)	0.1542 (17)	0.037 (4)*
H6C	0.544 (3)	0.819 (3)	0.1069 (16)	0.034 (4)*
H7A	0.226 (3)	0.138 (2)	0.0662 (15)	0.025 (4)*
H7B	0.479 (3)	0.238 (3)	0.1626 (16)	0.030 (4)*
H7C	0.286 (2)	0.050 (2)	0.2014 (14)	0.023 (3)*
H8A	-0.073 (3)	0.360 (2)	0.2338 (14)	0.021 (3)*
H8B	-0.046 (3)	0.123 (3)	0.2390 (15)	0.028 (4)*
H8C	-0.097 (3)	0.216 (2)	0.1031 (15)	0.024 (4)*
H9A	0.744 (3)	0.043 (2)	0.5000 (14)	0.024 (4)*
H9B	1.000 (3)	0.144 (2)	0.6036 (15)	0.027 (4)*
H9C	0.940 (2)	0.256 (2)	0.4774 (14)	0.021 (3)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0147 (5)	0.0146 (5)	0.0137 (5)	0.0060 (4)	0.0038 (4)	0.0032 (4)
C2	0.0169 (5)	0.0136 (5)	0.0134 (5)	0.0052 (4)	0.0052 (4)	0.0005 (4)
C3	0.0140 (5)	0.0128 (5)	0.0117 (5)	0.0025 (4)	0.0039 (4)	0.0012 (3)
C4	0.0161 (5)	0.0155 (5)	0.0113 (5)	0.0049 (4)	0.0027 (4)	0.0001 (4)
C5	0.0227 (6)	0.0191 (5)	0.0127 (5)	0.0061 (4)	0.0046 (4)	0.0025 (4)
C6	0.0246 (6)	0.0294 (6)	0.0248 (6)	0.0034 (5)	0.0089 (5)	0.0072 (5)
C7	0.0241 (6)	0.0199 (5)	0.0138 (5)	0.0087 (4)	0.0024 (4)	-0.0029 (4)
C8	0.0168 (5)	0.0222 (6)	0.0166 (5)	0.0028 (4)	0.0020 (4)	0.0005 (4)
C9	0.0215 (5)	0.0214 (5)	0.0179 (5)	0.0129 (4)	0.0056 (4)	0.0021 (4)
O1	0.0255 (4)	0.0258 (4)	0.0137 (4)	0.0173 (3)	0.0028 (3)	0.0007 (3)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C1—O1	1.3812 (14)	C6—H6A	1.043 (17)
C1—C2	1.3941 (15)	C6—H6B	1.037 (17)
C1—C3 <sup>i</sup>	1.4051 (15)	C6—H6C	1.038 (18)
C2—C3	1.3988 (16)	C7—H7A	0.976 (15)
C2—H2	0.963 (15)	C7—H7B	0.984 (17)
C3—C1 <sup>i</sup>	1.4051 (15)	C7—H7C	1.012 (16)
C3—C4	1.5371 (15)	C8—H8A	0.973 (15)
C4—C7	1.5376 (16)	C8—H8B	1.003 (17)
C4—C8	1.5428 (19)	C8—H8C	0.990 (15)
C4—C5	1.5494 (17)	C9—O1	1.4229 (14)
C5—C6	1.5172 (19)	C9—H9A	1.014 (16)
C5—H5A	0.991 (15)	C9—H9B	0.955 (16)
C5—H5B	0.985 (15)	C9—H9C	0.992 (15)
O1—C1—C2	121.95 (10)	H6A—C6—H6B	107.7 (13)
O1—C1—C3 <sup>i</sup>	117.05 (9)	C5—C6—H6C	110.9 (9)
C2—C1—C3 <sup>i</sup>	121.00 (10)	H6A—C6—H6C	107.8 (13)
C1—C2—C3	122.99 (10)	H6B—C6—H6C	107.0 (13)
C1—C2—H2	117.7 (9)	C4—C7—H7A	109.6 (9)
C3—C2—H2	119.3 (9)	C4—C7—H7B	112.6 (9)
C2—C3—C1 <sup>i</sup>	116.02 (10)	H7A—C7—H7B	107.7 (13)
C2—C3—C4	121.35 (9)	C4—C7—H7C	112.1 (8)
C1 <sup>i</sup> —C3—C4	122.63 (9)	H7A—C7—H7C	106.2 (12)
C3—C4—C7	111.63 (9)	H7B—C7—H7C	108.4 (12)
C3—C4—C8	109.87 (9)	C4—C8—H8A	111.4 (9)
C7—C4—C8	106.83 (9)	C4—C8—H8B	109.7 (9)
C3—C4—C5	111.30 (9)	H8A—C8—H8B	110.2 (13)
C7—C4—C5	108.72 (10)	C4—C8—H8C	108.6 (9)
C8—C4—C5	108.33 (9)	H8A—C8—H8C	108.9 (12)
C6—C5—C4	115.63 (10)	H8B—C8—H8C	107.9 (13)
C6—C5—H5A	110.5 (9)	O1—C9—H9A	109.7 (9)
C4—C5—H5A	106.6 (9)	O1—C9—H9B	104.8 (9)

---

C6—C5—H5B	107.9 (9)	H9A—C9—H9B	111.0 (12)
C4—C5—H5B	109.7 (9)	O1—C9—H9C	111.1 (8)
H5A—C5—H5B	106.2 (12)	H9A—C9—H9C	110.1 (12)
C5—C6—H6A	111.1 (10)	H9B—C9—H9C	110.1 (13)
C5—C6—H6B	112.1 (10)	C1—O1—C9	118.00 (9)

---

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .