

Methyl (*E*)-3-(2-bromo-4,5-dimethoxybenzylidene)dithiocarbazate

Zheng Fan,^a Yan-Lan Huang,^b Fang Tong,^b Yi-Wei Huang^b
and Shang Shan^{b*}

^aCollege of Biological and Environmental Engineering, Zhejiang University of Technology, People's Republic of China, and ^bCollege of Chemical Engineering and Materials Science, Zhejiang University of Technology, People's Republic of China
Correspondence e-mail: shanshang@mail.hz.zj.cn


Received 23 September 2011; accepted 25 September 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$;
 R factor = 0.035; wR factor = 0.079; data-to-parameter ratio = 15.4.

The title compound, $\text{C}_{11}\text{H}_{13}\text{BrN}_2\text{O}_2\text{S}_2$, was obtained from the condensation reaction of methyl dithiocarbazate and 2-bromo-4,5-dimethoxybenzaldehyde. In the molecule, the benzene ring and dithiocarbazate fragment are located on opposite sides of the $\text{C}=\text{N}$ bond, showing an *E* conformation. The dithiocarbazate fragment is approximately planar (r.m.s. deviation = 0.0281 \AA) and the mean plane is oriented at a dihedral angle of $11.38(15)^\circ$ with respect to the benzene ring. In the crystal, pairs of $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds link the molecules into centrosymmetric dimers.

Related literature

For applications of hydrazone and its derivatives in the biological field, see: Okabe *et al.* (1993); Hu *et al.* (2001). For related structures, see: Shan *et al.* (2008a,b,c). For the synthesis, see: Hu *et al.* (2001).



Experimental

Crystal data



$M_r = 349.26$

Triclinic, $P\bar{1}$	$V = 710.4(4)\text{ \AA}^3$
$a = 5.2460(12)\text{ \AA}$	$Z = 2$
$b = 11.781(5)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 12.400(5)\text{ \AA}$	$\mu = 3.18\text{ mm}^{-1}$
$\alpha = 102.347(3)^\circ$	$T = 293\text{ K}$
$\beta = 100.930(4)^\circ$	$0.42 \times 0.28 \times 0.25\text{ mm}$
$\gamma = 101.874(4)^\circ$	

Data collection

Rigaku R-AXIS RAPID IP diffractometer	5185 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	2553 independent reflections
$R_{\text{int}} = 0.026$	2051 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.73$, $T_{\max} = 0.82$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	166 parameters
$wR(F^2) = 0.079$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\max} = 0.32\text{ e \AA}^{-3}$
2553 reflections	$\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\text{N}\cdots\text{S}1^i$	0.86	2.62	3.456 (4)	166

Symmetry code: (i) $-x + 2, -y + 1, -z$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2002); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The work was supported by the Natural Science Foundation of Zhejiang Province, China (No. M203027).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5332).

References

- Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Hu, W., Sun, N. & Yang, Z. (2001). *Chem. J. Chin. Univ.* **22**, 2014–2017.
- Okabe, N., Nakamura, T. & Fukuda, H. (1993). *Acta Cryst.* **C49**, 1678–1680.
- Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
- Shan, S., Tian, Y.-L., Wang, S.-H., Wang, W.-L. & Xu, Y.-L. (2008a). *Acta Cryst.* **E64**, o1014.
- Shan, S., Tian, Y.-L., Wang, S.-H., Wang, W.-L. & Xu, Y.-L. (2008b). *Acta Cryst.* **E64**, o1024.
- Shan, S., Wang, S.-H., Tian, Y.-L., Wang, W.-L. & Xu, Y.-L. (2008c). *Acta Cryst.* **E64**, o1015.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, o2919 [doi:10.1107/S1600536811039304]

Methyl (*E*)-3-(2-bromo-4,5-dimethoxybenzylidene)dithiocarbazate

Zheng Fan, Yan-Lan Huang, Fang Tong, Yi-Wei Huang and Shang Shan

S1. Comment

Hydrazone and its derivatives have shown the potential application in the biological field (Okabe *et al.*, 1993; Hu *et al.*, 2001). As part of the ongoing investigation on anti-cancer compounds, the title compound has recently been prepared in our laboratory and its crystal structure is presented here.


In the molecules, the benzene ring and dithiocarbazate fragment are located on the opposite sides of the C=N bond, showing the *E*-configuration. This agrees with those found in the structures reported previously (Shan *et al.*, 2008*a,b*). The dithiocarbazate fragment is approximately planar, the r.m.s deviation being 0.0281 Å; its mean plane is oriented with respect to the benzene ring at 11.38 (15)°, similar to that found in a related structure (Shan *et al.* 2008*c*). In the crystal structure, intermolecular N—H···S hydrogen bonding links molecules to form the centro-symmetric dimers (Table 1).

S2. Experimental

Methyl dithiocarbazate was synthesized as described previously by Hu *et al.* (2001). Methyl dithiocarbazate (0.24 g, 2 mmol) and 2-bromo-4,5-dimethoxybenzaldehyde (0.49 g, 2 mmol) were dissolved in ethanol (20 ml), then acetic acid (0.2 ml) was added to the ethanol solution with stirring. The mixture solution was refluxed for 6 h. After cooling to room temperature, microcrystals appeared. The microcrystals were separated from the solution and washed with cold water three times. Recrystallization was performed twice with absolute methanol to obtain colourless single crystals of the title compound.

S3. Refinement

H atoms were placed in calculated positions with C—H = 0.93–0.96 Å and N—H = 0.86 Å, and refined in riding mode with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $1.2U_{\text{eq}}(\text{C},\text{N})$ for the others.

**Figure 1**

The molecular structure of the title compound with 40% probability displacement (arbitrary spheres for H atoms).

Methyl (*E*)-3-(2-bromo-4,5-dimethoxybenzylidene)dithiocarbazate

Crystal data



$M_r = 349.26$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.2460 (12)$ Å

$b = 11.781 (5)$ Å

$c = 12.400 (5)$ Å

$\alpha = 102.347 (3)^\circ$

$\beta = 100.930 (4)^\circ$

$\gamma = 101.874 (4)^\circ$

$V = 710.4 (4)$ Å³

$Z = 2$

$F(000) = 352$

$D_x = 1.633 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2553 reflections

$\theta = 2.8\text{--}25.2^\circ$

$\mu = 3.18 \text{ mm}^{-1}$

$T = 293$ K

Prism, colorless

$0.42 \times 0.28 \times 0.25$ mm

Data collection

Rigaku R-AXIS RAPID IP

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 10.0 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.73$, $T_{\max} = 0.82$

5185 measured reflections

2553 independent reflections

2051 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 2.8^\circ$

$h = -5 \rightarrow 6$

$k = -14 \rightarrow 11$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

$wR(F^2) = 0.079$

$S = 1.02$

2553 reflections

166 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.037P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.32 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.35 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.29930 (7)	0.52287 (3)	0.29304 (3)	0.06212 (15)
S1	1.14891 (18)	0.35920 (7)	-0.09979 (7)	0.0586 (2)
S2	0.95417 (16)	0.13601 (7)	-0.02534 (7)	0.0499 (2)
N1	0.7198 (5)	0.2821 (2)	0.11142 (18)	0.0411 (5)
N2	0.8444 (5)	0.3398 (2)	0.04224 (18)	0.0438 (6)
H2N	0.8360	0.4121	0.0428	0.053*
O1	0.3795 (4)	0.0498 (2)	0.38554 (17)	0.0620 (6)
O2	0.1403 (4)	0.1803 (2)	0.49916 (17)	0.0676 (6)
C1	0.3386 (5)	0.3734 (3)	0.3172 (2)	0.0438 (7)
C2	0.2229 (5)	0.3338 (3)	0.3994 (2)	0.0487 (7)
H2	0.1332	0.3810	0.4402	0.058*
C3	0.2405 (6)	0.2269 (3)	0.4200 (2)	0.0480 (7)
C4	0.3727 (5)	0.1538 (3)	0.3573 (2)	0.0459 (7)
C5	0.4848 (5)	0.1938 (3)	0.2762 (2)	0.0424 (7)
H5	0.5721	0.1457	0.2348	0.051*
C6	0.4722 (5)	0.3035 (2)	0.2539 (2)	0.0399 (6)
C7	0.6023 (5)	0.3466 (3)	0.1714 (2)	0.0418 (6)
H7	0.6001	0.4227	0.1621	0.050*
C8	0.9769 (5)	0.2856 (2)	-0.0251 (2)	0.0392 (6)
C9	1.1501 (7)	0.0933 (3)	-0.1230 (3)	0.0603 (9)
H9A	1.0899	0.1146	-0.1923	0.090*
H9B	1.1296	0.0081	-0.1393	0.090*
H9C	1.3358	0.1345	-0.0899	0.090*
C10	0.5363 (7)	-0.0214 (3)	0.3356 (3)	0.0635 (9)
H10A	0.4618	-0.0497	0.2548	0.095*
H10B	0.5356	-0.0888	0.3675	0.095*
H10C	0.7174	0.0263	0.3506	0.095*
C11	0.0060 (7)	0.2495 (4)	0.5675 (3)	0.0780 (12)
H11A	0.1296	0.3245	0.6114	0.117*
H11B	-0.0597	0.2057	0.6179	0.117*

H11C	-0.1419	0.2648	0.5192	0.117*
------	---------	--------	--------	--------

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.0681 (3)	0.0539 (2)	0.0700 (2)	0.02466 (17)	0.02423 (18)	0.01301 (17)
S1	0.0871 (6)	0.0518 (5)	0.0646 (5)	0.0304 (4)	0.0508 (5)	0.0325 (4)
S2	0.0635 (5)	0.0409 (4)	0.0588 (5)	0.0197 (4)	0.0304 (4)	0.0217 (4)
N1	0.0497 (14)	0.0445 (14)	0.0364 (12)	0.0116 (11)	0.0202 (11)	0.0177 (11)
N2	0.0615 (16)	0.0401 (13)	0.0443 (13)	0.0190 (12)	0.0286 (12)	0.0215 (11)
O1	0.0794 (16)	0.0694 (15)	0.0599 (13)	0.0274 (12)	0.0413 (12)	0.0352 (12)
O2	0.0721 (15)	0.0933 (17)	0.0550 (13)	0.0250 (13)	0.0420 (12)	0.0298 (12)
C1	0.0391 (16)	0.0493 (17)	0.0406 (15)	0.0114 (13)	0.0083 (13)	0.0083 (13)
C2	0.0431 (17)	0.070 (2)	0.0359 (15)	0.0200 (15)	0.0177 (13)	0.0067 (15)
C3	0.0424 (17)	0.070 (2)	0.0361 (15)	0.0117 (15)	0.0180 (13)	0.0177 (15)
C4	0.0435 (17)	0.0566 (19)	0.0369 (15)	0.0064 (14)	0.0137 (13)	0.0139 (14)
C5	0.0459 (17)	0.0515 (18)	0.0346 (14)	0.0148 (14)	0.0190 (13)	0.0112 (13)
C6	0.0382 (15)	0.0470 (17)	0.0354 (14)	0.0094 (13)	0.0133 (12)	0.0106 (13)
C7	0.0481 (17)	0.0421 (16)	0.0391 (15)	0.0117 (13)	0.0159 (13)	0.0142 (13)
C8	0.0474 (16)	0.0425 (16)	0.0346 (14)	0.0158 (13)	0.0144 (13)	0.0166 (12)
C9	0.073 (2)	0.0517 (19)	0.073 (2)	0.0287 (17)	0.0390 (18)	0.0213 (17)
C10	0.083 (2)	0.062 (2)	0.064 (2)	0.0289 (19)	0.0369 (19)	0.0264 (17)
C11	0.067 (2)	0.124 (3)	0.0492 (19)	0.024 (2)	0.0377 (18)	0.016 (2)

Geometric parameters (\AA , $^\circ$)

Br—C1	1.893 (3)	C3—C4	1.411 (4)
S1—C8	1.662 (3)	C4—C5	1.376 (4)
S2—C8	1.741 (3)	C5—C6	1.390 (4)
S2—C9	1.789 (3)	C5—H5	0.9300
N1—C7	1.280 (3)	C6—C7	1.454 (4)
N1—N2	1.381 (3)	C7—H7	0.9300
N2—C8	1.328 (3)	C9—H9A	0.9600
N2—H2N	0.8600	C9—H9B	0.9600
O1—C4	1.349 (3)	C9—H9C	0.9600
O1—C10	1.421 (4)	C10—H10A	0.9600
O2—C3	1.360 (3)	C10—H10B	0.9600
O2—C11	1.431 (4)	C10—H10C	0.9600
C1—C2	1.395 (4)	C11—H11A	0.9600
C1—C6	1.398 (4)	C11—H11B	0.9600
C2—C3	1.355 (4)	C11—H11C	0.9600
C2—H2	0.9300		
C8—S2—C9	101.93 (13)	N1—C7—C6	121.3 (3)
C7—N1—N2	113.4 (2)	N1—C7—H7	119.3
C8—N2—N1	121.0 (2)	C6—C7—H7	119.3
C8—N2—H2N	119.5	N2—C8—S1	120.9 (2)
N1—N2—H2N	119.5	N2—C8—S2	114.23 (19)

C4—O1—C10	118.1 (2)	S1—C8—S2	124.83 (16)
C3—O2—C11	117.8 (3)	S2—C9—H9A	109.5
C2—C1—C6	120.9 (3)	S2—C9—H9B	109.5
C2—C1—Br	117.3 (2)	H9A—C9—H9B	109.5
C6—C1—Br	121.8 (2)	S2—C9—H9C	109.5
C3—C2—C1	120.3 (3)	H9A—C9—H9C	109.5
C3—C2—H2	119.8	H9B—C9—H9C	109.5
C1—C2—H2	119.8	O1—C10—H10A	109.5
C2—C3—O2	125.5 (3)	O1—C10—H10B	109.5
C2—C3—C4	120.3 (2)	H10A—C10—H10B	109.5
O2—C3—C4	114.2 (3)	O1—C10—H10C	109.5
O1—C4—C5	126.1 (3)	H10A—C10—H10C	109.5
O1—C4—C3	115.1 (2)	H10B—C10—H10C	109.5
C5—C4—C3	118.7 (3)	O2—C11—H11A	109.5
C4—C5—C6	122.3 (3)	O2—C11—H11B	109.5
C4—C5—H5	118.9	H11A—C11—H11B	109.5
C6—C5—H5	118.9	O2—C11—H11C	109.5
C5—C6—C1	117.5 (2)	H11A—C11—H11C	109.5
C5—C6—C7	121.5 (2)	H11B—C11—H11C	109.5
C1—C6—C7	121.0 (3)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2N···S1 ⁱ	0.86	2.62	3.456 (4)	166

Symmetry code: (i) $-x+2, -y+1, -z$.