

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[(1,2,5,6- η)-Cycloocta-1,5-diene]bis(1-isopropyl-3-methylimidazolin-2-ylidene)-rhodium(I) tetrafluoroborate

Gary S. Nichol,^{a*} Jonathan Rajaseelan,^b David P. Walton^c and Edward Rajaseelan^c

^aDepartment of Chemistry and Biochemistry, The University of Arizona, Tucson, AZ 85716, USA, ^bLancaster Country Day School, Lancaster, PA 17603, USA, and ^cDepartment of Chemistry, Millersville University, Millersville, PA 17551, USA
Correspondence e-mail: gsnichol@email.arizona.edu


Received 10 November 2011; accepted 21 November 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.020; wR factor = 0.059; data-to-parameter ratio = 64.3.

In the title compound, $[\text{Rh}(\text{C}_8\text{H}_{12})(\text{C}_7\text{H}_{12}\text{N}_2)_2]\text{BF}_4$, the square-planar Rh complex cation and the BF_4^- anion are both bisected by a crystallographic twofold rotation axis. The Rh and B atoms lie on this axis and all others are in general positions. In the crystal, two unique C—H...F hydrogen-bonding interactions are present, which involve both imidazolin-2-ylidene H atoms. They form two separate $C(5)$ motifs, the combination of which is a rippled hydrogen-bonded sheet structure in the ab plane.

Related literature

For the structure and dynamics of related N -heterocyclic carbene rhodium and iridium complexes, see: Chianese *et al.* (2003); Köcher & Herrmann (1997); Leung *et al.* (2006); Nichol *et al.* (2009, 2010); Herrmann *et al.* (2006). For the catalytic properties of these complexes, see: Albrecht *et al.* (2002); Frey *et al.* (2006); Gnanamgari *et al.* (2007); Voutchkova *et al.* (2008).



Experimental

Crystal data

$[\text{Rh}(\text{C}_8\text{H}_{12})(\text{C}_7\text{H}_{12}\text{N}_2)_2]\text{BF}_4$
 $M_r = 546.27$
 Orthorhombic, $Pccn$
 $a = 11.7508$ (6) Å
 $b = 11.9283$ (6) Å
 $c = 17.3129$ (9) Å
 $V = 2426.7$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.75$ mm⁻¹
 $T = 100$ K
 $0.38 \times 0.37 \times 0.37$ mm

Data collection

Bruker Kappa APEXII DUO CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.763$, $T_{\max} = 0.771$
 234794 measured reflections
 14018 independent reflections
 10241 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.020$
 $wR(F^2) = 0.059$
 $S = 1.13$
 14018 reflections
 218 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 1.55$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.92$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C2}-\text{H2}\cdots\text{F1}^i$	0.909 (11)	2.496 (11)	3.3975 (8)	171.4 (10)
$\text{C3}-\text{H3}\cdots\text{F2}^{ii}$	0.877 (12)	2.478 (12)	3.2415 (8)	145.9 (11)

Symmetry codes: (i) $x - \frac{1}{2}, -y + 1, -z + \frac{1}{2}$; (ii) $x, y - 1, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXTL and publCIF (Westrip, 2010).

JR and DPW thank the Department of Chemistry, Millersville University, for project funding. The diffractometer was purchased with funding from NSF grant CHE-0741837.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2478).

References

- Albrecht, M., Miecznikowski, J. R., Samuel, A., Faller, J. W. & Crabtree, R. H. (2002). *Organometallics*, **21**, 3596–3604.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chianese, A. R., Li, X., Jansen, M. C., Faller, J. W. & Crabtree, R. H. (2003). *Organometallics*, **22**, 1663–1667.
- Frey, G. D., Rentzsch, C. F., von Preysing, D., Scherg, T., Mühlhofer, M., Herdtweck, E. & Herrmann, W. A. (2006). *J. Organomet. Chem.* **691**, 5725–5738.
- Gnanamgari, D., Moores, A., Rajaseelan, E. & Crabtree, R. H. (2007). *Organometallics*, **26**, 1226–1230.
- Herrmann, W. A., Schütz, J., Frey, G. D. & Herdtweck, E. (2006). Köcher, C. & Herrmann, W. A. (1997). *J. Organomet. Chem.* **532**, 261–265.
- Leung, C. H., Incarvito, C. D. & Crabtree, R. H. (2006). *Organometallics*, **25**, 6099–6107.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nichol, G. S., Rajaseelan, J., Anna, L. J. & Rajaseelan, E. (2009). *Eur. J. Inorg. Chem.* pp. 4320–4328.
- Nichol, G. S., Stasiw, D., Anna, L. J. & Rajaseelan, E. (2010). *Acta Cryst.* **E66**, m1114.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Voutchkova, A. M., Gnanamgari, D., Jakobsche, C. E., Butler, C., Miller, S. J., Parr, J. & Crabtree, R. H. (2008). *J. Organomet. Chem.* **693**, 1815–1821.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, m1860–m1861 [https://doi.org/10.1107/S1600536811049890]

[(1,2,5,6- η)-Cycloocta-1,5-diene]bis(1-isopropyl-3-methylimidazolin-2-ylidene)rhodium(I) tetrafluoridoborate

Gary S. Nichol, Jonathan Rajaseelan, David P. Walton and Edward Rajaseelan

S1. Comment

We are interested in rhodium and iridium complexes with N-heterocyclic carbene ligands, in particular ligands derived from 1,2,4-triazole-derived compounds (Nichol *et al.*, 2009, 2010). The title compound, (I), was prepared as part of this study (Figure 1). The Rh center has an expected square planar geometry and bond distances are unexceptional. Both the Rh and B atoms lie on a crystallographic twofold rotation axis, which bisects the complex and BF_4^- counterion. C–H \cdots F hydrogen bonding interactions, which involve both imidazolin-2-ylidene H atoms and all four F atoms, form a thick two-dimensional sheet structure in the *ab* plane (Figure 2).

S2. Experimental

The title compound was synthesized by transmetalation. 1-Isopropyl-3-methylimidazolium bromide (268 mg, 1.31 mmol) was mixed with Ag_2O (152 mg, 0.654 mmol), and was stirred under dark at room temperature for 90 minutes in 10 ml of CH_2Cl_2 . The resulting mixture was filtered through Celite into a new flask containing the neutral compound [(cod)Rh(NHC)Cl] (585 mg, 1.31 mmol), and AgBF_4 (254 mg, 1.31 mmol) and stirred for an additional 90 minutes under dark. The mixture was filtered once more through Celite to remove silver bromide and silver chloride, and the solvent was removed under pressure to give a yellow solid (93%). Crystals of the resulting solid of the title compound, (I), were obtained by slow diffusion of pentane into dichloromethane solution of the compound. ^1H NMR (400 MHz, CDCl_3): δ (p.p.m.) = 7.15 (s, 2 H, NCH), 6.93 (s, 2 H, NCH), 5.03 (m, $^3J_{\text{H-H}} = 6.8$ Hz, 2 H, CH of ^iPr), 4.63 (br, 2 H, CH of COD), 4.21 (s, 6 H, N— CH_3), 3.92 (m, 2 H, CH of COD), 2.63 (m, 2 H, CH_2 of COD), 2.42 – 1.92 (m, 6 H, CH_2 of COD), 1.46 (d, $^3J_{\text{H-H}} = 6.8$ Hz, 6 H, CH_3 of ^iPr), 1.00 (d, $^3J_{\text{H-H}} = 6.8$ Hz, 6 H, CH_3 of ^iPr). ^{13}C NMR: $\delta = 178.76, 178.22$ (Ir—C), 124, 117 (N—CH—N), 91.34, 91.25 (N— CHMe_3), 86.36, 86.28 (N— CH_3), 52.60 (CH of COD), 38.10, 33.75, 27.99, (CH $_2$ of COD), 23.5, 22.90 (CH $_3$ of ^iPr).

S3. Refinement

H atoms were located from a difference Fourier map and are freely refined.




Figure 1

Twice the asymmetric unit of (I), with H atoms omitted. Displacement ellipsoids are at the 50% probability level. Unlabeled atoms are related to labeled atoms by twofold rotation symmetry.




Figure 2

A *c*-axis projection showing C–H···F interactions (blue dotted lines) in (I). Red dotted lines indicate H bond continuation.

[(1,2,5,6- η)-Cycloocta-1,5-diene]bis(1-isopropyl-3-methylimidazolin-2-ylidene)rhodium(I) tetrafluoroborate

Crystal data

[Rh(C₈H₁₂)(C₇H₁₂N₂)₂]BF₄

M_r = 546.27

Orthorhombic, *Pccn*

Hall symbol: -P 2ab 2ac

a = 11.7508 (6) Å

b = 11.9283 (6) Å

c = 17.3129 (9) Å

V = 2426.7 (2) Å³

Z = 4

F(000) = 1128

D_x = 1.495 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 9624 reflections

θ = 4.2–51.7°

μ = 0.75 mm⁻¹

T = 100 K

Block, yellow

0.38 × 0.37 × 0.37 mm

Data collection

Bruker Kappa APEXII DUO CCD
diffractometer

Radiation source: fine-focus sealed tube with
Miracol optics

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

T_{min} = 0.763, *T_{max}* = 0.771

234794 measured reflections

14018 independent reflections

10241 reflections with *I* > 2σ(*I*)

R_{int} = 0.033

θ_{max} = 52.3°, θ_{min} = 2.9°

h = -26→25

k = -26→26

l = -37→38

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.020$ $wR(F^2) = 0.059$ $S = 1.13$

14018 reflections

218 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
map

Hydrogen site location: difference Fourier map

All H-atom parameters refined

 $w = 1/[\sigma^2(F_o^2) + (0.0182P)^2 + 0.6772P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.002$ $\Delta\rho_{\max} = 1.55 \text{ e } \text{Å}^{-3}$ $\Delta\rho_{\min} = -0.92 \text{ e } \text{Å}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Rh1	0.7500	0.2500	0.507678 (3)	0.01052 (1)
N1	0.56243 (4)	0.20105 (4)	0.38873 (3)	0.01399 (6)
N2	0.66738 (4)	0.05648 (4)	0.40496 (3)	0.01512 (6)
C1	0.65651 (4)	0.16534 (4)	0.42664 (3)	0.01295 (6)
C2	0.51512 (5)	0.11598 (5)	0.34425 (4)	0.01757 (8)
H2	0.4526 (10)	0.1269 (10)	0.3143 (7)	0.022 (3)*
C3	0.58158 (5)	0.02502 (5)	0.35458 (4)	0.01811 (8)
H3	0.5769 (10)	-0.0434 (10)	0.3364 (7)	0.028 (3)*
C4	0.51719 (5)	0.31568 (5)	0.39338 (3)	0.01602 (7)
H4	0.5633 (9)	0.3517 (9)	0.4313 (6)	0.019 (3)*
C5	0.53212 (7)	0.37474 (6)	0.31623 (4)	0.02376 (11)
H5A	0.4910 (11)	0.3362 (11)	0.2772 (7)	0.030 (3)*
H5B	0.6089 (11)	0.3778 (11)	0.3028 (8)	0.031 (3)*
H5C	0.5034 (11)	0.4517 (11)	0.3183 (8)	0.035 (3)*
C6	0.39373 (6)	0.31435 (7)	0.41936 (5)	0.02579 (12)
H6A	0.3478 (11)	0.2754 (11)	0.3834 (8)	0.028 (3)*
H6B	0.3635 (11)	0.3876 (11)	0.4230 (8)	0.035 (3)*
H6C	0.3844 (12)	0.2775 (12)	0.4678 (9)	0.033 (3)*
C7	0.75478 (6)	-0.02030 (5)	0.43195 (4)	0.02033 (9)
H7A	0.8005 (10)	0.0166 (10)	0.4689 (7)	0.024 (3)*
H7B	0.7196 (11)	-0.0845 (12)	0.4540 (8)	0.031 (3)*
H7C	0.8014 (11)	-0.0430 (11)	0.3906 (8)	0.032 (3)*
C8	0.61291 (5)	0.22486 (5)	0.59096 (3)	0.01603 (7)
H8	0.5433 (10)	0.2098 (10)	0.5610 (7)	0.021 (3)*
C9	0.68830 (5)	0.13645 (5)	0.60051 (3)	0.01646 (8)

H9	0.6691 (10)	0.0669 (10)	0.5751 (7)	0.025 (3)*
C10	0.77717 (6)	0.12488 (6)	0.66365 (4)	0.01943 (9)
H10A	0.7517 (9)	0.1609 (11)	0.7107 (8)	0.023 (3)*
H10B	0.7852 (10)	0.0442 (11)	0.6775 (7)	0.026 (3)*
C11	0.89369 (5)	0.16926 (6)	0.63829 (4)	0.01888 (9)
H11A	0.9440 (9)	0.1813 (9)	0.6833 (7)	0.020 (2)*
H11B	0.9335 (10)	0.1111 (10)	0.6049 (7)	0.023 (3)*
B1	0.7500	0.7500	0.29593 (6)	0.01705 (12)
F1	0.76439 (5)	0.84479 (5)	0.24966 (3)	0.03003 (10)
F2	0.65419 (4)	0.76347 (4)	0.34218 (3)	0.02659 (9)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Rh1	0.01063 (2)	0.00980 (2)	0.01114 (2)	−0.00124 (1)	0.000	0.000
N1	0.01412 (14)	0.01361 (14)	0.01424 (15)	0.00070 (11)	−0.00193 (11)	−0.00109 (12)
N2	0.01684 (16)	0.01164 (14)	0.01689 (16)	0.00010 (12)	−0.00365 (13)	−0.00144 (12)
C1	0.01345 (15)	0.01194 (15)	0.01347 (16)	−0.00035 (12)	−0.00093 (12)	−0.00044 (12)
C2	0.01794 (19)	0.01713 (19)	0.0176 (2)	−0.00031 (15)	−0.00510 (16)	−0.00262 (15)
C3	0.0211 (2)	0.01475 (18)	0.0185 (2)	−0.00131 (16)	−0.00536 (17)	−0.00293 (15)
C4	0.01615 (18)	0.01608 (18)	0.01583 (18)	0.00398 (14)	−0.00011 (14)	−0.00123 (14)
C5	0.0321 (3)	0.0195 (2)	0.0197 (2)	0.0075 (2)	0.0032 (2)	0.00306 (19)
C6	0.0185 (2)	0.0318 (3)	0.0270 (3)	0.0065 (2)	0.0043 (2)	−0.0023 (2)
C7	0.0218 (2)	0.01268 (16)	0.0265 (3)	0.00234 (17)	−0.0073 (2)	−0.00142 (16)
C8	0.01421 (17)	0.01788 (18)	0.01599 (18)	−0.00244 (14)	0.00177 (14)	−0.00026 (15)
C9	0.01790 (19)	0.01465 (17)	0.01685 (19)	−0.00334 (15)	0.00120 (15)	0.00176 (14)
C10	0.0215 (2)	0.0200 (2)	0.0168 (2)	−0.00095 (18)	−0.00002 (17)	0.00507 (17)
C11	0.0179 (2)	0.0210 (2)	0.0178 (2)	0.00064 (17)	−0.00287 (16)	0.00310 (17)
B1	0.0149 (3)	0.0145 (3)	0.0218 (3)	0.0006 (2)	0.000	0.000
F1	0.0346 (3)	0.0235 (2)	0.0320 (2)	0.00074 (18)	0.00329 (18)	0.01073 (17)
F2	0.01926 (17)	0.02301 (19)	0.0375 (3)	0.00023 (13)	0.00938 (16)	−0.00171 (16)

Geometric parameters (Å, °)

Rh1—C1	2.0482 (5)	C6—H6A	0.946 (13)
Rh1—C1 ⁱ	2.0482 (5)	C6—H6B	0.946 (14)
Rh1—C8	2.1826 (6)	C6—H6C	0.952 (15)
Rh1—C8 ⁱ	2.1826 (6)	C7—H7A	0.944 (12)
Rh1—C9	2.2233 (6)	C7—H7B	0.950 (14)
Rh1—C9 ⁱ	2.2233 (6)	C7—H7C	0.941 (13)
N1—C1	1.3544 (7)	C8—H8	0.985 (12)
N1—C2	1.3899 (7)	C8—C9	1.3872 (9)
N1—C4	1.4692 (7)	C8—C11 ⁱ	1.5074 (9)
N2—C1	1.3577 (7)	C9—H9	0.965 (12)
N2—C3	1.3850 (7)	C9—C10	1.5181 (9)
N2—C7	1.4532 (8)	C10—H10A	0.968 (13)
C2—H2	0.909 (11)	C10—H10B	0.997 (13)
C2—C3	1.3487 (9)	C10—C11	1.5322 (9)

C3—H3	0.877 (12)	C11—C8 ⁱ	1.5075 (9)
C4—H4	0.954 (11)	C11—H11A	0.989 (11)
C4—C5	1.5202 (9)	C11—H11B	1.017 (12)
C4—C6	1.5190 (9)	B1—F1	1.3960 (8)
C5—H5A	0.950 (13)	B1—F1 ⁱⁱ	1.3960 (8)
C5—H5B	0.933 (13)	B1—F2	1.3908 (8)
C5—H5C	0.979 (14)	B1—F2 ⁱⁱ	1.3909 (8)
C1—Rh1—C1 ⁱ	93.53 (3)	C4—C6—H6A	110.8 (8)
C1—Rh1—C8	89.36 (2)	C4—C6—H6B	111.7 (8)
C1 ⁱ —Rh1—C8 ⁱ	89.36 (2)	C4—C6—H6C	112.0 (8)
C1—Rh1—C8 ⁱ	156.36 (2)	H6A—C6—H6B	106.4 (11)
C1 ⁱ —Rh1—C8	156.36 (2)	H6A—C6—H6C	106.7 (11)
C1—Rh1—C9	91.14 (2)	H6B—C6—H6C	108.9 (11)
C1 ⁱ —Rh1—C9	166.03 (2)	N2—C7—H7A	109.0 (7)
C1 ⁱ —Rh1—C9 ⁱ	91.14 (2)	N2—C7—H7B	109.3 (8)
C1—Rh1—C9 ⁱ	166.03 (2)	N2—C7—H7C	110.4 (8)
C8—Rh1—C8 ⁱ	97.31 (3)	H7A—C7—H7B	110.6 (11)
C8—Rh1—C9	36.69 (2)	H7A—C7—H7C	108.6 (11)
C8 ⁱ —Rh1—C9	81.19 (2)	H7B—C7—H7C	109.0 (11)
C8 ⁱ —Rh1—C9 ⁱ	36.69 (2)	Rh1—C8—H8	106.8 (7)
C8—Rh1—C9 ⁱ	81.19 (2)	Rh1—C8—C9	73.25 (3)
C9—Rh1—C9 ⁱ	87.42 (3)	Rh1—C8—C11 ⁱ	106.38 (4)
C1—N1—C2	111.43 (5)	H8—C8—C9	117.0 (7)
C1—N1—C4	124.16 (5)	H8—C8—C11 ⁱ	113.3 (7)
C2—N1—C4	124.41 (5)	C9—C8—C11 ⁱ	127.23 (5)
C1—N2—C3	111.39 (5)	Rh1—C9—C8	70.06 (3)
C1—N2—C7	125.48 (5)	Rh1—C9—H9	105.7 (7)
C3—N2—C7	123.10 (5)	Rh1—C9—C10	110.59 (4)
Rh1—C1—N1	127.93 (4)	C8—C9—H9	116.7 (7)
Rh1—C1—N2	127.62 (4)	C8—C9—C10	126.46 (6)
N1—C1—N2	104.10 (4)	H9—C9—C10	114.2 (7)
N1—C2—H2	122.4 (7)	C9—C10—H10A	110.7 (7)
N1—C2—C3	106.40 (5)	C9—C10—H10B	109.0 (7)
H2—C2—C3	131.2 (7)	C9—C10—C11	112.14 (5)
N2—C3—C2	106.68 (5)	H10A—C10—H10B	104.8 (11)
N2—C3—H3	121.6 (8)	H10A—C10—C11	111.3 (7)
C2—C3—H3	131.7 (8)	H10B—C10—C11	108.6 (7)
N1—C4—H4	104.6 (7)	C8 ⁱ —C11—C10	113.53 (5)
N1—C4—C5	109.96 (5)	C8 ⁱ —C11—H11A	109.7 (7)
N1—C4—C6	110.61 (5)	C8 ⁱ —C11—H11B	106.6 (7)
H4—C4—C5	109.3 (7)	C10—C11—H11A	111.0 (7)
H4—C4—C6	110.0 (7)	C10—C11—H11B	109.8 (7)
C5—C4—C6	112.04 (6)	H11A—C11—H11B	105.8 (9)
C4—C5—H5A	110.0 (8)	F1—B1—F1 ⁱⁱ	109.97 (9)
C4—C5—H5B	110.4 (8)	F1—B1—F2	109.56 (3)
C4—C5—H5C	111.2 (8)	F1 ⁱⁱ —B1—F2	109.02 (3)
H5A—C5—H5B	109.5 (11)	F1 ⁱⁱ —B1—F2 ⁱⁱ	109.56 (3)

H5A—C5—H5C	107.7 (11)	F1—B1—F2 ⁱⁱ	109.02 (3)
H5B—C5—H5C	107.9 (11)	F2—B1—F2 ⁱⁱ	109.70 (9)
C2—N1—C1—Rh1	-173.45 (4)	C2—N1—C4—C6	56.65 (8)
C2—N1—C1—N2	0.13 (6)	C1—Rh1—C8—C9	92.76 (4)
C4—N1—C1—Rh1	7.28 (8)	C1 ⁱ —Rh1—C8—C9	-169.87 (5)
C4—N1—C1—N2	-179.14 (5)	C1 ⁱ —Rh1—C8—C11 ⁱ	-45.11 (7)
C3—N2—C1—Rh1	173.54 (4)	C1—Rh1—C8—C11 ⁱ	-142.49 (4)
C3—N2—C1—N1	-0.07 (6)	C8 ⁱ —Rh1—C8—C9	-64.50 (3)
C7—N2—C1—Rh1	-4.70 (8)	C8 ⁱ —Rh1—C8—C11 ⁱ	60.26 (4)
C7—N2—C1—N1	-178.31 (6)	C9 ⁱ —Rh1—C8—C9	-97.57 (4)
C1 ⁱ —Rh1—C1—N1	-83.65 (5)	C9—Rh1—C8—C11 ⁱ	124.75 (6)
C1 ⁱ —Rh1—C1—N2	104.22 (5)	C9 ⁱ —Rh1—C8—C11 ⁱ	27.19 (4)
C8—Rh1—C1—N1	72.88 (5)	Rh1—C8—C9—C10	101.36 (6)
C8 ⁱ —Rh1—C1—N1	179.85 (5)	C11 ⁱ —C8—C9—Rh1	-98.12 (6)
C8—Rh1—C1—N2	-99.26 (5)	C11 ⁱ —C8—C9—C10	3.25 (9)
C8 ⁱ —Rh1—C1—N2	7.72 (8)	C1—Rh1—C9—C8	-87.40 (4)
C9—Rh1—C1—N1	109.52 (5)	C1 ⁱ —Rh1—C9—C8	163.01 (8)
C9 ⁱ —Rh1—C1—N1	25.66 (11)	C1—Rh1—C9—C10	149.98 (4)
C9—Rh1—C1—N2	-62.61 (5)	C1 ⁱ —Rh1—C9—C10	40.40 (11)
C9 ⁱ —Rh1—C1—N2	-146.48 (8)	C8 ⁱ —Rh1—C9—C8	115.05 (4)
C1—N1—C2—C3	-0.15 (7)	C8—Rh1—C9—C10	-122.61 (6)
C4—N1—C2—C3	179.12 (5)	C8 ⁱ —Rh1—C9—C10	-7.56 (4)
N1—C2—C3—N2	0.09 (7)	C9 ⁱ —Rh1—C9—C8	78.69 (3)
C1—N2—C3—C2	-0.01 (7)	C9 ⁱ —Rh1—C9—C10	-43.92 (4)
C7—N2—C3—C2	178.27 (6)	Rh1—C9—C10—C11	-13.94 (7)
C1—N1—C4—C5	111.55 (6)	C8—C9—C10—C11	-93.84 (7)
C1—N1—C4—C6	-124.17 (6)	C9—C10—C11—C8 ⁱ	39.69 (8)
C2—N1—C4—C5	-67.63 (8)		

Symmetry codes: (i) $-x+3/2, -y+1/2, z$; (ii) $-x+3/2, -y+3/2, z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H2...F1 ⁱⁱⁱ	0.909 (11)	2.496 (11)	3.3975 (8)	171.4 (10)
C3—H3...F2 ^{iv}	0.877 (12)	2.478 (12)	3.2415 (8)	145.9 (11)

Symmetry codes: (iii) $x-1/2, -y+1, -z+1/2$; (iv) $x, y-1, z$.