

(2-Bromoacetyl)ferrocene

Xiang-Xiang Wu,^a Xin Zhu,^a Qiu-Juan Ma,^a Seik Weng Ng^{b,c} and Edward R. T. Tiekkink^{b*}

^aHenan University of Traditional Medicine, Zhengzhou 450008, People's Republic of China, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: Edward.Tiekkink@gmail.com


Received 23 November 2011; accepted 24 November 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004 \text{ \AA}$; R factor = 0.035; wR factor = 0.079; data-to-parameter ratio = 18.4.

In the title molecule, $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_7\text{H}_6\text{BrO})]$, the C atoms of the substituted ring have disparate Fe–C bond lengths compared with the unsubstituted ring. In the bromoacetyl residue, the Br and O atoms are co-planar [the $\text{O}-\text{C}-\text{C}-\text{Br}$ torsion angle is $5.7(4)^\circ$] and are *syn* to each other. Helical supramolecular chains along the *b* axis are formed in the crystal structure mediated by $\text{C}-\text{H}\cdots\text{O}$ contacts; the carbonyl-O atom is bifurcated. The chains are linked into layers by $\text{C}-\text{H}\cdots\pi$ (unsubstituted ring) interactions that stack along the *a*-axis direction.

Related literature

For background to the potential applications of ferrocenyl derivatives in medicine and as biosensors, see: Arezki *et al.* (2011); Huang *et al.* (2008); Yang *et al.* (2007).



Experimental

Crystal data

$[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_7\text{H}_6\text{BrO})]$
 $M_r = 306.97$
Monoclinic, $P2_1/c$
 $a = 7.7095(3) \text{ \AA}$
 $b = 9.6609(4) \text{ \AA}$
 $c = 14.7464(7) \text{ \AA}$
 $\beta = 98.061(4)^\circ$
 $V = 1087.47(8) \text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 5.03 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 $0.30 \times 0.10 \times 0.03 \text{ mm}$

Data collection

Agilent SuperNova Dual diffractometer with Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.314$, $T_{\max} = 0.864$
7692 measured reflections
2497 independent reflections
2004 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.079$
 $S = 1.02$
2497 reflections
136 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.73 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.58 \text{ e \AA}^{-3}$

Table 1
Selected bond lengths (Å).

Fe1–C1	2.053 (3)	Fe1–C6	2.037 (3)
Fe1–C2	2.051 (3)	Fe1–C7	2.060 (3)
Fe1–C3	2.042 (3)	Fe1–C8	2.059 (3)
Fe1–C4	2.040 (3)	Fe1–C9	2.043 (3)
Fe1–C5	2.040 (3)	Fe1–C10	2.027 (3)

Table 2

Hydrogen-bond geometry (Å, °).

$Cg1$ is the centroid of the C1–C5 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1–H1 \cdots O1 ⁱ	1.00	2.49	3.327 (4)	140
C12–H12a \cdots O1 ⁱ	0.99	2.35	3.291 (4)	158
C12–H12b \cdots Cg1 ⁱⁱ	0.99	2.64	3.445 (3)	139

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

We thank Henan University of Traditional Medicine and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5144).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Arezki, A., Chabot, G. G., Quentin, L., Scherman, D., Jaouena, G. & Brulé, E. (2011). *Med. Chem. Commun.*, **2**, 190–195.
- Barbour, L. J. (2001). *J. Supramol. Chem.*, **1**, 189–191.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Huang, K., Yang, H., Zhou, Z., Yu, M., Li, F., Gao, X., Yi, T. & Huang, C. (2008). *Org. Lett.*, **10**, 2557–2560.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yang, H., Zhou, Z., Huang, K., Yu, M., Li, F., Yi, T. & Huang, C. (2007). *Org. Lett.*, **9**, 4729–4732.

supporting information

Acta Cryst. (2011). E67, m1875 [https://doi.org/10.1107/S1600536811050434]

(2-Bromoacetyl)ferrocene

Xiang-Xiang Wu, Xin Zhu, Qiu-Juan Ma, Seik Weng Ng and Edward R. T. Tiekink

S1. Comment

Ferrocenyl derivatives continue to attract attention owing to their applications as biosensors and drugs (Arezki *et al.*, 2011; Huang *et al.*, 2008; Yang *et al.*, 2007). The title compound, (I), is a synthetic precursor for such molecules.

The molecular structure of (I), Fig. 1, features substituted and unsubstituted cyclopentadienyl rings which form a dihedral angle of 1.14 (18) $^{\circ}$ indicating an almost parallel relationship; the rings are almost eclipsed when viewed down the ring centroid–Fe–ring centroid axis. The Fe–C bond distances formed by the C atoms in the unsubstituted ring are equal within experimental error, *i.e.* 2.040 (3) to 2.053 (3) Å, but a disparity is evident in the Fe–C bond distances formed by the substituted ring, range of Fe–C is 2.027 (3) Å, for the C atom carrying the substituent, to 2.060 (3) Å; Table 1. The average Fe–C bond distances are experimentally equivalent as is reflected in the Fe–ring centroid distances of 1.6480 (14), for the C1-ring, compared to 1.6447 (14) Å for the C6-ring; the ring centroid–Fe–ring centroid angle is 179.68 (7) $^{\circ}$. In the bromoacetyl residue, the Br and O atoms are co-planar [the O1–C11–C12–Br1 torsion angle is 5.7 (4) $^{\circ}$] and are *syn* to each other.

Supramolecular helical chains along the *b* axis and mediated by C–H···O contacts involving the bifurcated carbonyl–O atom feature in the crystal packing, Fig. 2. These are connected into layers in the *bc* plane by C–H··· π contacts, Fig. 3. The layers are undulating and stack along the *a* axis, Fig. 4.

S2. Experimental

A flask was charged with ferrocene (281 mg, 1.51 mmol) and 2-bromoacetyl bromide (305 mg, 1.51 mmol) in dichloromethane (30 ml); the flask was chilled in an ice-bath. To the solution was added anhydrous aluminium chloride (221 mg, 1.66 mmol) under a nitrogen atmosphere. The reaction mixture, which turned deep-violet, was stirred for an hour. Water was added and the organic phase was extracted with dichloromethane (3×15 ml). The organic layer was dried over anhydrous sodium sulfate. The solvent was removed and the product purified by flash column chromatography (ethyl acetate/hexane, 1:9) to give the product as a brown solid; crystals were grown from its solution in dichloromethane.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [C–H 0.99 to 1.00 Å, $U_{\text{iso}}(\text{H}) 1.2U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation.




Figure 1

Molecular structure of (I) showing atom-labelling scheme and displacement ellipsoids at the 70% probability level.




Figure 2

Supramolecular helical chain along the *b* axis in (I). The C—H···O contacts are shown as orange dashed lines.





Figure 3

Supramolecular layer in the bc plane in (I). The C—H···O and C—H··· π contacts are shown as orange and purple dashed lines, respectively.

**Figure 4**

Stacking of layers along the *a* axis in (I). The C—H···O and C—H···π contacts are shown as orange and purple dashed lines, respectively.

(2-Bromoacetyl)ferrocene

Crystal data

[Fe(C₅H₅)(C₇H₆BrO)]

M_r = 306.97

Monoclinic, *P*2₁/*c*

Hall symbol: -P 2ybc

a = 7.7095 (3) Å

b = 9.6609 (4) Å

c = 14.7464 (7) Å

β = 98.061 (4)°

V = 1087.47 (8) Å³

Z = 4

F(000) = 608

D_x = 1.875 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2673 reflections

θ = 2.5–27.5°

μ = 5.03 mm⁻¹

T = 100 K

Plate, brown

0.30 × 0.10 × 0.03 mm

Data collection

Agilent SuperNova Dual
diffractometer with Atlas detector
Radiation source: SuperNova (Mo) X-ray
Source
Mirror monochromator
Detector resolution: 10.4041 pixels mm⁻¹
 ω scan
Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2010)

$T_{\min} = 0.314$, $T_{\max} = 0.864$
7692 measured reflections
2497 independent reflections
2004 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$
 $\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -7 \rightarrow 10$
 $k = -12 \rightarrow 12$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.079$
 $S = 1.02$
2497 reflections
136 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0276P)^2 + 0.4313P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.73 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.73500 (4)	0.70286 (3)	0.88244 (3)	0.02647 (12)
Fe1	0.18625 (5)	0.71721 (4)	0.60388 (3)	0.01114 (12)
O1	0.4210 (3)	0.5296 (2)	0.80778 (15)	0.0200 (5)
C1	0.3624 (4)	0.8207 (3)	0.5363 (2)	0.0201 (7)
H1	0.4199	0.9111	0.5548	0.024*
C2	0.2000 (4)	0.8024 (3)	0.4777 (2)	0.0186 (7)
H2	0.1235	0.8781	0.4483	0.022*
C3	0.1643 (4)	0.6583 (3)	0.4698 (2)	0.0160 (6)
H3	0.0591	0.6149	0.4333	0.019*
C4	0.3048 (4)	0.5867 (3)	0.5228 (2)	0.0174 (7)
H4	0.3161	0.4840	0.5301	0.021*
C5	0.4263 (4)	0.6876 (3)	0.5640 (2)	0.0218 (7)
H5	0.5375	0.6678	0.6055	0.026*
C6	0.1556 (4)	0.8318 (3)	0.7165 (2)	0.0154 (6)
H6	0.2173	0.9202	0.7359	0.019*
C7	-0.0085 (4)	0.8188 (3)	0.6598 (2)	0.0191 (7)
H7	-0.0807	0.8971	0.6307	0.023*
C8	-0.0498 (4)	0.6754 (3)	0.6484 (2)	0.0181 (7)
H8	-0.1560	0.6362	0.6103	0.022*
C9	0.0873 (4)	0.5979 (3)	0.6989 (2)	0.0155 (6)
H9	0.0938	0.4947	0.7034	0.019*
C10	0.2161 (4)	0.6937 (3)	0.7418 (2)	0.0140 (6)
C11	0.3856 (4)	0.6510 (3)	0.7928 (2)	0.0145 (6)
C12	0.5074 (4)	0.7682 (3)	0.8252 (2)	0.0161 (7)
H12B	0.4538	0.8252	0.8697	0.019*

H12A	0.5232	0.8278	0.7723	0.019*
------	--------	--------	--------	--------

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.01576 (18)	0.0226 (2)	0.0391 (2)	0.00046 (12)	-0.00282 (16)	-0.00281 (15)
Fe1	0.0115 (2)	0.0123 (2)	0.0098 (2)	0.00045 (16)	0.00201 (18)	-0.00059 (17)
O1	0.0241 (12)	0.0117 (12)	0.0221 (13)	0.0019 (9)	-0.0036 (10)	-0.0007 (9)
C1	0.0239 (17)	0.0225 (17)	0.0161 (16)	-0.0087 (14)	0.0102 (15)	-0.0031 (14)
C2	0.0256 (18)	0.0183 (17)	0.0128 (16)	0.0034 (13)	0.0063 (14)	0.0024 (13)
C3	0.0206 (16)	0.0173 (16)	0.0098 (15)	0.0000 (13)	0.0005 (13)	-0.0041 (13)
C4	0.0176 (15)	0.0181 (16)	0.0170 (16)	0.0055 (12)	0.0040 (14)	-0.0004 (13)
C5	0.0124 (15)	0.036 (2)	0.0177 (17)	0.0007 (14)	0.0029 (14)	-0.0019 (15)
C6	0.0184 (16)	0.0116 (15)	0.0165 (16)	0.0019 (12)	0.0034 (14)	-0.0038 (13)
C7	0.0183 (16)	0.0231 (17)	0.0163 (16)	0.0099 (13)	0.0036 (14)	-0.0005 (14)
C8	0.0130 (15)	0.0247 (17)	0.0178 (17)	-0.0023 (13)	0.0066 (14)	-0.0057 (14)
C9	0.0175 (15)	0.0178 (16)	0.0122 (15)	-0.0033 (12)	0.0056 (13)	0.0008 (13)
C10	0.0173 (15)	0.0170 (16)	0.0080 (14)	-0.0014 (12)	0.0031 (13)	-0.0001 (12)
C11	0.0208 (16)	0.0155 (16)	0.0071 (14)	-0.0013 (13)	0.0019 (13)	-0.0016 (12)
C12	0.0138 (15)	0.0175 (16)	0.0160 (16)	0.0025 (12)	-0.0020 (13)	0.0000 (13)

Geometric parameters (\AA , $^\circ$)

Br1—C12	1.943 (3)	C3—H3	1.0000
Fe1—C1	2.053 (3)	C4—C5	1.428 (4)
Fe1—C2	2.051 (3)	C4—H4	1.0000
Fe1—C3	2.042 (3)	C5—H5	1.0000
Fe1—C4	2.040 (3)	C6—C7	1.421 (5)
Fe1—C5	2.040 (3)	C6—C10	1.445 (4)
Fe1—C6	2.037 (3)	C6—H6	1.0000
Fe1—C7	2.060 (3)	C7—C8	1.426 (4)
Fe1—C8	2.059 (3)	C7—H7	1.0000
Fe1—C9	2.043 (3)	C8—C9	1.418 (4)
Fe1—C10	2.027 (3)	C8—H8	1.0000
O1—C11	1.217 (3)	C9—C10	1.438 (4)
C1—C5	1.416 (4)	C9—H9	1.0000
C1—C2	1.429 (5)	C10—C11	1.473 (4)
C1—H1	1.0000	C11—C12	1.506 (4)
C2—C3	1.421 (4)	C12—H12B	0.9900
C2—H2	1.0000	C12—H12A	0.9900
C3—C4	1.423 (4)		
C10—Fe1—C6	41.65 (12)	C4—C3—C2	107.9 (3)
C10—Fe1—C4	121.43 (12)	C4—C3—Fe1	69.53 (17)
C6—Fe1—C4	157.89 (13)	C2—C3—Fe1	70.02 (17)
C10—Fe1—C5	107.09 (13)	C4—C3—H3	126.1
C6—Fe1—C5	121.68 (13)	C2—C3—H3	126.1
C4—Fe1—C5	40.97 (12)	Fe1—C3—H3	126.1

C10—Fe1—C3	157.32 (12)	C3—C4—C5	107.8 (3)
C6—Fe1—C3	159.69 (12)	C3—C4—Fe1	69.65 (16)
C4—Fe1—C3	40.82 (12)	C5—C4—Fe1	69.50 (17)
C5—Fe1—C3	68.70 (13)	C3—C4—H4	126.1
C10—Fe1—C9	41.37 (12)	C5—C4—H4	126.1
C6—Fe1—C9	69.52 (12)	Fe1—C4—H4	126.1
C4—Fe1—C9	107.11 (12)	C1—C5—C4	108.4 (3)
C5—Fe1—C9	124.08 (13)	C1—C5—Fe1	70.27 (18)
C3—Fe1—C9	121.24 (12)	C4—C5—Fe1	69.53 (17)
C10—Fe1—C2	160.42 (12)	C1—C5—H5	125.8
C6—Fe1—C2	123.26 (12)	C4—C5—H5	125.8
C4—Fe1—C2	68.40 (12)	Fe1—C5—H5	125.8
C5—Fe1—C2	68.27 (13)	C7—C6—C10	107.4 (3)
C3—Fe1—C2	40.63 (11)	C7—C6—Fe1	70.56 (18)
C9—Fe1—C2	156.99 (13)	C10—C6—Fe1	68.79 (16)
C10—Fe1—C1	123.48 (13)	C7—C6—H6	126.3
C6—Fe1—C1	106.94 (13)	C10—C6—H6	126.3
C4—Fe1—C1	68.62 (13)	Fe1—C6—H6	126.3
C5—Fe1—C1	40.49 (13)	C6—C7—C8	108.7 (3)
C3—Fe1—C1	68.68 (13)	C6—C7—Fe1	68.85 (17)
C9—Fe1—C1	160.67 (13)	C8—C7—Fe1	69.73 (16)
C2—Fe1—C1	40.75 (13)	C6—C7—H7	125.6
C10—Fe1—C7	68.82 (12)	C8—C7—H7	125.6
C6—Fe1—C7	40.58 (13)	Fe1—C7—H7	125.6
C4—Fe1—C7	160.14 (13)	C9—C8—C7	108.2 (3)
C5—Fe1—C7	157.53 (13)	C9—C8—Fe1	69.15 (16)
C3—Fe1—C7	123.65 (13)	C7—C8—Fe1	69.75 (17)
C9—Fe1—C7	68.35 (12)	C9—C8—H8	125.9
C2—Fe1—C7	107.77 (13)	C7—C8—H8	125.9
C1—Fe1—C7	122.06 (13)	Fe1—C8—H8	125.9
C10—Fe1—C8	68.87 (13)	C8—C9—C10	108.0 (3)
C6—Fe1—C8	68.78 (12)	C8—C9—Fe1	70.40 (17)
C4—Fe1—C8	123.64 (12)	C10—C9—Fe1	68.70 (16)
C5—Fe1—C8	160.53 (13)	C8—C9—H9	126.0
C3—Fe1—C8	107.26 (13)	C10—C9—H9	126.0
C9—Fe1—C8	40.45 (12)	Fe1—C9—H9	126.0
C2—Fe1—C8	121.89 (14)	C9—C10—C6	107.6 (3)
C1—Fe1—C8	157.56 (13)	C9—C10—C11	123.6 (3)
C7—Fe1—C8	40.52 (12)	C6—C10—C11	128.4 (3)
C5—C1—C2	107.6 (3)	C9—C10—Fe1	69.93 (17)
C5—C1—Fe1	69.25 (17)	C6—C10—Fe1	69.57 (17)
C2—C1—Fe1	69.53 (17)	C11—C10—Fe1	120.8 (2)
C5—C1—H1	126.2	O1—C11—C10	121.5 (3)
C2—C1—H1	126.2	O1—C11—C12	123.6 (3)
Fe1—C1—H1	126.2	C10—C11—C12	114.9 (3)
C3—C2—C1	108.3 (3)	C11—C12—Br1	112.2 (2)
C3—C2—Fe1	69.35 (17)	C11—C12—H12B	109.2
C1—C2—Fe1	69.72 (18)	Br1—C12—H12B	109.2

C3—C2—H2	125.8	C11—C12—H12A	109.2
C1—C2—H2	125.8	Br1—C12—H12A	109.2
Fe1—C2—H2	125.8	H12B—C12—H12A	107.9
C10—Fe1—C1—C5	76.6 (2)	C9—Fe1—C6—C10	38.23 (17)
C6—Fe1—C1—C5	119.3 (2)	C2—Fe1—C6—C10	-163.33 (18)
C4—Fe1—C1—C5	-37.78 (19)	C1—Fe1—C6—C10	-121.69 (19)
C3—Fe1—C1—C5	-81.8 (2)	C7—Fe1—C6—C10	118.5 (3)
C9—Fe1—C1—C5	42.9 (5)	C8—Fe1—C6—C10	81.65 (19)
C2—Fe1—C1—C5	-119.1 (3)	C10—C6—C7—C8	-0.7 (3)
C7—Fe1—C1—C5	161.02 (19)	Fe1—C6—C7—C8	58.4 (2)
C8—Fe1—C1—C5	-165.4 (3)	C10—C6—C7—Fe1	-59.2 (2)
C10—Fe1—C1—C2	-164.34 (17)	C10—Fe1—C7—C6	38.79 (18)
C6—Fe1—C1—C2	-121.66 (19)	C4—Fe1—C7—C6	163.8 (3)
C4—Fe1—C1—C2	81.30 (19)	C5—Fe1—C7—C6	-44.8 (4)
C5—Fe1—C1—C2	119.1 (3)	C3—Fe1—C7—C6	-162.66 (18)
C3—Fe1—C1—C2	37.33 (17)	C9—Fe1—C7—C6	83.4 (2)
C9—Fe1—C1—C2	162.0 (3)	C2—Fe1—C7—C6	-120.74 (19)
C7—Fe1—C1—C2	-79.9 (2)	C1—Fe1—C7—C6	-78.3 (2)
C8—Fe1—C1—C2	-46.3 (4)	C8—Fe1—C7—C6	120.6 (3)
C5—C1—C2—C3	0.2 (4)	C10—Fe1—C7—C8	-81.9 (2)
Fe1—C1—C2—C3	-58.8 (2)	C6—Fe1—C7—C8	-120.6 (3)
C5—C1—C2—Fe1	59.0 (2)	C4—Fe1—C7—C8	43.1 (5)
C10—Fe1—C2—C3	162.0 (3)	C5—Fe1—C7—C8	-165.4 (3)
C6—Fe1—C2—C3	-163.31 (17)	C3—Fe1—C7—C8	76.7 (2)
C4—Fe1—C2—C3	37.94 (18)	C9—Fe1—C7—C8	-37.27 (19)
C5—Fe1—C2—C3	82.18 (19)	C2—Fe1—C7—C8	118.6 (2)
C9—Fe1—C2—C3	-45.0 (4)	C1—Fe1—C7—C8	161.04 (19)
C1—Fe1—C2—C3	119.8 (3)	C6—C7—C8—C9	0.7 (3)
C7—Fe1—C2—C3	-121.34 (19)	Fe1—C7—C8—C9	58.6 (2)
C8—Fe1—C2—C3	-79.1 (2)	C6—C7—C8—Fe1	-57.9 (2)
C10—Fe1—C2—C1	42.2 (4)	C10—Fe1—C8—C9	-38.11 (18)
C6—Fe1—C2—C1	76.9 (2)	C6—Fe1—C8—C9	-82.9 (2)
C4—Fe1—C2—C1	-81.9 (2)	C4—Fe1—C8—C9	76.4 (2)
C5—Fe1—C2—C1	-37.65 (18)	C5—Fe1—C8—C9	43.4 (5)
C3—Fe1—C2—C1	-119.8 (3)	C3—Fe1—C8—C9	118.20 (19)
C9—Fe1—C2—C1	-164.8 (3)	C2—Fe1—C8—C9	160.24 (18)
C7—Fe1—C2—C1	118.8 (2)	C1—Fe1—C8—C9	-166.0 (3)
C8—Fe1—C2—C1	161.02 (18)	C7—Fe1—C8—C9	-119.8 (3)
C1—C2—C3—C4	-0.4 (3)	C10—Fe1—C8—C7	81.7 (2)
Fe1—C2—C3—C4	-59.4 (2)	C6—Fe1—C8—C7	36.90 (19)
C1—C2—C3—Fe1	59.0 (2)	C4—Fe1—C8—C7	-163.81 (19)
C10—Fe1—C3—C4	-45.4 (4)	C5—Fe1—C8—C7	163.2 (4)
C6—Fe1—C3—C4	162.8 (3)	C3—Fe1—C8—C7	-121.98 (19)
C5—Fe1—C3—C4	37.96 (18)	C9—Fe1—C8—C7	119.8 (3)
C9—Fe1—C3—C4	-79.9 (2)	C2—Fe1—C8—C7	-79.9 (2)
C2—Fe1—C3—C4	119.0 (3)	C1—Fe1—C8—C7	-46.2 (4)
C1—Fe1—C3—C4	81.6 (2)	C7—C8—C9—C10	-0.4 (3)

C7—Fe1—C3—C4	-163.30 (18)	Fe1—C8—C9—C10	58.58 (19)
C8—Fe1—C3—C4	-121.84 (18)	C7—C8—C9—Fe1	-59.0 (2)
C10—Fe1—C3—C2	-164.4 (3)	C10—Fe1—C9—C8	119.4 (3)
C6—Fe1—C3—C2	43.8 (4)	C6—Fe1—C9—C8	80.94 (19)
C4—Fe1—C3—C2	-119.0 (3)	C4—Fe1—C9—C8	-122.16 (19)
C5—Fe1—C3—C2	-81.0 (2)	C5—Fe1—C9—C8	-163.95 (19)
C9—Fe1—C3—C2	161.13 (19)	C3—Fe1—C9—C8	-79.8 (2)
C1—Fe1—C3—C2	-37.43 (19)	C2—Fe1—C9—C8	-47.3 (4)
C7—Fe1—C3—C2	77.7 (2)	C1—Fe1—C9—C8	163.8 (3)
C8—Fe1—C3—C2	119.16 (19)	C7—Fe1—C9—C8	37.33 (18)
C2—C3—C4—C5	0.5 (3)	C6—Fe1—C9—C10	-38.48 (17)
Fe1—C3—C4—C5	-59.3 (2)	C4—Fe1—C9—C10	118.42 (18)
C2—C3—C4—Fe1	59.7 (2)	C5—Fe1—C9—C10	76.6 (2)
C10—Fe1—C4—C3	161.21 (17)	C3—Fe1—C9—C10	160.74 (17)
C6—Fe1—C4—C3	-164.1 (3)	C2—Fe1—C9—C10	-166.7 (3)
C5—Fe1—C4—C3	-119.1 (3)	C1—Fe1—C9—C10	44.4 (4)
C9—Fe1—C4—C3	118.28 (18)	C7—Fe1—C9—C10	-82.09 (19)
C2—Fe1—C4—C3	-37.77 (18)	C8—Fe1—C9—C10	-119.4 (3)
C1—Fe1—C4—C3	-81.7 (2)	C8—C9—C10—C6	0.0 (3)
C7—Fe1—C4—C3	44.8 (4)	Fe1—C9—C10—C6	59.60 (19)
C8—Fe1—C4—C3	77.0 (2)	C8—C9—C10—C11	-174.0 (3)
C10—Fe1—C4—C5	-79.7 (2)	Fe1—C9—C10—C11	-114.3 (3)
C6—Fe1—C4—C5	-45.1 (4)	C8—C9—C10—Fe1	-59.6 (2)
C3—Fe1—C4—C5	119.1 (3)	C7—C6—C10—C9	0.5 (3)
C9—Fe1—C4—C5	-122.7 (2)	Fe1—C6—C10—C9	-59.82 (19)
C2—Fe1—C4—C5	81.3 (2)	C7—C6—C10—C11	174.0 (3)
C1—Fe1—C4—C5	37.35 (19)	Fe1—C6—C10—C11	113.7 (3)
C7—Fe1—C4—C5	163.8 (3)	C7—C6—C10—Fe1	60.3 (2)
C8—Fe1—C4—C5	-163.94 (19)	C6—Fe1—C10—C9	118.7 (2)
C2—C1—C5—C4	0.1 (3)	C4—Fe1—C10—C9	-80.1 (2)
Fe1—C1—C5—C4	59.2 (2)	C5—Fe1—C10—C9	-122.53 (18)
C2—C1—C5—Fe1	-59.2 (2)	C3—Fe1—C10—C9	-47.0 (4)
C3—C4—C5—C1	-0.3 (3)	C2—Fe1—C10—C9	164.4 (3)
Fe1—C4—C5—C1	-59.7 (2)	C1—Fe1—C10—C9	-163.89 (18)
C3—C4—C5—Fe1	59.4 (2)	C7—Fe1—C10—C9	80.88 (19)
C10—Fe1—C5—C1	-121.9 (2)	C8—Fe1—C10—C9	37.29 (17)
C6—Fe1—C5—C1	-78.7 (2)	C4—Fe1—C10—C6	161.21 (18)
C4—Fe1—C5—C1	119.5 (3)	C5—Fe1—C10—C6	118.76 (19)
C3—Fe1—C5—C1	81.7 (2)	C3—Fe1—C10—C6	-165.7 (3)
C9—Fe1—C5—C1	-164.20 (19)	C9—Fe1—C10—C6	-118.7 (2)
C2—Fe1—C5—C1	37.88 (19)	C2—Fe1—C10—C6	45.7 (4)
C7—Fe1—C5—C1	-46.2 (4)	C1—Fe1—C10—C6	77.4 (2)
C8—Fe1—C5—C1	163.2 (4)	C7—Fe1—C10—C6	-37.82 (18)
C10—Fe1—C5—C4	118.55 (19)	C8—Fe1—C10—C6	-81.41 (19)
C6—Fe1—C5—C4	161.75 (18)	C6—Fe1—C10—C11	-123.4 (3)
C3—Fe1—C5—C4	-37.82 (18)	C4—Fe1—C10—C11	37.8 (3)
C9—Fe1—C5—C4	76.3 (2)	C5—Fe1—C10—C11	-4.6 (3)
C2—Fe1—C5—C4	-81.6 (2)	C3—Fe1—C10—C11	70.9 (4)

C1—Fe1—C5—C4	−119.5 (3)	C9—Fe1—C10—C11	117.9 (3)
C7—Fe1—C5—C4	−165.7 (3)	C2—Fe1—C10—C11	−77.7 (4)
C8—Fe1—C5—C4	43.7 (5)	C1—Fe1—C10—C11	−46.0 (3)
C10—Fe1—C6—C7	−118.5 (3)	C7—Fe1—C10—C11	−161.2 (3)
C4—Fe1—C6—C7	−165.4 (3)	C8—Fe1—C10—C11	155.2 (3)
C5—Fe1—C6—C7	161.56 (19)	C9—C10—C11—O1	−5.8 (4)
C3—Fe1—C6—C7	45.6 (4)	C6—C10—C11—O1	−178.4 (3)
C9—Fe1—C6—C7	−80.3 (2)	Fe1—C10—C11—O1	−91.0 (3)
C2—Fe1—C6—C7	78.2 (2)	C9—C10—C11—C12	174.9 (3)
C1—Fe1—C6—C7	119.8 (2)	C6—C10—C11—C12	2.3 (4)
C8—Fe1—C6—C7	−36.84 (18)	Fe1—C10—C11—C12	89.7 (3)
C4—Fe1—C6—C10	−46.9 (4)	O1—C11—C12—Br1	5.7 (4)
C5—Fe1—C6—C10	−80.0 (2)	C10—C11—C12—Br1	−175.0 (2)
C3—Fe1—C6—C10	164.1 (3)		

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C1—C5 ring.

D—H···A	D—H	H···A	D···A	D—H···A
C1—H1···O1 ⁱ	1.00	2.49	3.327 (4)	140
C12—H12a···O1 ⁱ	0.99	2.35	3.291 (4)	158
C12—H12b···Cg1 ⁱⁱ	0.99	2.64	3.445 (3)	139

Symmetry codes: (i) $-x+1, y+1/2, -z+3/2$; (ii) $x, -y+1/2, z-1/2$.