

An orthorhombic polymorph of pyrazino-[2,3-*f*][1,10]phenanthroline-2,3-dicarbonitrile

Wei Yang,^a Qi-Ming Qiu,^a Li-Li Zhou,^b Qiong-Hua Jin^{a*} and Cun-Lin Zhang^c

^aDepartment of Chemistry, Capital Normal University, Beijing 100048, People's Republic of China, ^bResearch Center for Import–Export Chemicals Safety of the General Administration of Quality Supervision, Inspection and Quarantine of the People's Republic of China (AQSIQ), Chinese Academy of Inspection and Quarantine, Beijing 100123, People's Republic of China, and ^cKey Laboratory of Terahertz Optoelectronics, Ministry of Education, Department of Physics, Capital Normal University, Beijing 100048, People's Republic of China
Correspondence e-mail: jinqh204@163.com


Received 5 October 2011; accepted 7 November 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}–\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.119; data-to-parameter ratio = 11.1.

The title compound, $\text{C}_{16}\text{H}_6\text{N}_6$, is a polymorph of the previously reported structure [Kozlov & Goldberg (2008). *Acta Cryst.* **C64**, o498–o501]. Unlike the previously reported monoclinic polymorph (space group $P2_1/c$, $Z = 8$), the title compound reveals orthorhombic symmetry (space group $Pnma$, $Z = 4$). The molecule shows crystallographic mirror symmetry, while the previously reported structure exhibits two independent molecules per asymmetric unit. In the title compound, adjacent molecules are essentially parallel along the c axis and tend to be vertical along the b axis with dihedral angles of 72.02 (6)°. However, in the reported polymorph, the entire crystal structure shows an antiparallel arrangement of adjacent columns related by inversion centers and the two independent molecules are nearly parallel with a dihedral angle of 2.48 (6)°.

Related literature

For ligands based on 1,10-phenanthroline in coordination chemistry, see: Rabaca *et al.* (2008); Stephenson *et al.* (2008). For reports of the title compound in coordination chemistry, see: Kulkarni *et al.* (2004); Stephenson & Hardie (2006); Xiao *et al.* (2011); Xu *et al.* (2002). For examples of polymorphism, see: Demirtaş *et al.* (2011); Jiang *et al.* (2000); Okabe *et al.* (2001); Pan & Chen (2009); Ramos Silva *et al.* (2011); Thal-lapally *et al.* (2004). For the previously reported polymorph, see: Kozlov & Goldberg (2008). For related structures, see: Kozlov *et al.* (2008).



Experimental

Crystal data

$\text{C}_{16}\text{H}_6\text{N}_6$	$V = 1214.00$ (18) Å ³
$M_r = 282.27$	$Z = 4$
Orthorhombic, $Pnma$	Mo $K\alpha$ radiation
$a = 14.1055$ (13) Å	$\mu = 0.10$ mm ⁻¹
$b = 16.3331$ (14) Å	$T = 293$ K
$c = 5.2694$ (4) Å	$0.45 \times 0.30 \times 0.26$ mm

Data collection

Bruker SMART CCD area-detector diffractometer	4543 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2007)	1113 independent reflections
$T_{\min} = 0.956$, $T_{\max} = 0.974$	759 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	100 parameters
$wR(F^2) = 0.119$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\max} = 0.20$ e Å ⁻³
1113 reflections	$\Delta\rho_{\min} = -0.24$ e Å ⁻³

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT-Plus* (Bruker, 2007); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *ORTEPIII* (Burnett & Johnson, 1996); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Natural Science Foundation of China (No. 21171119). It was granted by the research project of recycling PET of food contact materials (No. 2010 J K022) from the Chinese Academy of Inspection and Quarantine, National Keystone Basic Research Program (973 Program) under grant Nos. 2007CB310408 and 2006CB302901, and the Funding Project for Academic Human Resources Development in Institutions of Higher Learning Under the Jurisdiction of Beijing Municipality.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2127).

References

- Bruker (2007). *SMART, SAINT-Plus and SADABS*. Bruker AXS Inc., Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. OakRidge National Laboratory, Tennessee, USA.
- Demirtaş, G., Dege, N. & Büyükgüngör, O. (2011). *Acta Cryst. E* **67**, o1509–o1510.
- Jiang, R.-W., Ming, D.-S., But, P. P. H. & Mak, T. C. W. (2000). *Acta Cryst. C* **56**, 594–595.
- Kozlov, L. & Goldberg, I. (2008). *Acta Cryst. C* **64**, o498–o501.
- Kozlov, L., Rubin-Preminger, J. M. & Goldberg, I. (2008). *Acta Cryst. C* **64**, o1–o3.
- Kulkarni, M. S., Rao, B. S. M., Sastri, C. V., Maiya, B. G., Mohan, H. & Mittal, J. P. (2004). *J. Photochem. Photobiol. A Chem.* **167**, 101–109.
- Okabe, N., Kyoyama, H. & Suzuki, M. (2001). *Acta Cryst. E* **57**, o764–o766.
- Pan, J. X. & Chen, Q. W. (2009). *Acta Cryst. E* **65**, o652.
- Rabaca, S., Duarte, M. C., Santos, I. C., Pereira, L. C. J., Fourmigue, M., Henriques, R. T. & Almeida, M. (2008). *Polyhedron*, **27**, 1999–2006.
- Ramos Silva, M., Pereira Silva, P. S., Yuste, C. & Paixão, J. A. (2011). *Acta Cryst. E* **67**, o340.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stephenson, M. D. & Hardie, M. J. (2006). *Cryst. Growth Des.* **6**(2), 423–432.
- Stephenson, M. D., Prior, T. J. & Hardie, M. J. (2008). *Cryst. Growth Des.* **8**, 643–653.
- Thallapally, P. K., Jetti, R. K. R., Katz, A. K., Carrell, H. L., Singh, K., Lahiri, K., Kotha, S., Boese, R. & Desiraju, G. R. (2004). *Angew. Chem. Int. Ed.* **43**, 1149–1155.
- Xiao, Y. L., Jin, Q. H., Deng, Y. H., Li, Z. F., Yang, W., Wu, M. H. & Zhang, C. L. (2011). *Inorg. Chem. Commun.* Accepted, doi:10.1016/j.inoche.2011.10.011.
- Xu, Z. D., Liu, H., Wang, M., Xiao, S. L., Yang, M. & Bu, X. H. (2002). *J. Inorg. Biochem.* **92**, 149–155.

supporting information

Acta Cryst. (2011). E67, o3250–o3251 [https://doi.org/10.1107/S1600536811047039]

An orthorhombic polymorph of pyrazino[2,3-*f*][1,10]phenanthroline-2,3-dicarbonitrile

Wei Yang, Qi-Ming Qiu, Li-Li Zhou, Qiong-Hua Jin and Cun-Lin Zhang

S1. Comment

Ligands based on 1,10-phenanthroline are attractive building blocks for the formation of metal-organic complexes bearing diverse dimensionalities (Rabaca *et al.*, 2008; Stephenson *et al.*, 2008). Among 1,10-phenanthroline derivatives, the DICNQ (DICNQ = pyrazino[2,3-*f*][1,10]phenanthroline-2,3-dicarbonitrile) (Xu *et al.*, 2002; Kulkarni *et al.*, 2004) has a big conjugated system and multiple coordination including strong chelating phenanthroline ring, bridging pyrazine ring and cyanite groups, so it is good in synthesizing luminescent complexes containing multifunctional ligands. Up to date only a few complexes of DICNQ are reported, such as [Co(dicnq)₂Br₂] and [Cu(dicnq)₂Br₂][Cu(dicnq)₂Br]Br·(CH₃CN)·2(H₂O) (Stephenson & Hardie, 2006). Very recently, we synthesized five Cu(I) complexes [Cu(PPh₃)₂(C₁₆H₆N₆)X] (C₁₆H₆N₆ = pyrazino[2,3-*f*][1,10]phenanthroline-2,3-dicarbonitrile; X = Cl, Br, I, CN, SCN) (Xiao *et al.*, 2011). During our attempts to synthesize Cu(II) complexes of DICNQ and 4,4'-bipyridine, we obtained crystals of a new orthorhombic polymorph of the solvent-free DICNQ.

Polymorphism can potentially be found in any crystalline material including polymers, minerals, and metals, and is related to allotropy, which refers to elemental solids. When polymorphism exists as a result of difference in crystal packing, it is called packing polymorphism. The different crystal types probably are the result of hydration, solvation or the effect of synthesis methods. There are recently reported examples of organic polymorphs: 2-(pyrimidin-2-ylsulfanyl)acetic acid, which is able to form monoclinic and orthorhombic crystals (Ramos Silva *et al.*, 2011; Pan & Chen, 2009). 3,4,5-Trihydroxybenzoic acid monohydrate is found to form three polymorphs (Demirtaş *et al.*, 2011; Jiang *et al.*, 2000; Okabe *et al.*, 2001). We herein report a new orthorhombic polymorph of DICNQ (1).

ORTEP (Burnett & Johnson, 1996) representation of the title compound (1) is shown in Fig. 1. The title compound belongs to orthorhombic system, space group *Pnma*, consisting of one symmetric unit of single molecule, unlike the reported monoclinic polymorph (space group *P2₁/c*), (2), (Kozlov & Goldberg, 2008), which consists of two crystallographically independent molecules.

The phenanthroline fragment of both polymorphs is nearly planar, and a similar bending of the cyano groups from the plane of the phenanthroline residues can be observed. The deviating distance of the cyano groups from the plane in (1) (0.182 (7) Å) is obviously bigger than the corresponding deviating distances in (2) (0.0345 Å and 0.0374 Å).

The packing pattern differs from the reported polymorph. In the title compound, there are two packing arrangements (Fig. 2), adjacent molecules are essentially parallel along the *c* axis and tend to be vertical along the *b* axis with dihedral angles of 72.02 (6)°. However, in the reported polymorph the entire crystal structure shows an antiparallel arrangement of the adjacent columns related by inversion centers, and the two independent molecules are nearly parallel with a dihedral angle of 2.48 (6)°.

In addition, the intermolecular assembly of (1) is dominated by π - π stacking of overlaps among phenanthroline fragments and π - π stacking between phenanthroline fragment and cyano group of adjacent molecules (Fig. 2), with an interplanar distance between the parallel phenanthroline rings in the two adjacent molecules of 3.030 (1) Å. In contrast, in (2) and the ethanol solvate of DICNQ ($C_{16}H_6N_6$).(C₂H₆O) (3) (Kozlov *et al.*, 2008), adjacent molecules overlap only through their phenanthroline fragments with the shortest interplanar distance of 3.380 (3) Å for (2), 3.27 Å and 3.40 Å for (3).

Various methods and conditions of the crystallization process are the main reason responsible for the formation of different polymorphic forms (Thallapally *et al.*, 2004). The two polymorphs (1) and (2) are prepared under different catalysts. (1) was prepared under the catalysis of CuCl₂.2H₂O and 4,4'-bipy, while (2) was prepared under the catalysis of [Pd(PPh₃)Cl₂].

S2. Experimental

A mixture of CuCl₂.2H₂O (0.0178 g, 0.1 mmol) in 5 ml CH₂Cl₂ and DICNQ (0.0290 g, 0.1 mmol) in 3 ml CH₃CN was stirred for 30 min, then a solution of 4,4'-bipyridine (0.0202 g, 0.1 mmol) in 2 ml CH₃CN was added under continuous stirring. The resulting solution was refluxed for 5 h and filtered. The filtrate was allowed to evaporate slowly at room temperature to yield yellow block crystals of the title complex. Crystals suitable for single-crystal X-ray diffraction were selected directly from the sample as prepared.

IR (KBr, disc: ν/cm^{-1}): 3466.82 s, 1587.57m, 1571.17m, 1505.59m, 1459.84m, 1444.23m, 1384.77m, 1372.26m, 1331.97w, 1264.07w, 1219.29m, 1141.93m, 1123.84w, 1073.56w, 1028.64w, 975.09w, 817.56m, 743.23m, 688.54w, 618.37w, 527.98w, 441.00w, 420.02w, 409.42w.

S3. Refinement

H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 Å and with $U_{iso}(H) = 1.2U_{eq}(C)$.




Figure 1

Perspective view of a basic unit of the title complex. Atoms are displayed as ellipsoids at the 50% probability level at *ca* 293 K.




Figure 2

The crystal packing of the title complex.

pyrazino[2,3-*f*][1,10]phenanthroline-2,3-dicarbonitrile

Crystal data

C₁₆H₆N₆ $M_r = 282.27$ Orthorhombic, *Pnma*

Hall symbol: -P 2ac 2n

 $a = 14.1055$ (13) Å $b = 16.3331$ (14) Å $c = 5.2694$ (4) Å $V = 1214.00$ (18) Å³ $Z = 4$ $F(000) = 576$ $D_x = 1.544$ Mg m⁻³Mo *K*α radiation, $\lambda = 0.71073$ Å

Cell parameters from 1184 reflections

 $\theta = 2.9$ – 27.5° $\mu = 0.10$ mm⁻¹ $T = 293$ K

Block, yellow

 $0.45 \times 0.30 \times 0.26$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scansAbsorption correction: multi-scan
(*SADABS*; Bruker, 2007) $T_{\min} = 0.956$, $T_{\max} = 0.974$

4543 measured reflections

1113 independent reflections

759 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.046$ $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.9^\circ$ $h = -16 \rightarrow 16$ $k = -19 \rightarrow 7$ $l = -6 \rightarrow 6$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.041$ $wR(F^2) = 0.119$ $S = 1.06$

1113 reflections

100 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0556P)^2 + 0.3018P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.20$ e Å⁻³ $\Delta\rho_{\min} = -0.24$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.56932 (11)	0.33317 (10)	-0.1078 (3)	0.0347 (5)
N2	0.77953 (11)	0.33585 (9)	0.6140 (3)	0.0322 (4)
N3	0.92249 (13)	0.37680 (13)	1.1146 (4)	0.0539 (6)
C1	0.61818 (12)	0.29508 (11)	0.0786 (3)	0.0279 (5)
C2	0.67134 (12)	0.33802 (11)	0.2603 (3)	0.0281 (5)

C3	0.67273 (13)	0.42373 (12)	0.2476 (4)	0.0349 (5)
H3	0.7060	0.4542	0.3673	0.042*
C4	0.62440 (14)	0.46176 (13)	0.0564 (4)	0.0381 (6)
H4	0.6249	0.5185	0.0422	0.046*
C5	0.57451 (13)	0.41428 (12)	-0.1161 (4)	0.0369 (5)
H5	0.5426	0.4410	-0.2463	0.044*
C6	0.72648 (12)	0.29328 (11)	0.4470 (3)	0.0285 (5)
C7	0.83106 (12)	0.29288 (12)	0.7773 (4)	0.0317 (5)
C8	0.88437 (14)	0.33918 (13)	0.9640 (4)	0.0372 (5)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0333 (9)	0.0368 (10)	0.0339 (10)	0.0021 (7)	-0.0025 (8)	0.0040 (8)
N2	0.0305 (9)	0.0357 (9)	0.0303 (9)	-0.0020 (7)	0.0014 (8)	-0.0007 (8)
N3	0.0476 (11)	0.0703 (14)	0.0437 (12)	-0.0084 (10)	-0.0041 (10)	-0.0142 (11)
C1	0.0249 (9)	0.0318 (10)	0.0271 (11)	0.0011 (8)	0.0041 (8)	0.0010 (9)
C2	0.0255 (9)	0.0291 (10)	0.0296 (11)	0.0002 (8)	0.0036 (8)	0.0018 (9)
C3	0.0335 (11)	0.0312 (10)	0.0401 (13)	-0.0031 (9)	-0.0007 (10)	-0.0031 (10)
C4	0.0353 (11)	0.0302 (10)	0.0488 (14)	0.0018 (9)	0.0026 (11)	0.0059 (10)
C5	0.0354 (11)	0.0370 (12)	0.0382 (12)	0.0052 (9)	-0.0005 (10)	0.0082 (10)
C6	0.0253 (9)	0.0319 (9)	0.0283 (10)	-0.0016 (8)	0.0028 (8)	-0.0029 (8)
C7	0.0264 (10)	0.0428 (11)	0.0261 (11)	-0.0027 (8)	0.0008 (9)	-0.0014 (9)
C8	0.0314 (11)	0.0480 (13)	0.0321 (12)	-0.0018 (9)	0.0013 (10)	-0.0013 (11)

Geometric parameters (Å, °)

N1—C5	1.327 (3)	C3—C4	1.366 (3)
N1—C1	1.352 (2)	C3—H3	0.9300
N2—C7	1.327 (2)	C4—C5	1.387 (3)
N2—C6	1.348 (2)	C4—H4	0.9300
N3—C8	1.138 (3)	C5—H5	0.9300
C1—C2	1.404 (2)	C6—C6 ⁱ	1.414 (4)
C1—C1 ⁱ	1.473 (4)	C7—C7 ⁱ	1.401 (4)
C2—C3	1.402 (3)	C7—C8	1.451 (3)
C2—C6	1.451 (2)		
C5—N1—C1	117.07 (17)	C3—C4—H4	120.6
C7—N2—C6	117.02 (16)	C5—C4—H4	120.6
N1—C1—C2	122.55 (17)	N1—C5—C4	124.37 (19)
N1—C1—C1 ⁱ	117.41 (10)	N1—C5—H5	117.8
C2—C1—C1 ⁱ	119.97 (11)	C4—C5—H5	117.8
C3—C2—C1	118.29 (17)	N2—C6—C6 ⁱ	121.05 (10)
C3—C2—C6	121.85 (17)	N2—C6—C2	118.69 (16)
C1—C2—C6	119.80 (17)	C6 ⁱ —C6—C2	120.23 (10)
C4—C3—C2	118.85 (18)	N2—C7—C7 ⁱ	121.93 (10)
C4—C3—H3	120.6	N2—C7—C8	116.61 (17)
C2—C3—H3	120.6	C7 ⁱ —C7—C8	121.41 (11)

C3—C4—C5	118.85 (18)	N3—C8—C7	177.0 (2)
C5—N1—C1—C2	1.0 (3)	C3—C4—C5—N1	0.7 (3)
C5—N1—C1—C1 ⁱ	-175.87 (13)	C7—N2—C6—C6 ⁱ	-0.18 (19)
N1—C1—C2—C3	0.5 (3)	C7—N2—C6—C2	-178.16 (15)
C1 ⁱ —C1—C2—C3	177.37 (12)	C3—C2—C6—N2	0.7 (3)
N1—C1—C2—C6	-176.57 (15)	C1—C2—C6—N2	177.73 (15)
C1 ⁱ —C1—C2—C6	0.26 (19)	C3—C2—C6—C6 ⁱ	-177.26 (13)
C1—C2—C3—C4	-1.5 (3)	C1—C2—C6—C6 ⁱ	-0.26 (19)
C6—C2—C3—C4	175.52 (17)	C6—N2—C7—C7 ⁱ	0.18 (19)
C2—C3—C4—C5	1.0 (3)	C6—N2—C7—C8	-177.14 (15)
C1—N1—C5—C4	-1.7 (3)		

Symmetry code: (i) $x, -y+1/2, z$.