

## catena-Poly[[bis( $\mu$ -2-formyl-6-methoxyphenolato-1:2 $\kappa^4$ O<sup>1</sup>,O<sup>6</sup>:O<sup>1</sup>,O<sup>2</sup>)copper(II)-sodium]- $\mu$ -tetrafluoridoborate-1:1' $\kappa^2$ F:F]

Yu Yang, Po Gao, Jing-Lin Yang, Hai-Ge Hou and Ting Gao\*

School of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China

Correspondence e-mail: gaoting1218@yahoo.com.cn


Received 16 November 2011; accepted 1 December 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$ ; disorder in main residue;  $R$  factor = 0.061; wR factor = 0.143; data-to-parameter ratio = 14.4.

In the title heterodinuclear complex,  $[\text{CuNa}(\text{BF}_4)(\text{C}_8\text{H}_7\text{O}_3)_2]_n$ , the Cu<sup>II</sup> ion is four-coordinated by four O atoms of two 2-formyl-6-methoxyphenolate ligands, giving rise to a square-planar geometry. The Na<sup>+</sup> ion is six-coordinated by four O atoms from the two ligands and two F atoms of two tetrafluoridoborate anions. The tetrafluoridoborate anion links the Na<sup>+</sup> ions, forming a one-dimensional structure along [001]. Three F atoms of the tetrafluoridoborate anion are disordered over two sets of sites, with an occupancy ratio of 0.790 (11):0.210 (11).

### Related literature

For related heterodinuclear complexes, see: Gao *et al.* (2011); Kajiwara *et al.* (2008).



### Experimental

#### Crystal data

$[\text{CuNa}(\text{BF}_4)(\text{C}_8\text{H}_7\text{O}_3)_2]$

$M_r = 475.62$

Monoclinic,  $P2_1/c$

$a = 9.932 (2)\text{ \AA}$

$b = 19.349 (4)\text{ \AA}$

$c = 9.940 (2)\text{ \AA}$

$\beta = 105.16 (3)^\circ$

$V = 1843.7 (7)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 1.28\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.21 \times 0.18 \times 0.16\text{ mm}$

#### Data collection

Rigaku R-AXIS RAPID diffractometer

Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.779$ ,  $T_{\max} = 0.820$

17729 measured reflections

4201 independent reflections

2278 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.100$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$

$wR(F^2) = 0.143$

$S = 1.04$

4201 reflections

292 parameters

30 restraints

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.50\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.34\text{ e \AA}^{-3}$

**Table 1**  
Selected bond lengths (Å).

Cu1—O2	1.888 (3)	Na1—O4	2.614 (4)
Cu1—O3	1.929 (3)	Na1—O5	2.370 (3)
Cu1—O5	1.889 (3)	Na1—F1	2.215 (7)
Cu1—O6	1.936 (3)	Na1—F3'	2.344 (17)
Na1—O1	2.615 (3)	Na1—F4'	2.243 (4)
Na1—O2	2.377 (3)		

Symmetry code: (i)  $x, -y + \frac{3}{2}, z + \frac{1}{2}$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors gratefully acknowledge financial support from Heilongjiang Province (11551334) and the Students Innovation Programme of Heilongjiang University (CX11073).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2490).

### References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Gao, P., Hou, H.-G., Gao, T., Yang, J.-L. & Yang, Y. (2011). *Acta Cryst. E67*, m1522.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Kajiwara, T., Nakano, M., Takaishi, S. & Yamashita, M. (2008). *Inorg. Chem.* **47**, 8604–8606.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalStructure*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

# supporting information

*Acta Cryst.* (2012). E68, m37 [doi:10.1107/S1600536811051713]

## **catena-Poly[[bis( $\mu$ -2-formyl-6-methoxy-phenolato-1:2 $\kappa^4$ O<sup>1</sup>,O<sup>6</sup>:O<sup>1</sup>,O<sup>2</sup>)copper(II)sodium]- $\mu$ -tetrafluoridoborate-1:1' $\kappa^2$ F:F']**

**Yu Yang, Po Gao, Jing-Lin Yang, Hai-Ge Hou and Ting Gao**

### **S1. Comment**


Orthovanillin is a commercial ligand that can yeild heterodinuclear 3d-4f complexes with two different coordination sites; 3d ions have a marked affinity for the inner O<sub>2</sub>O<sub>2</sub> site, whereas 4f ions prefer the larger outer O<sub>2</sub>O<sub>2</sub> coordination site (Kajiwara *et al.*, 2008). Recently, we were interested in the nature of the products obtained by reacting a 3d complex with alkali metal ions (Gao *et al.*, 2011). In this paper we reacted a Cu complex with sodium tetrafluoridoborate to yield a heterodinuclear complex. As shown in Fig. 1, the Cu<sup>II</sup> ion is four-coordinated by two aldehyde O atoms and two phenolate O atoms from two orthovanillin ligands (Table 1). The Cu<sup>II</sup> ion is inserted to inner cavity in a square-planar geometry. The Na<sup>+</sup> ion is ligated by two phenolate O atoms, two methoxyl O atoms and two F atoms from two tetrafluoridoborate anions. The Cu<sup>II</sup> and Na<sup>+</sup> ions are bridged by the phenolate O atoms.

### **S2. Experimental**

To a solution of *o*-vanillin (0.046 g, 0.20 mmol) in dichloromethane (5 ml) was added to a solution of copper(II) acetate monohydrate (0.040 g, 0.20 mmol) and sodium tetrafluoridoborate (0.034 g, 0.20 mmol) in ethanol (5 ml). The mixture was stirred, heated under reflux (30 min) and then allowed to cool to room temperature (yield: 70%). The crystals suitable for X-ray determination were obtained by slow diffusion of diethylether into the solution for one week. Analysis, calculated for C<sub>16</sub>H<sub>14</sub>BCuF<sub>4</sub>NaO<sub>6</sub>: C 40.40, H 2.97%; found: C 40.38, H 2.98%.

### **S3. Refinement**

Three F atoms were disordered each in two positions with an occupancy factors of 0.790 (11) for F1, F2, F3 and 0.210 (11) for F1', F2', F3'. H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93 (aromatic and aldehyde) and 0.96 (methyl) Å and with U<sub>iso</sub>(H) = 1.2(1.5 for methyl)U<sub>eq</sub>(C).

**Figure 1**

The molecular structure of the title compound, showing the 30% probability displacement ellipsoids. [Symmetry code: (i)  $x, 3/2-y, 1/2+z$ .]

**catena-Poly[[bis( $\mu$ -2-formyl-6-methoxyphenolato- 1:2 $\kappa^4\text{O}^1,\text{O}^6:\text{O}^1,\text{O}^2)$ copper(II)sodium]-  $\mu$ -tetrafluoridoborate-1:1' $\kappa^2\text{F}:F$ ]**

#### Crystal data

$[\text{CuNa}(\text{BF}_4)(\text{C}_8\text{H}_7\text{O}_3)_2]$

$M_r = 475.62$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.932$  (2) Å

$b = 19.349$  (4) Å

$c = 9.940$  (2) Å

$\beta = 105.16$  (3)°

$V = 1843.7$  (7) Å<sup>3</sup>

$Z = 4$

$F(000) = 956$

$D_x = 1.714 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 9827 reflections

$\theta = 3.0\text{--}27.5^\circ$

$\mu = 1.28 \text{ mm}^{-1}$

$T = 293$  K

Block, colorless

$0.21 \times 0.18 \times 0.16$  mm

*Data collection*

Rigaku R-AXIS RAPID  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(*ABSCOR*; Higashi, 1995)  
 $T_{\min} = 0.779$ ,  $T_{\max} = 0.820$

17729 measured reflections  
4201 independent reflections  
2278 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.100$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.0^\circ$   
 $h = -12 \rightarrow 12$   
 $k = -25 \rightarrow 25$   
 $l = -11 \rightarrow 12$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.143$   
 $S = 1.04$   
4201 reflections  
292 parameters  
30 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0582P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.50 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
F1'	0.821 (3)	0.8431 (15)	0.453 (4)	0.135 (10)	0.210 (11)
F2'	0.979 (4)	0.8027 (19)	0.428 (4)	0.174 (14)	0.210 (11)
F3'	0.874 (2)	0.7798 (9)	0.6030 (15)	0.075 (6)	0.210 (11)
F1	0.7789 (9)	0.8066 (5)	0.5476 (9)	0.144 (3)	0.790 (11)
F2	0.8679 (9)	0.8364 (4)	0.3784 (7)	0.132 (3)	0.790 (11)
F3	0.9995 (6)	0.7795 (3)	0.5612 (7)	0.131 (3)	0.790 (11)
F4	0.8288 (4)	0.72587 (18)	0.4087 (4)	0.1037 (13)	
B1	0.8662 (8)	0.7866 (3)	0.4740 (7)	0.0628 (17)	
C1	1.1275 (4)	0.9403 (3)	0.8798 (5)	0.0456 (11)	
C2	1.2649 (5)	0.9573 (3)	0.9383 (5)	0.0550 (13)	
H2	1.3310	0.9224	0.9656	0.066*	
C3	1.3060 (5)	1.0264 (3)	0.9568 (5)	0.0615 (14)	
H3	1.3991	1.0372	0.9971	0.074*	
C4	1.2118 (5)	1.0780 (3)	0.9166 (5)	0.0549 (13)	
H4	1.2412	1.1238	0.9266	0.066*	
C5	1.0682 (4)	1.0625 (2)	0.8591 (4)	0.0427 (11)	
C6	1.0246 (4)	0.9926 (2)	0.8378 (4)	0.0402 (10)	
C7	0.9744 (5)	1.1182 (3)	0.8252 (4)	0.0503 (12)	
H7	1.0138	1.1621	0.8362	0.060*	
C8	1.1712 (5)	0.8188 (3)	0.8887 (6)	0.0687 (15)	
H8A	1.2427	0.8250	0.8409	0.103*	
H8B	1.1231	0.7761	0.8600	0.103*	
H8C	1.2127	0.8178	0.9875	0.103*	
C9	0.4449 (5)	0.8729 (3)	0.6389 (5)	0.0559 (13)	
C10	0.3042 (5)	0.8629 (3)	0.5859 (5)	0.0650 (15)	

H10	0.2700	0.8181	0.5687	0.078*
C11	0.2110 (5)	0.9186 (3)	0.5569 (5)	0.0659 (16)
H11	0.1160	0.9107	0.5212	0.079*
C12	0.2593 (5)	0.9832 (3)	0.5809 (5)	0.0621 (15)
H12	0.1971	1.0201	0.5627	0.075*
C13	0.4038 (5)	0.9963 (3)	0.6338 (4)	0.0477 (12)
C14	0.4998 (4)	0.9407 (2)	0.6613 (4)	0.0435 (11)
C15	0.4493 (5)	1.0659 (3)	0.6616 (5)	0.0598 (14)
H15	0.3793	1.0992	0.6435	0.072*
C16	0.5011 (7)	0.7525 (3)	0.6450 (9)	0.120 (3)
H16A	0.4328	0.7411	0.6941	0.180*
H16B	0.5804	0.7225	0.6753	0.180*
H16C	0.4614	0.7465	0.5467	0.180*
Cu1	0.73679 (5)	1.03129 (3)	0.74231 (6)	0.0460 (2)
Na1	0.80631 (19)	0.85912 (9)	0.75096 (19)	0.0555 (5)
O1	1.0749 (3)	0.87466 (17)	0.8560 (3)	0.0542 (8)
O2	0.8957 (3)	0.97361 (15)	0.7830 (3)	0.0468 (8)
O3	0.8430 (3)	1.11575 (16)	0.7817 (3)	0.0538 (8)
O4	0.5441 (3)	0.82251 (18)	0.6728 (4)	0.0732 (11)
O5	0.6356 (3)	0.94771 (15)	0.7049 (3)	0.0482 (8)
O6	0.5707 (3)	1.08808 (17)	0.7071 (3)	0.0589 (9)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
F1'	0.144 (13)	0.108 (12)	0.148 (14)	0.046 (9)	0.031 (9)	0.023 (9)
F2'	0.165 (16)	0.181 (17)	0.184 (17)	-0.004 (10)	0.061 (10)	-0.001 (10)
F3'	0.108 (11)	0.074 (9)	0.043 (7)	0.021 (8)	0.021 (7)	-0.014 (6)
F1	0.160 (6)	0.160 (6)	0.149 (6)	-0.009 (5)	0.106 (5)	-0.052 (5)
F2	0.175 (6)	0.118 (5)	0.116 (4)	0.009 (4)	0.058 (4)	0.059 (4)
F3	0.110 (5)	0.084 (4)	0.151 (6)	-0.006 (3)	-0.051 (4)	0.006 (3)
F4	0.133 (3)	0.070 (3)	0.095 (3)	-0.002 (2)	0.008 (2)	-0.034 (2)
B1	0.083 (4)	0.038 (3)	0.068 (4)	0.008 (3)	0.020 (4)	0.009 (3)
C1	0.044 (2)	0.049 (3)	0.044 (2)	0.003 (2)	0.010 (2)	0.006 (2)
C2	0.047 (3)	0.063 (4)	0.053 (3)	0.006 (2)	0.010 (2)	0.003 (3)
C3	0.041 (2)	0.082 (4)	0.061 (3)	-0.009 (3)	0.014 (2)	-0.006 (3)
C4	0.050 (3)	0.063 (4)	0.052 (3)	-0.018 (3)	0.015 (2)	-0.010 (3)
C5	0.045 (2)	0.046 (3)	0.037 (2)	-0.006 (2)	0.013 (2)	-0.001 (2)
C6	0.043 (2)	0.042 (3)	0.035 (2)	-0.007 (2)	0.0077 (19)	-0.0005 (19)
C7	0.063 (3)	0.039 (3)	0.048 (3)	-0.008 (2)	0.012 (2)	-0.005 (2)
C8	0.073 (3)	0.052 (3)	0.079 (4)	0.022 (3)	0.017 (3)	0.008 (3)
C9	0.047 (3)	0.062 (4)	0.055 (3)	-0.011 (2)	0.007 (2)	-0.006 (3)
C10	0.055 (3)	0.074 (4)	0.069 (3)	-0.018 (3)	0.021 (3)	-0.012 (3)
C11	0.042 (3)	0.095 (5)	0.060 (3)	-0.014 (3)	0.012 (2)	0.000 (3)
C12	0.041 (3)	0.089 (5)	0.054 (3)	0.007 (3)	0.008 (2)	0.010 (3)
C13	0.046 (2)	0.055 (3)	0.040 (2)	-0.003 (2)	0.009 (2)	-0.001 (2)
C14	0.043 (2)	0.050 (3)	0.036 (2)	-0.002 (2)	0.007 (2)	-0.005 (2)
C15	0.049 (3)	0.064 (4)	0.062 (3)	0.014 (3)	0.007 (2)	0.002 (3)

C16	0.079 (4)	0.054 (4)	0.212 (8)	-0.016 (3)	0.010 (5)	-0.023 (5)
Cu1	0.0427 (3)	0.0384 (3)	0.0514 (3)	-0.0002 (3)	0.0023 (2)	-0.0014 (3)
Na1	0.0595 (10)	0.0392 (11)	0.0616 (11)	-0.0008 (9)	0.0046 (9)	0.0018 (9)
O1	0.0483 (17)	0.041 (2)	0.068 (2)	0.0085 (15)	0.0063 (15)	0.0036 (16)
O2	0.0418 (15)	0.0359 (17)	0.0562 (18)	-0.0042 (14)	0.0011 (14)	-0.0001 (15)
O3	0.0541 (19)	0.042 (2)	0.058 (2)	-0.0009 (15)	0.0031 (16)	0.0009 (15)
O4	0.055 (2)	0.045 (2)	0.110 (3)	-0.0091 (17)	0.0063 (19)	-0.014 (2)
O5	0.0379 (15)	0.0449 (19)	0.0561 (19)	-0.0041 (13)	0.0019 (14)	-0.0034 (15)
O6	0.0485 (18)	0.049 (2)	0.074 (2)	0.0037 (16)	0.0064 (17)	-0.0013 (18)

Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )

F1'—B1	1.18 (3)	C10—C11	1.400 (7)
F2'—B1	1.35 (3)	C10—H10	0.9300
F3'—B1	1.270 (16)	C11—C12	1.339 (8)
F1—B1	1.331 (9)	C11—H11	0.9300
F2—B1	1.357 (9)	C12—C13	1.416 (6)
F3—B1	1.387 (8)	C12—H12	0.9300
F4—B1	1.346 (7)	C13—C14	1.415 (6)
C1—O1	1.370 (5)	C13—C15	1.425 (7)
C1—C2	1.376 (6)	C14—O5	1.311 (5)
C1—C6	1.422 (6)	C15—O6	1.248 (5)
C2—C3	1.397 (7)	C15—H15	0.9300
C2—H2	0.9300	C16—O4	1.426 (6)
C3—C4	1.354 (7)	C16—H16A	0.9600
C3—H3	0.9300	C16—H16B	0.9600
C4—C5	1.423 (6)	C16—H16C	0.9600
C4—H4	0.9300	Cu1—O2	1.888 (3)
C5—C7	1.407 (6)	Cu1—O3	1.929 (3)
C5—C6	1.418 (6)	Cu1—O5	1.889 (3)
C6—O2	1.306 (5)	Cu1—O6	1.936 (3)
C7—O3	1.263 (5)	Cu1—Na1	3.399 (2)
C7—H7	0.9300	Na1—O1	2.615 (3)
C8—O1	1.423 (5)	Na1—O2	2.377 (3)
C8—H8A	0.9600	Na1—O4	2.614 (4)
C8—H8B	0.9600	Na1—O5	2.370 (3)
C8—H8C	0.9600	Na1—F1	2.215 (7)
C9—O4	1.364 (6)	Na1—F3'	2.344 (17)
C9—C10	1.373 (6)	Na1—F4 <sup>i</sup>	2.243 (4)
C9—C14	1.415 (6)		
B1—F3'—Na1	127.8 (12)	O6—C15—H15	115.7
B1—F1—Na1	133.9 (6)	C13—C15—H15	115.7
B1—F4—Na1 <sup>ii</sup>	160.3 (4)	O4—C16—H16A	109.5
F1'—B1—F3'	101 (2)	O4—C16—H16B	109.5
F1'—B1—F2'	92 (2)	H16A—C16—H16B	109.5
F3'—B1—F2'	121.7 (19)	O4—C16—H16C	109.5
F1'—B1—F4	132.8 (15)	H16A—C16—H16C	109.5

F3'—B1—F4	109.5 (10)	H16B—C16—H16C	109.5
F4—B1—F2'	100.8 (15)	O2—Cu1—O5	84.72 (13)
F1—B1—F2	107.9 (7)	O2—Cu1—O3	94.30 (13)
F4—B1—F2	109.6 (6)	O5—Cu1—O3	179.02 (12)
F1—B1—F3	109.8 (7)	O2—Cu1—O6	177.29 (14)
F4—B1—F3	108.3 (5)	O5—Cu1—O6	93.78 (13)
F2—B1—F3	108.9 (6)	O3—Cu1—O6	87.19 (13)
F1—B1—F4	112.4 (7)	O2—Cu1—Na1	42.47 (9)
O1—C1—C2	125.9 (4)	O5—Cu1—Na1	42.27 (9)
O1—C1—C6	113.4 (4)	O3—Cu1—Na1	136.76 (10)
C2—C1—C6	120.7 (5)	O6—Cu1—Na1	135.94 (11)
C1—C2—C3	120.5 (5)	F1—Na1—F4 <sup>i</sup>	105.5 (3)
C1—C2—H2	119.7	F1—Na1—F3'	27.6 (4)
C3—C2—H2	119.7	F4 <sup>i</sup> —Na1—F3'	88.1 (4)
C4—C3—C2	120.7 (4)	F1—Na1—O5	104.2 (3)
C4—C3—H3	119.6	F4 <sup>i</sup> —Na1—O5	126.93 (16)
C2—C3—H3	119.6	F3'—Na1—O5	131.4 (5)
C3—C4—C5	120.4 (5)	F1—Na1—O2	120.2 (3)
C3—C4—H4	119.8	F4 <sup>i</sup> —Na1—O2	128.62 (14)
C5—C4—H4	119.8	F3'—Na1—O2	122.4 (5)
C7—C5—C6	122.5 (4)	O5—Na1—O2	64.85 (11)
C7—C5—C4	117.8 (4)	F1—Na1—O4	74.3 (2)
C6—C5—C4	119.7 (4)	F4 <sup>i</sup> —Na1—O4	85.33 (15)
O2—C6—C5	124.0 (4)	F3'—Na1—O4	93.7 (6)
O2—C6—C1	118.2 (4)	O5—Na1—O4	62.07 (11)
C5—C6—C1	117.8 (4)	O2—Na1—O4	126.91 (13)
O3—C7—C5	127.8 (4)	F1—Na1—O1	106.6 (2)
O3—C7—H7	116.1	F4 <sup>i</sup> —Na1—O1	84.28 (14)
C5—C7—H7	116.1	F3'—Na1—O1	83.9 (6)
O1—C8—H8A	109.5	O5—Na1—O1	126.69 (12)
O1—C8—H8B	109.5	O2—Na1—O1	62.17 (10)
H8A—C8—H8B	109.5	O4—Na1—O1	169.41 (13)
O1—C8—H8C	109.5	F1—Na1—Cu1	116.8 (3)
H8A—C8—H8C	109.5	F4 <sup>i</sup> —Na1—Cu1	135.96 (13)
H8B—C8—H8C	109.5	F3'—Na1—Cu1	135.7 (4)
O4—C9—C10	126.1 (5)	O5—Na1—Cu1	32.42 (7)
O4—C9—C14	113.6 (4)	O2—Na1—Cu1	32.44 (7)
C10—C9—C14	120.2 (5)	O4—Na1—Cu1	94.48 (10)
C9—C10—C11	121.4 (5)	O1—Na1—Cu1	94.40 (9)
C9—C10—H10	119.3	C1—O1—C8	117.4 (4)
C11—C10—H10	119.3	C1—O1—Na1	118.6 (2)
C12—C11—C10	119.7 (5)	C8—O1—Na1	124.0 (3)
C12—C11—H11	120.1	C6—O2—Cu1	126.6 (3)
C10—C11—H11	120.1	C6—O2—Na1	127.5 (3)
C11—C12—C13	120.9 (5)	Cu1—O2—Na1	105.09 (12)
C11—C12—H12	119.5	C7—O3—Cu1	124.2 (3)
C13—C12—H12	119.5	C9—O4—C16	118.2 (4)
C14—C13—C12	120.1 (5)	C9—O4—Na1	118.6 (3)

---

C14—C13—C15	121.2 (4)	C16—O4—Na1	122.6 (3)
C12—C13—C15	118.7 (5)	C14—O5—Cu1	127.0 (3)
O5—C14—C9	117.9 (4)	C14—O5—Na1	127.7 (3)
O5—C14—C13	124.6 (4)	Cu1—O5—Na1	105.31 (13)
C9—C14—C13	117.5 (4)	C15—O6—Cu1	124.6 (3)
O6—C15—C13	128.5 (5)		

---

Symmetry codes: (i)  $x, -y+3/2, z+1/2$ ; (ii)  $x, -y+3/2, z-1/2$ .