

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tetraaquabis[4-(4*H*-1,2,4-triazol-4-yl)-benzoato- κ N¹]nickel(II) decahydrateWeixuan Sun,^a Yaqin Yu,^a Guanjun Wang^{b*} and Xiaohui Wu^c

^aDepartment of Preventive Medicine, School of Public Health, Jilin University, Changchun 130021, People's Republic of China, ^bDepartment of Hematology, the First Hospital of Jilin University, Changchun 130021, People's Republic of China, and ^cSchool of Public Health, Jilin University, Changchun 130021, People's Republic of China

Correspondence e-mail: doctorwanggj@163.com


Received 19 November 2011; accepted 28 November 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.049; wR factor = 0.119; data-to-parameter ratio = 12.9.

In the title compound, $[\text{Ni}(\text{C}_9\text{H}_6\text{N}_3\text{O}_2)_2(\text{H}_2\text{O})_4] \cdot 10\text{H}_2\text{O}$, the Ni^{II} ion lies on a twofold rotation axis and displays a slightly distorted octahedral geometry defined by two N atoms from two monodentate 4-(1,2,4-triazol-4-yl)benzoate ligands and four water molecules, two of which also lie on the twofold rotation axis. In the crystal, the complex molecules and uncoordinated water molecules are linked *via* intermolecular O—H...N and O—H...O hydrogen bonds, forming a three-dimensional supramolecular network. π - π interactions between the benzene rings provide additional stability of the crystal packing [centroid-centroid distance = 3.792 (2) Å].

Related literature

For general background to the applications and structures of metal coordination polymers, see: Rowsell & Yaghi (2005); Su *et al.* (2010); Wang *et al.* (2009); Zhang & Chen (2008). For a related structure, see: Cui & Zhao (2011).



Experimental

Crystal data

$[\text{Ni}(\text{C}_9\text{H}_6\text{N}_3\text{O}_2)_2(\text{H}_2\text{O})_4] \cdot 10\text{H}_2\text{O}$
 $M_r = 687.25$
 Monoclinic, $C2/c$

$a = 25.840$ (3) Å
 $b = 7.8664$ (8) Å
 $c = 16.8013$ (17) Å

$\beta = 112.712$ (1)°
 $V = 3150.3$ (6) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.70$ mm⁻¹
 $T = 293$ K
 $0.22 \times 0.20 \times 0.19$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2001)
 $T_{\min} = 0.83$, $T_{\max} = 0.90$

8290 measured reflections
 3079 independent reflections
 2273 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.057$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.119$
 $S = 1.04$
 3079 reflections
 238 parameters
 14 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.55$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.63$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3—H3A...O1 ⁱ	0.84 (2)	1.93 (2)	2.751 (3)	167 (4)
O4—H4A...O1 ⁱⁱ	0.84 (2)	1.86 (2)	2.692 (3)	172 (4)
O5—H5A...O8	0.82 (2)	1.94 (2)	2.752 (3)	169 (4)
O5—H5B...O7	0.83 (2)	1.84 (2)	2.670 (3)	171 (4)
O6—H6A...O7	0.82 (2)	1.95 (2)	2.773 (4)	178 (4)
O6—H6B...O10	0.84 (2)	1.91 (3)	2.747 (4)	177 (6)
O7—H7A...O2 ⁱⁱⁱ	0.84 (2)	1.84 (2)	2.674 (3)	170 (4)
O7—H7B...O9 ^{iv}	0.84 (2)	1.88 (2)	2.715 (4)	171 (4)
O8—H8A...N3 ^v	0.82 (2)	2.16 (2)	2.943 (3)	160 (4)
O8—H8B...O2 ^{vi}	0.85 (2)	1.92 (2)	2.763 (3)	175 (4)
O9—H9A...O1 ⁱⁱ	0.84 (2)	1.93 (2)	2.751 (3)	169 (4)
O9—H9B...O8	0.85 (2)	1.93 (2)	2.757 (3)	164 (4)

Symmetry codes: (i) $-x + \frac{3}{2}, -y + \frac{3}{2}, -z + 2$; (ii) $-x + \frac{3}{2}, -y + \frac{1}{2}, -z + 2$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $x, y + 1, z$; (v) $x, -y + 1, z - \frac{1}{2}$; (vi) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors thank Jilin University for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2491).

References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cui, F.-H. & Zhao, P.-Z. (2011). *Acta Cryst.* **E67**, m1341–m1342.
 Rowsell, J. L. C. & Yaghi, O. M. (2005). *Angew. Chem. Int. Ed.* **44**, 4670–4679.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Su, S.-Q., Guo, Z.-Y., Li, G.-H., Deng, R.-P., Song, S.-Y., Qin, C., Pan, C.-L., Guo, H.-D., Gao, F., Wang, S. & Zhang, H.-J. (2010). *Dalton Trans.* **39**, 9123–9130.
 Wang, G.-H., Li, Z.-G., Jia, H.-Q., Hu, N.-H. & Xu, J.-W. (2009). *Acta Cryst.* **E65**, m1568–m1569.
 Zhang, J.-P. & Chen, X.-M. (2008). *J. Am. Chem. Soc.* **130**, 6010–6017.

supporting information

Acta Cryst. (2012). E68, m10 [doi:10.1107/S1600536811051063]

Tetraaquabis[4-(4*H*-1,2,4-triazol-4-yl)benzoato- κ N¹]nickel(II) decahydrate**Weixuan Sun, Yaqin Yu, Guanjun Wang and Xiaohui Wu****S1. Comment**

Recently, the chemists have devoted themselves to design and synthesize coordination polymers, not only due to their potential applications in the realm of gas adsorption and separation, catalysis, magnetism, luminescence, host–guest chemistry *etc.*, but also for their aesthetic and often complicated architectures and topologies (Su *et al.*, 2010; Wang *et al.*, 2009). It is well known that carboxylic acids are excellent building blocks for the construction of coordination polymers because the carboxylate groups may induce core aggregation and link these discrete clusters into an extended framework by virtue of its bridging ability (Rowse & Yaghi, 2005; Zhang & Chen, 2008). Taking these into account, we chose a carboxylate ligand, 4-(1,2,4-triazol-4-yl)benzoic acid, generating the title compound, which is reported here.

In the title compound, the Ni^{II} ion lies on a twofold rotation axis and displays a slightly distorted octahedral geometry defined by two N atoms from two 4-(1,2,4-triazol-4-yl)benzoate ligands and four water molecules, two of which lie on the twofold rotation axis (Fig. 1). The bond lengths and angles are in a normal range (Cui & Zhao, 2011). In the crystal, the complex molecules and uncoordinated water molecules are linked *via* intermolecular O—H \cdots N and O—H \cdots O hydrogen bonds, forming a three-dimensional supramolecular network (Fig. 2). π – π interactions between the benzene rings, with a centroid–centroid distance of 3.792 (2) Å, provide additional stability of the crystal packing.

S2. Experimental

The synthesis was performed under hydrothermal conditions. A mixture of Ni(CH₃COO)₂·4H₂O (0.2 mmol, 0.050 g), 4-(1,2,4-triazol-4-yl)benzoic acid (0.4 mmol, 0.075 g), NaOH (0.4 mmol, 0.016 g) and H₂O (15 ml) in a 25 ml stainless steel reactor with a Teflon liner was heated from 293 to 443 K in 2 h and a constant temperature was maintained at 443 K for 72 h. After the mixture was cooled to 298 K, pink crystals of the title compound were obtained from the reaction.

S3. Refinement

H atoms on C atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atoms bonded to O atoms were located in a difference Fourier map and refined with O—H distance restraints of 0.85 (2) Å and with $U_{\text{iso}}(\text{H}) = 0.054 \text{ \AA}^2$.




Figure 1

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity. [Symmetry code: (i) 1-x, y, 3/2-z.]





Figure 2

View of the three-dimensional structure of the title compound built by hydrogen bonds (dashed lines).

Tetraaquabis[4-(4*H*-1,2,4-triazol-4-yl)benzoato- κ N¹]nickel(II) decahydrate

Crystal data



$M_r = 687.25$

Monoclinic, $C2/c$

Hall symbol: -C 2yc

$a = 25.840$ (3) Å

$b = 7.8664$ (8) Å

$c = 16.8013$ (17) Å

$\beta = 112.712$ (1)°

$V = 3150.3$ (6) Å³

$Z = 4$

$F(000) = 1448$

$D_x = 1.449$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3079 reflections

$\theta = 1.0$ – 25.9 °

$\mu = 0.70$ mm⁻¹

$T = 293$ K

Block, pink

$0.22 \times 0.20 \times 0.19$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2001)
 $T_{\min} = 0.83$, $T_{\max} = 0.90$

8290 measured reflections
3079 independent reflections
2273 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.057$
 $\theta_{\max} = 25.9^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -31 \rightarrow 31$
 $k = -6 \rightarrow 9$
 $l = -20 \rightarrow 17$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.119$
 $S = 1.04$
3079 reflections
238 parameters
14 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0451P)^2 + 2.4003P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.55 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.63 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	0.59822 (8)	0.7500	0.01918 (19)
C1	0.83480 (12)	0.4306 (4)	1.04342 (19)	0.0186 (7)
C2	0.79863 (12)	0.3835 (4)	0.96082 (19)	0.0200 (7)
H2	0.8126	0.3236	0.9257	0.024*
C3	0.74225 (13)	0.4242 (4)	0.93015 (19)	0.0205 (7)
H3	0.7182	0.3890	0.8755	0.025*
C4	0.72201 (12)	0.5180 (4)	0.98169 (19)	0.0163 (7)
C5	0.75769 (13)	0.5702 (4)	1.0635 (2)	0.0209 (7)
H5	0.7441	0.6351	1.0975	0.025*
C6	0.81360 (12)	0.5246 (4)	1.0939 (2)	0.0194 (7)
H6	0.8374	0.5575	1.1491	0.023*
C7	0.89538 (13)	0.3748 (4)	1.0778 (2)	0.0196 (7)
C8	0.62534 (12)	0.5537 (4)	0.86818 (19)	0.0192 (7)
H8	0.6334	0.5222	0.8209	0.023*
C9	0.63313 (13)	0.6112 (4)	0.9971 (2)	0.0231 (7)
H9	0.6480	0.6275	1.0565	0.028*
N1	0.66358 (10)	0.5587 (3)	0.95053 (15)	0.0170 (6)
N2	0.57580 (10)	0.5986 (3)	0.86385 (15)	0.0193 (6)
N3	0.58082 (11)	0.6353 (4)	0.94733 (16)	0.0230 (7)
O1	0.92384 (9)	0.3974 (3)	1.15821 (13)	0.0218 (5)
O2	0.91424 (9)	0.3070 (3)	1.02773 (14)	0.0302 (6)
O3	0.5000	0.8594 (4)	0.7500	0.0294 (8)
O4	0.5000	0.3349 (4)	0.7500	0.0228 (7)
O5	0.54970 (9)	0.5800 (3)	0.67831 (14)	0.0211 (5)

O6	0.70477 (12)	0.6665 (4)	0.74921 (19)	0.0430 (7)
O7	0.60752 (10)	0.8271 (3)	0.64103 (15)	0.0280 (6)
O8	0.52147 (10)	0.3127 (3)	0.56372 (15)	0.0255 (6)
O9	0.60244 (10)	0.1461 (3)	0.69923 (16)	0.0315 (6)
O10	0.69525 (11)	0.3274 (4)	0.70686 (17)	0.0363 (7)
H3A	0.5254 (13)	0.921 (4)	0.784 (2)	0.054*
H4A	0.5231 (14)	0.265 (4)	0.783 (2)	0.054*
H5A	0.5411 (17)	0.510 (4)	0.639 (2)	0.054*
H5B	0.5653 (16)	0.656 (4)	0.661 (3)	0.054*
H6A	0.6758 (12)	0.715 (5)	0.718 (2)	0.054*
H6B	0.7026 (18)	0.562 (3)	0.738 (3)	0.054*
H7A	0.6022 (17)	0.833 (5)	0.5885 (14)	0.054*
H7B	0.6041 (18)	0.929 (3)	0.654 (3)	0.054*
H8A	0.5304 (17)	0.319 (6)	0.5218 (19)	0.054*
H8B	0.4879 (10)	0.279 (5)	0.550 (3)	0.054*
H9A	0.5962 (17)	0.146 (6)	0.7445 (18)	0.054*
H9B	0.5732 (12)	0.191 (5)	0.662 (2)	0.054*
H10A	0.7237 (12)	0.268 (5)	0.727 (2)	0.054*
H10B	0.6695 (14)	0.275 (5)	0.712 (3)	0.054*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0147 (3)	0.0216 (3)	0.0196 (3)	0.000	0.0049 (2)	0.000
C1	0.0117 (15)	0.0238 (18)	0.0189 (16)	0.0005 (13)	0.0044 (13)	0.0017 (14)
C2	0.0179 (17)	0.0281 (19)	0.0144 (16)	0.0013 (14)	0.0069 (13)	-0.0005 (14)
C3	0.0155 (16)	0.0305 (19)	0.0121 (15)	-0.0005 (14)	0.0014 (12)	0.0006 (14)
C4	0.0105 (15)	0.0197 (16)	0.0172 (16)	0.0021 (13)	0.0035 (13)	0.0023 (14)
C5	0.0188 (17)	0.0258 (19)	0.0174 (16)	0.0041 (14)	0.0062 (13)	-0.0042 (14)
C6	0.0153 (16)	0.0223 (17)	0.0170 (16)	-0.0007 (13)	0.0024 (13)	-0.0029 (14)
C7	0.0145 (16)	0.0217 (18)	0.0214 (17)	0.0004 (13)	0.0057 (14)	0.0031 (15)
C8	0.0144 (16)	0.0263 (18)	0.0148 (15)	0.0000 (13)	0.0034 (13)	-0.0012 (14)
C9	0.0170 (17)	0.037 (2)	0.0146 (16)	0.0034 (15)	0.0055 (13)	-0.0029 (15)
N1	0.0123 (13)	0.0229 (15)	0.0136 (13)	0.0014 (11)	0.0026 (10)	-0.0014 (11)
N2	0.0151 (13)	0.0272 (15)	0.0155 (13)	-0.0008 (12)	0.0058 (11)	-0.0021 (12)
N3	0.0144 (14)	0.0370 (17)	0.0149 (14)	0.0017 (12)	0.0026 (11)	-0.0043 (13)
O1	0.0167 (11)	0.0254 (12)	0.0180 (11)	0.0023 (10)	0.0010 (9)	0.0010 (10)
O2	0.0169 (13)	0.0498 (17)	0.0221 (13)	0.0123 (12)	0.0056 (10)	0.0020 (12)
O3	0.0210 (19)	0.0179 (18)	0.032 (2)	0.000	-0.0087 (15)	0.000
O4	0.0194 (18)	0.0150 (17)	0.0248 (19)	0.000	-0.0015 (14)	0.000
O5	0.0195 (12)	0.0248 (13)	0.0213 (12)	-0.0055 (10)	0.0106 (10)	-0.0034 (10)
O6	0.0341 (17)	0.0381 (17)	0.0489 (18)	0.0012 (14)	0.0073 (14)	0.0007 (15)
O7	0.0318 (14)	0.0289 (14)	0.0249 (13)	-0.0017 (12)	0.0126 (12)	0.0015 (12)
O8	0.0190 (13)	0.0378 (15)	0.0209 (12)	-0.0077 (11)	0.0090 (10)	-0.0023 (11)
O9	0.0255 (14)	0.0388 (15)	0.0315 (15)	0.0064 (12)	0.0123 (12)	0.0047 (13)
O10	0.0282 (16)	0.0390 (17)	0.0373 (16)	-0.0016 (12)	0.0080 (13)	0.0061 (13)

Geometric parameters (Å, °)

Ni1—O3	2.054 (3)	C8—H8	0.9300
Ni1—O4	2.071 (3)	C9—N3	1.300 (4)
Ni1—O5	2.077 (2)	C9—N1	1.370 (4)
Ni1—N2	2.145 (2)	C9—H9	0.9300
C1—C6	1.387 (4)	N2—N3	1.388 (3)
C1—C2	1.391 (4)	O3—H3A	0.836 (18)
C1—C7	1.510 (4)	O4—H4A	0.840 (18)
C2—C3	1.382 (4)	O5—H5A	0.822 (19)
C2—H2	0.9300	O5—H5B	0.835 (19)
C3—C4	1.385 (4)	O6—H6A	0.824 (19)
C3—H3	0.9300	O6—H6B	0.838 (19)
C4—C5	1.389 (4)	O7—H7A	0.840 (18)
C4—N1	1.430 (4)	O7—H7B	0.844 (19)
C5—C6	1.381 (4)	O8—H8A	0.823 (18)
C5—H5	0.9300	O8—H8B	0.848 (19)
C6—H6	0.9300	O9—H9A	0.836 (19)
C7—O2	1.243 (4)	O9—H9B	0.847 (19)
C7—O1	1.277 (4)	O10—H10A	0.824 (19)
C8—N2	1.303 (4)	O10—H10B	0.817 (19)
C8—N1	1.355 (4)		
O3—Ni1—O4	180.000 (2)	C5—C6—C1	121.1 (3)
O3—Ni1—O5 ⁱ	93.97 (6)	C5—C6—H6	119.5
O4—Ni1—O5 ⁱ	86.03 (7)	C1—C6—H6	119.5
O3—Ni1—O5	93.97 (7)	O2—C7—O1	124.0 (3)
O4—Ni1—O5	86.03 (7)	O2—C7—C1	119.0 (3)
O5 ⁱ —Ni1—O5	172.07 (13)	O1—C7—C1	116.9 (3)
O3—Ni1—N2	89.93 (7)	N2—C8—N1	111.3 (3)
O4—Ni1—N2	90.07 (7)	N2—C8—H8	124.4
O5 ⁱ —Ni1—N2	92.24 (9)	N1—C8—H8	124.4
O5—Ni1—N2	87.77 (9)	N3—C9—N1	111.2 (3)
O3—Ni1—N2 ⁱ	89.93 (7)	N3—C9—H9	124.4
O4—Ni1—N2 ⁱ	90.07 (7)	N1—C9—H9	124.4
O5 ⁱ —Ni1—N2 ⁱ	87.77 (9)	C8—N1—C9	103.8 (3)
O5—Ni1—N2 ⁱ	92.24 (9)	C8—N1—C4	128.2 (3)
N2—Ni1—N2 ⁱ	179.85 (16)	C9—N1—C4	128.1 (2)
C6—C1—C2	118.7 (3)	C8—N2—N3	107.1 (2)
C6—C1—C7	121.1 (3)	C8—N2—Ni1	126.1 (2)
C2—C1—C7	120.1 (3)	N3—N2—Ni1	126.73 (18)
C3—C2—C1	121.0 (3)	C9—N3—N2	106.7 (2)
C3—C2—H2	119.5	Ni1—O3—H3A	125 (3)
C1—C2—H2	119.5	Ni1—O4—H4A	131 (3)
C2—C3—C4	119.3 (3)	Ni1—O5—H5A	118 (3)
C2—C3—H3	120.4	Ni1—O5—H5B	130 (3)
C4—C3—H3	120.4	H5A—O5—H5B	103 (4)
C3—C4—C5	120.6 (3)	H6A—O6—H6B	110 (4)

C3—C4—N1	119.4 (3)	H7A—O7—H7B	103 (4)
C5—C4—N1	120.0 (3)	H8A—O8—H8B	112 (4)
C6—C5—C4	119.3 (3)	H9A—O9—H9B	103 (4)
C6—C5—H5	120.4	H10A—O10—H10B	108 (4)
C4—C5—H5	120.4		
C6—C1—C2—C3	-2.1 (5)	C3—C4—N1—C8	-17.2 (5)
C7—C1—C2—C3	175.9 (3)	C5—C4—N1—C8	163.8 (3)
C1—C2—C3—C4	2.0 (5)	C3—C4—N1—C9	162.5 (3)
C2—C3—C4—C5	-0.3 (5)	C5—C4—N1—C9	-16.5 (5)
C2—C3—C4—N1	-179.3 (3)	N1—C8—N2—N3	-0.1 (4)
C3—C4—C5—C6	-1.3 (5)	N1—C8—N2—Ni1	-176.2 (2)
N1—C4—C5—C6	177.7 (3)	O3—Ni1—N2—C8	-109.7 (3)
C4—C5—C6—C1	1.2 (5)	O4—Ni1—N2—C8	70.3 (3)
C2—C1—C6—C5	0.4 (5)	O5 ⁱ —Ni1—N2—C8	156.4 (3)
C7—C1—C6—C5	-177.5 (3)	O5—Ni1—N2—C8	-15.7 (3)
C6—C1—C7—O2	-172.3 (3)	O3—Ni1—N2—N3	75.0 (2)
C2—C1—C7—O2	9.8 (5)	O4—Ni1—N2—N3	-105.0 (2)
C6—C1—C7—O1	9.3 (5)	O5 ⁱ —Ni1—N2—N3	-19.0 (3)
C2—C1—C7—O1	-168.6 (3)	O5—Ni1—N2—N3	169.0 (3)
N2—C8—N1—C9	-0.1 (4)	N1—C9—N3—N2	-0.4 (4)
N2—C8—N1—C4	179.7 (3)	C8—N2—N3—C9	0.3 (4)
N3—C9—N1—C8	0.3 (4)	Ni1—N2—N3—C9	176.4 (2)
N3—C9—N1—C4	-179.5 (3)		

Symmetry code: (i) $-x+1, y, -z+3/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H3A \cdots O1 ⁱⁱ	0.84 (2)	1.93 (2)	2.751 (3)	167 (4)
O4—H4A \cdots O1 ⁱⁱⁱ	0.84 (2)	1.86 (2)	2.692 (3)	172 (4)
O5—H5A \cdots O8	0.82 (2)	1.94 (2)	2.752 (3)	169 (4)
O5—H5B \cdots O7	0.83 (2)	1.84 (2)	2.670 (3)	171 (4)
O6—H6A \cdots O7	0.82 (2)	1.95 (2)	2.773 (4)	178 (4)
O6—H6B \cdots O10	0.84 (2)	1.91 (3)	2.747 (4)	177 (6)
O7—H7A \cdots O2 ^{iv}	0.84 (2)	1.84 (2)	2.674 (3)	170 (4)
O7—H7B \cdots O9 ^v	0.84 (2)	1.88 (2)	2.715 (4)	171 (4)
O8—H8A \cdots N3 ^{vi}	0.82 (2)	2.16 (2)	2.943 (3)	160 (4)
O8—H8B \cdots O2 ^{vii}	0.85 (2)	1.92 (2)	2.763 (3)	175 (4)
O9—H9A \cdots O1 ⁱⁱⁱ	0.84 (2)	1.93 (2)	2.751 (3)	169 (4)
O9—H9B \cdots O8	0.85 (2)	1.93 (2)	2.757 (3)	164 (4)

Symmetry codes: (ii) $-x+3/2, -y+3/2, -z+2$; (iii) $-x+3/2, -y+1/2, -z+2$; (iv) $-x+3/2, y+1/2, -z+3/2$; (v) $x, y+1, z$; (vi) $x, -y+1, z-1/2$; (vii) $x-1/2, -y+1/2, z-1/2$.