

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Ethoxy-5-methylbis[1,2,4]triazolo-[1,5-a:4',3'-c]quinazoline

 Rashad Al-Salahi,<sup>a\*</sup> Detlef Geffken<sup>b</sup> and Ahmed Bari<sup>a</sup>

<sup>a</sup>Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, and <sup>b</sup>Department of Chemistry, Institute of Pharmacy, University of Hamburg, Bundesstrasse 45, 20146, Hamburg, Germany  
Correspondence e-mail: abari@ksu.edu.sa


Received 1 November 2011; accepted 7 December 2011

Key indicators: single-crystal X-ray study;  $T = 153$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.055;  $wR$  factor = 0.134; data-to-parameter ratio = 15.6.

The title compound,  $\text{C}_{13}\text{H}_{12}\text{N}_6\text{O}$ , is a functionalized ditriazoloquinazoline with substituted ethoxy and methyl groups attached at the 2-position of each triazole spacer. The fused-ring system is essentially planar [r.m.s. deviation = 0.016 (2) Å]. In the crystal, a weak  $\text{C}-\text{H}\cdots\text{N}$  hydrogen bond connects the molecules into a chain along [101].

### Related literature

For the biological activity of quinazoline derivatives, see: Alagarsamy *et al.* (2005, 2007). For our study of the chemical and pharmacological properties of triazolo quinazoline derivatives, see: Al-Salahi (2009); Al-Salahi *et al.* (2010, 2011); Berezank *et al.* (2008a,b). For related structures, see: El-Tombary *et al.* (1999).



### Experimental

#### Crystal data

$\text{C}_{13}\text{H}_{12}\text{N}_6\text{O}$   
 $M_r = 268.29$   
 Monoclinic,  $P2_1/c$   
 $a = 7.4121$  (11) Å  
 $b = 19.240$  (3) Å  
 $c = 9.3096$  (14) Å  
 $\beta = 109.051$  (2)°

$V = 1254.9$  (3) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.10$  mm<sup>-1</sup>  
 $T = 153$  K  
 $0.43 \times 0.21 \times 0.05$  mm

#### Data collection

Bruker SMART APEX CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 1998)  
 $T_{\min} = 0.959$ ,  $T_{\max} = 0.995$   
 2851 measured reflections  
 2851 independent reflections  
 1817 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.053$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$   
 $wR(F^2) = 0.134$   
 $S = 0.91$   
 2851 reflections  
 183 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.29$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3}\cdots\text{N4}^i$	0.95	2.60	3.533 (2)	166

 Symmetry code: (i)  $x - 1, y, z - 1$ .

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: publCIF (Westrip, 2010).

We wish to express our gratitude to the Department of Chemistry, X-ray crystallography division of Hamburg University for their valuable help in the determination of the X-ray crystal structure.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5002).

### References

- Alagarsamy, V., Giridhar, R. & Yadav, M. R. (2005). *Bioorg. Med. Chem. Lett.* **15**, 1877–1880.  
 Alagarsamy, V., Solomon, V. R. & Murugan, M. (2007). *Bioorg. Med. Chem.* **15**, 4009–4015.  
 Al-Salahi, R. (2009). PhD dissertation, Hamburg University, Germany.  
 Al-Salahi, R. & Geffken, D. (2010). *Molecules*, **15**, 7016–7034.  
 Al-Salahi, R., Geffken, D. & Koellner, M. (2011). *Chem. Pharm. Bull.* **59**, 730–733.  
 Berezank, J. F., Chan, D., Geffken, D., Hanagan, M. A., Lepone, G. E., Pasteris, R. S. & Swan, T. (2008a). World patent WO 2008103357.  
 Berezank, J. F., Chan, D., Geffken, D., Hanagan, M. A., Lepone, G. E., Pasteris, R. S. & Swan, T. (2008b). *Chem. Abstr.* 1042828.  
 Bruker (1998). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 El-Tombary, A. A., Ismail, K. A., Aboulwafa, O. M., Omar, A.-M., El-Azzouni, M. Z. & El-Mansoury, S. T. (1999). *Farmaco*, **54**, 486–488.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2012). E68, o101 [doi:10.1107/S1600536811052810]

**2-Ethoxy-5-methylbis[1,2,4]triazolo[1,5-a:4',3'-c]quinazoline****Rashad Al-Salahi, Detlef Geffken and Ahmed Bari****S1. Comment**


Recently, the triazolo-annelated quinazoline compounds were found to display remarkable pharmacological activities (Alagarsamy *et al.*, 2005, 2007). The novel 2-alkoxy-1,2,4-triazolo[1,5-*a*]quinazolines have been proven as excellent agents for controlling the plant growth diseases caused by fungal pathogen agents (Berezank *et al.*, 2008*a,b*), and described as adenosine receptor antagonists (Al-Salahi *et al.*, 2011). Furthermore, the bis-[1,2,4]triazolo[4,3 - *a*:4',3'-*c*]quinazoline has been reported to exhibit antitoxoplasmosis activity (El-Tombary *et al.*, 1999). This work has done in continuation of our program with the aim of obtaining an interesting series of bis-[1,2,4]triazoloquinazolines compounds which could be expected to contribute as a pharmacophore to the bioactivity of their triazoloquinazoline parent compounds (Al-Salahi *et al.*, 2009, 2010).

**S2. Experimental**

A mixture of 2-ethoxy-5-chloro-[1,2,4]triazolo[1,5-*a*]quinazoline (1 mmol) and acetohydrazide (2.2 mmol) was refluxed in absolute toluene (15 ml) in the presence of sodium hydride (0.8 mmol) for 18 h. The solvent was removed under reduced pressure, and the residue was crystallized from methanol to afford (C<sub>13</sub>H<sub>12</sub>N<sub>6</sub>O) as yellowish crystal.

**S3. Refinement**

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95–0.98 Å. The displacement parameters are  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$ , where  $x = 1.2$  or  $1.5$ .

**Figure 1**

The molecular structure of the title compound, showing 50% probability displacement ellipsoids for non-H atoms and the atom-numbering scheme.

### 2-Ethoxy-5-methylbis[1,2,4]triazolo[1,5-a:4',3'-c]quinazoline

#### Crystal data

$C_{13}H_{12}N_6O$

$M_r = 268.29$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 7.4121$  (11) Å

$b = 19.240$  (3) Å

$c = 9.3096$  (14) Å

$\beta = 109.051$  (2)°

$V = 1254.9$  (3) Å<sup>3</sup>

$Z = 4$

$F(000) = 560$

$D_x = 1.420$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 210 reflections

$\theta = 1.2$ – $27.0$ °

$\mu = 0.10$  mm<sup>-1</sup>

$T = 153$  K

Plate, colourless

$0.43 \times 0.21 \times 0.05$  mm

#### Data collection

Bruker SMART APEX CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scan

Absorption correction: multi-scan

(*SADABS*; Bruker, 1998)

$T_{\min} = 0.959$ ,  $T_{\max} = 0.995$

2851 measured reflections

2851 independent reflections

1817 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.053$

$\theta_{\max} = 27.5$ °,  $\theta_{\min} = 2.6$ °

$h = -9 \rightarrow 9$

$k = -24 \rightarrow 24$

$l = 0 \rightarrow 11$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.055$   
 $wR(F^2) = 0.134$   
 $S = 0.91$   
 2851 reflections  
 183 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.072P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.044$   
 $\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.26808 (18)	0.12586 (7)	0.60062 (14)	0.0292 (4)
N5	-0.0311 (2)	0.21218 (8)	0.29706 (16)	0.0243 (4)
N3	0.0101 (2)	0.33173 (8)	0.34526 (16)	0.0244 (4)
N4	0.1812 (2)	0.23979 (8)	0.52010 (16)	0.0233 (4)
N6	0.0364 (2)	0.14887 (8)	0.36339 (16)	0.0265 (4)
N1	-0.0267 (3)	0.44495 (9)	0.32026 (18)	0.0375 (4)
N2	-0.1545 (2)	0.41376 (9)	0.18987 (18)	0.0360 (4)
C6	-0.1751 (3)	0.22080 (10)	0.1548 (2)	0.0263 (5)
C1	-0.2270 (3)	0.28941 (10)	0.10826 (19)	0.0273 (5)
C8	0.0587 (2)	0.26359 (10)	0.39243 (19)	0.0223 (4)
C11	0.1609 (2)	0.16942 (10)	0.49439 (19)	0.0232 (4)
C5	-0.2620 (3)	0.16363 (11)	0.0665 (2)	0.0322 (5)
H5	-0.2268	0.1175	0.1003	0.039*
C7	-0.1306 (3)	0.34608 (11)	0.20684 (19)	0.0276 (5)
C4	-0.4021 (3)	0.17661 (12)	-0.0729 (2)	0.0379 (6)
H4	-0.4620	0.1386	-0.1356	0.045*
C3	-0.4562 (3)	0.24455 (12)	-0.1222 (2)	0.0379 (6)
H3	-0.5523	0.2522	-0.2174	0.045*
C9	0.0698 (3)	0.39608 (10)	0.4119 (2)	0.0292 (5)
C2	-0.3704 (3)	0.30040 (12)	-0.0332 (2)	0.0349 (5)
H2	-0.4079	0.3464	-0.0671	0.042*
C10	0.2187 (3)	0.40648 (10)	0.5633 (2)	0.0336 (5)
H10A	0.2326	0.4562	0.5868	0.050*
H10B	0.3407	0.3877	0.5610	0.050*
H10C	0.1805	0.3823	0.6414	0.050*

C12	0.2394 (3)	0.05207 (10)	0.5604 (2)	0.0371 (5)
H12A	0.2652	0.0432	0.4642	0.045*
H12B	0.1059	0.0386	0.5465	0.045*
C13	0.3737 (3)	0.01071 (11)	0.6867 (2)	0.0439 (6)
H13A	0.5055	0.0238	0.6983	0.066*
H13B	0.3556	-0.0389	0.6628	0.066*
H13C	0.3479	0.0202	0.7816	0.066*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0254 (7)	0.0221 (8)	0.0340 (7)	0.0003 (6)	0.0013 (6)	-0.0025 (6)
N5	0.0207 (8)	0.0270 (10)	0.0242 (8)	-0.0019 (7)	0.0060 (6)	-0.0033 (7)
N3	0.0219 (8)	0.0257 (10)	0.0265 (8)	0.0020 (7)	0.0092 (7)	0.0015 (7)
N4	0.0174 (8)	0.0253 (10)	0.0263 (8)	-0.0008 (7)	0.0059 (7)	-0.0020 (7)
N6	0.0241 (8)	0.0264 (10)	0.0272 (8)	-0.0008 (7)	0.0061 (7)	-0.0031 (7)
N1	0.0416 (11)	0.0319 (11)	0.0399 (10)	0.0074 (9)	0.0147 (8)	0.0028 (8)
N2	0.0360 (10)	0.0366 (12)	0.0348 (9)	0.0086 (9)	0.0108 (8)	0.0047 (8)
C6	0.0185 (9)	0.0407 (13)	0.0216 (9)	-0.0026 (9)	0.0091 (7)	-0.0014 (9)
C1	0.0190 (9)	0.0404 (13)	0.0248 (9)	0.0005 (9)	0.0100 (8)	0.0021 (8)
C8	0.0169 (9)	0.0267 (11)	0.0249 (9)	-0.0012 (8)	0.0092 (8)	-0.0029 (8)
C11	0.0179 (9)	0.0252 (11)	0.0264 (9)	-0.0011 (8)	0.0073 (7)	-0.0038 (8)
C5	0.0256 (10)	0.0441 (14)	0.0282 (10)	-0.0102 (10)	0.0107 (8)	-0.0060 (9)
C7	0.0226 (10)	0.0359 (13)	0.0258 (9)	0.0068 (9)	0.0099 (8)	0.0062 (9)
C4	0.0254 (11)	0.0629 (17)	0.0263 (10)	-0.0151 (11)	0.0097 (9)	-0.0098 (10)
C3	0.0200 (10)	0.0670 (17)	0.0250 (10)	-0.0045 (11)	0.0050 (8)	0.0011 (10)
C9	0.0296 (11)	0.0266 (12)	0.0350 (10)	0.0017 (9)	0.0153 (9)	-0.0033 (9)
C2	0.0220 (10)	0.0553 (15)	0.0281 (10)	0.0034 (10)	0.0092 (9)	0.0076 (10)
C10	0.0338 (12)	0.0270 (12)	0.0394 (11)	-0.0031 (10)	0.0110 (9)	-0.0077 (9)
C12	0.0357 (12)	0.0240 (12)	0.0473 (12)	-0.0033 (10)	0.0076 (10)	-0.0056 (10)
C13	0.0462 (14)	0.0250 (13)	0.0560 (14)	0.0026 (10)	0.0106 (11)	0.0014 (10)

*Geometric parameters (Å, °)*

O1—C11	1.341 (2)	C5—C4	1.395 (3)
O1—C12	1.466 (2)	C5—H5	0.9500
N5—C8	1.351 (2)	C4—C3	1.400 (3)
N5—N6	1.383 (2)	C4—H4	0.9500
N5—C6	1.414 (2)	C3—C2	1.379 (3)
N3—C9	1.391 (2)	C3—H3	0.9500
N3—C8	1.392 (2)	C9—C10	1.493 (3)
N3—C7	1.395 (2)	C2—H2	0.9500
N4—C8	1.320 (2)	C10—H10A	0.9800
N4—C11	1.375 (2)	C10—H10B	0.9800
N6—C11	1.327 (2)	C10—H10C	0.9800
N1—C9	1.313 (2)	C12—C13	1.497 (3)
N1—N2	1.407 (2)	C12—H12A	0.9900
N2—C7	1.317 (3)	C12—H12B	0.9900

C6—C5	1.398 (3)	C13—H13A	0.9800
C6—C1	1.403 (3)	C13—H13B	0.9800
C1—C2	1.412 (3)	C13—H13C	0.9800
C1—C7	1.454 (3)		
C11—O1—C12	114.54 (14)	C5—C4—H4	119.4
C8—N5—N6	108.83 (14)	C3—C4—H4	119.4
C8—N5—C6	126.11 (16)	C2—C3—C4	120.26 (19)
N6—N5—C6	125.03 (15)	C2—C3—H3	119.9
C9—N3—C8	133.29 (16)	C4—C3—H3	119.9
C9—N3—C7	105.59 (16)	N1—C9—N3	108.74 (17)
C8—N3—C7	121.08 (16)	N1—C9—C10	126.52 (18)
C8—N4—C11	100.42 (14)	N3—C9—C10	124.74 (17)
C11—N6—N5	100.92 (14)	C3—C2—C1	120.2 (2)
C9—N1—N2	108.99 (16)	C3—C2—H2	119.9
C7—N2—N1	106.98 (15)	C1—C2—H2	119.9
C5—C6—C1	122.08 (18)	C9—C10—H10A	109.5
C5—C6—N5	121.37 (18)	C9—C10—H10B	109.5
C1—C6—N5	116.54 (16)	H10A—C10—H10B	109.5
C6—C1—C2	118.41 (18)	C9—C10—H10C	109.5
C6—C1—C7	118.80 (16)	H10A—C10—H10C	109.5
C2—C1—C7	122.79 (19)	H10B—C10—H10C	109.5
N4—C8—N5	112.61 (16)	O1—C12—C13	108.09 (16)
N4—C8—N3	129.94 (16)	O1—C12—H12A	110.1
N5—C8—N3	117.45 (16)	C13—C12—H12A	110.1
N6—C11—O1	123.95 (17)	O1—C12—H12B	110.1
N6—C11—N4	117.22 (16)	C13—C12—H12B	110.1
O1—C11—N4	118.83 (15)	H12A—C12—H12B	108.4
C4—C5—C6	117.8 (2)	C12—C13—H13A	109.5
C4—C5—H5	121.1	C12—C13—H13B	109.5
C6—C5—H5	121.1	H13A—C13—H13B	109.5
N2—C7—N3	109.70 (17)	C12—C13—H13C	109.5
N2—C7—C1	130.31 (17)	H13A—C13—H13C	109.5
N3—C7—C1	119.98 (17)	H13B—C13—H13C	109.5
C5—C4—C3	121.25 (19)		
C8—N5—N6—C11	0.41 (17)	C8—N4—C11—O1	178.70 (15)
C6—N5—N6—C11	-177.68 (15)	C1—C6—C5—C4	1.1 (3)
C9—N1—N2—C7	-0.1 (2)	N5—C6—C5—C4	-179.63 (15)
C8—N5—C6—C5	-178.46 (16)	N1—N2—C7—N3	-0.09 (19)
N6—N5—C6—C5	-0.7 (3)	N1—N2—C7—C1	178.94 (18)
C8—N5—C6—C1	0.9 (2)	C9—N3—C7—N2	0.26 (19)
N6—N5—C6—C1	178.64 (15)	C8—N3—C7—N2	178.22 (14)
C5—C6—C1—C2	-0.6 (3)	C9—N3—C7—C1	-178.89 (15)
N5—C6—C1—C2	-179.99 (15)	C8—N3—C7—C1	-0.9 (2)
C5—C6—C1—C7	179.69 (16)	C6—C1—C7—N2	-179.24 (17)
N5—C6—C1—C7	0.3 (2)	C2—C1—C7—N2	1.1 (3)
C11—N4—C8—N5	0.72 (18)	C6—C1—C7—N3	-0.3 (2)

C11—N4—C8—N3	180.00 (16)	C2—C1—C7—N3	-179.94 (15)
N6—N5—C8—N4	-0.8 (2)	C6—C5—C4—C3	-0.9 (3)
C6—N5—C8—N4	177.30 (15)	C5—C4—C3—C2	0.2 (3)
N6—N5—C8—N3	179.86 (13)	N2—N1—C9—N3	0.3 (2)
C6—N5—C8—N3	-2.1 (2)	N2—N1—C9—C10	-179.39 (17)
C9—N3—C8—N4	0.1 (3)	C8—N3—C9—N1	-177.93 (16)
C7—N3—C8—N4	-177.21 (17)	C7—N3—C9—N1	-0.3 (2)
C9—N3—C8—N5	179.34 (16)	C8—N3—C9—C10	1.8 (3)
C7—N3—C8—N5	2.0 (2)	C7—N3—C9—C10	179.35 (16)
N5—N6—C11—O1	-179.07 (15)	C4—C3—C2—C1	0.2 (3)
N5—N6—C11—N4	0.04 (19)	C6—C1—C2—C3	0.0 (3)
C12—O1—C11—N6	0.7 (2)	C7—C1—C2—C3	179.65 (17)
C12—O1—C11—N4	-178.38 (14)	C11—O1—C12—C13	177.76 (15)
C8—N4—C11—N6	-0.47 (19)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C3—H3...N4 <sup>i</sup>	0.95	2.60	3.533 (2)	166

Symmetry code: (i)  $x-1, y, z-1$ .