

Hexakis(dimethyl sulfoxide- κO)nickel(II) bis(2,2-dicyanoethene-1,1-dithiolato- $\kappa^2 S,S'$)nickelate(II)

Meiju Niu,^{a*} Shumei Fan^b and Guihua Liu^a

^aSchool of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China, and ^bDongchang College Liaocheng University, Shandong 252000, People's Republic of China

Correspondence e-mail: niumeiju@163.com


Received 28 November 2011; accepted 14 December 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.038; wR factor = 0.095; data-to-parameter ratio = 17.0.

The reaction of $NiCl_2 \cdot 6H_2O$ with sodium 2,2-dicyanoethene-1,1-dithiolate [$Na_2(i\text{-mnt})$] in dimethyl sulfoxide produces the title complex, $[Ni(C_2H_6OS)_6][Ni(C_4N_2S_2)_2]$. There is half each of an $[Ni(C_2H_6OS)_6]^{2-}$ complex anion and an $[Ni\{(CH_3)_2SO\}_6]^{2+}$ complex cation in the asymmetric unit. The *i*-mnt ligand coordinates in a bidentate manner to the Ni atom in the anion through the two chelating S atoms in an approximate square-planar geometry. The Ni atom in the complex cation has an octahedral coordination environment with six dimethyl sulfoxide molecules as ligands.

Related literature

For related structures, see Gao *et al.* (2004, 2005); Yu *et al.* (2005); Chen & Yu (2005).



Experimental

Crystal data

$[Ni(C_2H_6OS)_6][Ni(C_4N_2S_2)_2]$	$V = 1930.6$ (4) \AA^3
$M_r = 866.55$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 8.3368$ (10) \AA	$\mu = 1.55 \text{ mm}^{-1}$
$b = 12.6763$ (17) \AA	$T = 298$ K
$c = 18.710$ (2) \AA	$0.50 \times 0.48 \times 0.05$ mm
$\beta = 102.466$ (2) $^\circ$	

Data collection

Bruker SMART CCD area-detector diffractometer	9458 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	3379 independent reflections
$T_{\min} = 0.511$, $T_{\max} = 0.921$	2328 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	199 parameters
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
3379 reflections	$\Delta\rho_{\text{min}} = -0.35 \text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Ni1–O3	2.052 (2)	Ni2–S4	2.2010 (11)
Ni1–O2	2.060 (3)	Ni2–S5	2.2030 (11)
Ni1–O1	2.064 (2)		
O3–Ni1–O2	90.14 (11)	S4–Ni2–S5	78.96 (4)

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Natural Science Foundation of China (Nos. 20671048, 21041002).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2141).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, H. M. & Yu, H. L. (2005). *Chin. J. Spectrosc. Lab.* **22**, 740–742.
- Gao, X. K., Dou, J. M. & Dong, F. Y. (2004). *J. Inorg. Organomet. Polym.* **14**, 227–237.
- Gao, X. K., Dou, J. M. & Li, D. C. (2005). *J. Mol. Struct.* **733**, 181–186.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yu, H. L., Chen, H. M. & Dou, J. M. (2005). *Chin. J. Spectrosc. Lab.* **22**, 556–558.

supporting information

Acta Cryst. (2012). E68, m67 [doi:10.1107/S1600536811053827]

Hexakis(dimethyl sulfoxide- κO)nickel(II) bis(2,2-dicyanoethene-1,1-dithiolato- $\kappa^2 S,S'$)nickelate(II)

Meiju Niu, Shumei Fan and Guihua Liu

S1. Comment

The bridging ligand 1,1-dicyanoethene-2,2-dithiolate has been attracting more and more attention due to its delocalized π electron system able to build special planar conjugated structures (Yu *et al.*, 2005). The title complex consists of one $[\text{Ni}(i\text{-mnt})_2]^{2-}$ (where *i*-mnt is 1,1-dicyanoethene-2,2-dithiolate) complex anion and one $[\text{Ni}((\text{CH}_3)_2\text{SO})_6]^{2+}$ complex cation. For the $[\text{Ni}(i\text{-mnt})_2]^{2-}$ complex anion, the NiS_4 group is square planar and tortured slightly at an angle of 2.90 (14) $^\circ$ with respect to the plane of *i*-mnt ligand (Chen *et al.*, 2005). Four Ni-S bonds present two comparable distances of 2.2010 (11) and 2.2030 (11) \AA , similar to the Ni-S distance of 2.172 (3)


\AA in $[\text{K}(\text{N}18\text{C}6)]_2[\text{Ni}(\text{mnt})_2]$ (Gao *et al.*, 2004). The average S-C, C≡N, C-C and C=C bond lengths were 1.717, 1.141, 1.418, 1.379 \AA , respectively (Gao *et al.*, 2005). The $[\text{Ni}((\text{CH}_3)_2\text{SO})_6]^{2+}$ complex cation contains a nickel with six-coordinated octahedral geometry.

S2. Experimental

The title compound, $[\text{Ni}((\text{CH}_3)_2\text{SO})_6][\text{Ni}(i\text{-mnt})_2]$ was synthesized by the reaction of 0.05 mmol $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ and 0.1 mmol $\text{Na}_2(i\text{-mnt})(1,1\text{-dicyanoethene-2,2-dithiolate sodium})$ in 5 ml water. The solution was stirred for 2 hours and then filtered. The precipitate was dissolved in 10ml Dimethyl Sulfoxide. After slow evaporation of the solution over one month, deep green crystals suitable for X-ray diffraction were obtained (56.8%, m.p. 527–529 K).

S3. Refinement


All H atoms were placed in geometrically idealized positions and treated as riding on their parent atoms, with C—H 0.96 \AA , with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$, and refined as riding on their parent atoms.

**Figure 1**

The molecular structure of the title compound with atom labels and 40% probability displacement ellipsoids for non-H atoms. Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$, (ii) $-x + 2, -y, -z + 1$.

Hexakis(dimethyl sulfoxide- κ O)nickel(II) bis(2,2-dicyanoethene-1,1-dithiolato- κ^2S,S')nickelate(II)

Crystal data



$M_r = 866.55$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.3368 (10)$ Å

$b = 12.6763 (17)$ Å

$c = 18.710 (2)$ Å

$\beta = 102.466 (2)^\circ$

$V = 1930.6 (4)$ Å³

$Z = 2$

$F(000) = 896$

$D_x = 1.491 \text{ Mg m}^{-3}$

Mo $\text{K}\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2760 reflections

$\theta = 2.5\text{--}24.3^\circ$

$\mu = 1.55 \text{ mm}^{-1}$

$T = 298$ K

Needle, green

$0.50 \times 0.48 \times 0.05$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.511$, $T_{\max} = 0.921$

9458 measured reflections

3379 independent reflections

2328 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.0^\circ$

$h = -9 \rightarrow 9$

$k = -12 \rightarrow 15$

$l = -22 \rightarrow 18$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.038$$

$$wR(F^2) = 0.095$$

$$S = 1.06$$

3379 reflections

199 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0275P)^2 + 1.0073P]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.30 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.35 \text{ e \AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	0.5000	0.5000	0.02973 (18)
Ni2	1.0000	0.0000	0.5000	0.0415 (2)
N1	0.7064 (6)	0.2565 (4)	0.2505 (2)	0.0863 (15)
N2	1.1494 (6)	0.0945 (4)	0.2174 (2)	0.0751 (13)
O1	0.4945 (3)	0.64165 (18)	0.44513 (13)	0.0367 (6)
O2	0.7372 (3)	0.47506 (19)	0.48840 (15)	0.0440 (7)
O3	0.4122 (3)	0.41769 (19)	0.40520 (13)	0.0436 (7)
S1	0.62727 (13)	0.72370 (8)	0.46849 (5)	0.0412 (3)
S2	0.78272 (12)	0.36249 (8)	0.47193 (5)	0.0385 (3)
S3	0.32535 (13)	0.47016 (8)	0.33476 (5)	0.0391 (3)
S4	0.81754 (14)	0.08901 (9)	0.41999 (6)	0.0489 (3)
S5	1.10676 (15)	-0.00988 (9)	0.40228 (6)	0.0533 (3)
C1	0.5310 (7)	0.8291 (4)	0.5044 (3)	0.095 (2)
H1A	0.4416	0.8553	0.4675	0.142*
H1B	0.6092	0.8846	0.5198	0.142*
H1C	0.4898	0.8047	0.5456	0.142*
C2	0.6505 (7)	0.7821 (4)	0.3864 (2)	0.0732 (16)
H2A	0.6983	0.7320	0.3586	0.110*
H2B	0.7208	0.8426	0.3971	0.110*
H2C	0.5449	0.8036	0.3586	0.110*
C3	0.9350 (5)	0.3786 (3)	0.4198 (2)	0.0503 (11)
H3A	0.8874	0.4120	0.3741	0.075*
H3B	0.9777	0.3108	0.4106	0.075*
H3C	1.0224	0.4216	0.4465	0.075*
C4	0.9093 (6)	0.3174 (4)	0.5552 (2)	0.0585 (13)
H4A	0.9998	0.3651	0.5701	0.088*
H4B	0.9504	0.2484	0.5480	0.088*
H4C	0.8462	0.3144	0.5924	0.088*
C5	0.2206 (6)	0.3635 (4)	0.2835 (2)	0.0676 (14)
H5A	0.2961	0.3066	0.2827	0.101*
H5B	0.1756	0.3863	0.2343	0.101*
H5C	0.1334	0.3399	0.3057	0.101*
C6	0.4776 (6)	0.4904 (4)	0.2835 (2)	0.0600 (13)
H6A	0.5526	0.5440	0.3065	0.090*

H6B	0.4265	0.5123	0.2348	0.090*
H6C	0.5365	0.4258	0.2813	0.090*
C7	0.9525 (5)	0.0750 (3)	0.3630 (2)	0.0409 (10)
C8	0.9395 (5)	0.1255 (3)	0.2967 (2)	0.0427 (10)
C9	0.8105 (6)	0.1979 (4)	0.2706 (2)	0.0526 (12)
C10	1.0561 (6)	0.1086 (3)	0.2528 (2)	0.0522 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0336 (4)	0.0269 (4)	0.0290 (4)	0.0013 (3)	0.0074 (3)	-0.0017 (3)
Ni2	0.0476 (5)	0.0370 (4)	0.0408 (4)	0.0016 (4)	0.0116 (3)	-0.0019 (3)
N1	0.092 (4)	0.077 (3)	0.076 (3)	0.030 (3)	-0.011 (3)	-0.008 (2)
N2	0.080 (3)	0.093 (3)	0.058 (3)	0.006 (3)	0.029 (2)	0.002 (2)
O1	0.0380 (16)	0.0297 (14)	0.0415 (14)	-0.0024 (12)	0.0067 (12)	0.0023 (12)
O2	0.0373 (17)	0.0336 (16)	0.0633 (18)	-0.0007 (12)	0.0155 (14)	-0.0079 (13)
O3	0.061 (2)	0.0358 (15)	0.0309 (14)	-0.0010 (14)	0.0039 (13)	-0.0047 (12)
S1	0.0402 (7)	0.0377 (6)	0.0461 (6)	-0.0039 (5)	0.0104 (5)	0.0007 (5)
S2	0.0339 (6)	0.0327 (6)	0.0497 (6)	0.0002 (4)	0.0104 (5)	-0.0077 (5)
S3	0.0409 (6)	0.0434 (6)	0.0325 (5)	0.0031 (5)	0.0071 (4)	-0.0050 (4)
S4	0.0502 (8)	0.0503 (7)	0.0486 (6)	0.0102 (5)	0.0157 (5)	-0.0005 (5)
S5	0.0537 (8)	0.0614 (8)	0.0473 (6)	0.0163 (6)	0.0161 (5)	0.0049 (6)
C1	0.094 (5)	0.053 (3)	0.155 (6)	-0.023 (3)	0.067 (4)	-0.048 (4)
C2	0.086 (4)	0.072 (3)	0.063 (3)	-0.032 (3)	0.018 (3)	0.015 (3)
C3	0.045 (3)	0.048 (3)	0.063 (3)	-0.003 (2)	0.022 (2)	-0.011 (2)
C4	0.067 (3)	0.049 (3)	0.056 (3)	0.014 (2)	0.004 (2)	0.000 (2)
C5	0.077 (4)	0.074 (3)	0.049 (3)	-0.025 (3)	0.008 (2)	-0.017 (3)
C6	0.062 (3)	0.064 (3)	0.061 (3)	-0.005 (3)	0.030 (3)	0.006 (2)
C7	0.045 (3)	0.035 (2)	0.041 (2)	-0.0014 (19)	0.0061 (19)	-0.0096 (18)
C8	0.045 (3)	0.042 (3)	0.041 (2)	0.002 (2)	0.008 (2)	-0.0053 (19)
C9	0.066 (3)	0.049 (3)	0.040 (2)	-0.003 (3)	0.005 (2)	-0.010 (2)
C10	0.065 (4)	0.051 (3)	0.040 (3)	-0.002 (2)	0.008 (2)	0.001 (2)

Geometric parameters (\AA , $^\circ$)

Ni1—O3	2.052 (2)	S5—C7	1.716 (4)
Ni1—O3 ⁱ	2.052 (2)	C1—H1A	0.9600
Ni1—O2 ⁱ	2.060 (3)	C1—H1B	0.9600
Ni1—O2	2.060 (3)	C1—H1C	0.9600
Ni1—O1	2.064 (2)	C2—H2A	0.9600
Ni1—O1 ⁱ	2.064 (2)	C2—H2B	0.9600
Ni2—S4	2.2010 (11)	C2—H2C	0.9600
Ni2—S4 ⁱⁱ	2.2010 (11)	C3—H3A	0.9600
Ni2—S5 ⁱⁱ	2.2030 (11)	C3—H3B	0.9600
Ni2—S5	2.2030 (11)	C3—H3C	0.9600
N1—C9	1.142 (6)	C4—H4A	0.9600
N2—C10	1.139 (5)	C4—H4B	0.9600
O1—S1	1.513 (3)	C4—H4C	0.9600

O2—S2	1.525 (3)	C5—H5A	0.9600
O3—S3	1.514 (3)	C5—H5B	0.9600
S1—C2	1.752 (4)	C5—H5C	0.9600
S1—C1	1.765 (5)	C6—H6A	0.9600
S2—C3	1.772 (4)	C6—H6B	0.9600
S2—C4	1.777 (4)	C6—H6C	0.9600
S3—C6	1.767 (4)	C7—C8	1.378 (5)
S3—C5	1.775 (4)	C8—C9	1.418 (6)
S4—C7	1.719 (4)	C8—C10	1.418 (6)
O3—Ni1—O3 ⁱ	180.000 (1)	H1B—C1—H1C	109.5
O3—Ni1—O2 ⁱ	89.86 (11)	S1—C2—H2A	109.5
O3 ⁱ —Ni1—O2 ⁱ	90.14 (11)	S1—C2—H2B	109.5
O3—Ni1—O2	90.14 (11)	H2A—C2—H2B	109.5
O3 ⁱ —Ni1—O2	89.86 (11)	S1—C2—H2C	109.5
O2 ⁱ —Ni1—O2	180.00 (15)	H2A—C2—H2C	109.5
O3—Ni1—O1	92.69 (9)	H2B—C2—H2C	109.5
O3 ⁱ —Ni1—O1	87.31 (9)	S2—C3—H3A	109.5
O2 ⁱ —Ni1—O1	90.01 (10)	S2—C3—H3B	109.5
O2—Ni1—O1	89.99 (10)	H3A—C3—H3B	109.5
O3—Ni1—O1 ⁱ	87.31 (9)	S2—C3—H3C	109.5
O3 ⁱ —Ni1—O1 ⁱ	92.69 (9)	H3A—C3—H3C	109.5
O2 ⁱ —Ni1—O1 ⁱ	89.99 (10)	H3B—C3—H3C	109.5
O2—Ni1—O1 ⁱ	90.01 (10)	S2—C4—H4A	109.5
O1—Ni1—O1 ⁱ	180.0	S2—C4—H4B	109.5
S4—Ni2—S4 ⁱⁱ	180.00 (5)	H4A—C4—H4B	109.5
S4—Ni2—S5 ⁱⁱ	101.04 (4)	S2—C4—H4C	109.5
S4 ⁱⁱ —Ni2—S5 ⁱⁱ	78.96 (4)	H4A—C4—H4C	109.5
S4—Ni2—S5	78.96 (4)	H4B—C4—H4C	109.5
S4 ⁱⁱ —Ni2—S5	101.04 (4)	S3—C5—H5A	109.5
S5 ⁱⁱ —Ni2—S5	180.0	S3—C5—H5B	109.5
S1—O1—Ni1	121.20 (14)	H5A—C5—H5B	109.5
S2—O2—Ni1	116.83 (15)	S3—C5—H5C	109.5
S3—O3—Ni1	122.93 (15)	H5A—C5—H5C	109.5
O1—S1—C2	104.45 (19)	H5B—C5—H5C	109.5
O1—S1—C1	105.4 (2)	S3—C6—H6A	109.5
C2—S1—C1	99.2 (3)	S3—C6—H6B	109.5
O2—S2—C3	104.06 (18)	H6A—C6—H6B	109.5
O2—S2—C4	104.50 (18)	S3—C6—H6C	109.5
C3—S2—C4	99.1 (2)	H6A—C6—H6C	109.5
O3—S3—C6	105.8 (2)	H6B—C6—H6C	109.5
O3—S3—C5	102.88 (19)	C8—C7—S5	125.7 (3)
C6—S3—C5	98.3 (2)	C8—C7—S4	125.1 (3)
C7—S4—Ni2	85.42 (14)	S5—C7—S4	109.2 (2)
C7—S5—Ni2	85.44 (14)	C7—C8—C9	121.2 (4)
S1—C1—H1A	109.5	C7—C8—C10	121.3 (4)
S1—C1—H1B	109.5	C9—C8—C10	117.5 (4)
H1A—C1—H1B	109.5	N1—C9—C8	179.0 (5)

S1—C1—H1C	109.5	N2—C10—C8	179.7 (6)
H1A—C1—H1C	109.5		
O3—Ni1—O1—S1	-142.43 (17)	Ni1—O3—S3—C5	-161.4 (2)
O3 ⁱ —Ni1—O1—S1	37.57 (17)	S4 ⁱⁱ —Ni2—S4—C7	-120 (100)
O2 ⁱ —Ni1—O1—S1	127.71 (17)	S5 ⁱⁱ —Ni2—S4—C7	173.58 (13)
O2—Ni1—O1—S1	-52.29 (17)	S5—Ni2—S4—C7	-6.42 (13)
O1 ⁱ —Ni1—O1—S1	-139 (100)	S4—Ni2—S5—C7	6.44 (13)
O3—Ni1—O2—S2	-48.30 (17)	S4 ⁱⁱ —Ni2—S5—C7	-173.56 (13)
O3 ⁱ —Ni1—O2—S2	131.70 (17)	S5 ⁱⁱ —Ni2—S5—C7	112 (14)
O2 ⁱ —Ni1—O2—S2	13 (48)	Ni2—S5—C7—C8	170.5 (3)
O1—Ni1—O2—S2	-140.99 (17)	Ni2—S5—C7—S4	-8.58 (17)
O1 ⁱ —Ni1—O2—S2	39.01 (17)	Ni2—S4—C7—C8	-170.5 (3)
O3 ⁱ —Ni1—O3—S3	-178 (100)	Ni2—S4—C7—S5	8.59 (18)
O2 ⁱ —Ni1—O3—S3	71.7 (2)	S5—C7—C8—C9	-177.2 (3)
O2—Ni1—O3—S3	-108.3 (2)	S4—C7—C8—C9	1.7 (6)
O1—Ni1—O3—S3	-18.3 (2)	S5—C7—C8—C10	1.6 (6)
O1 ⁱ —Ni1—O3—S3	161.7 (2)	S4—C7—C8—C10	-179.4 (3)
Ni1—O1—S1—C2	145.1 (2)	C7—C8—C9—N1	26 (32)
Ni1—O1—S1—C1	-110.9 (3)	C10—C8—C9—N1	-153 (32)
Ni1—O2—S2—C3	149.93 (19)	C7—C8—C10—N2	79 (100)
Ni1—O2—S2—C4	-106.6 (2)	C9—C8—C10—N2	-102 (100)
Ni1—O3—S3—C6	95.9 (2)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y, -z+1$.