

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-N-(1-Benzothiophen-3-ylmethylidene)-2,6-dimethylaniline

 Nermin Kahveci Yağcı,^a Necmi Dege,^{b*} Sümeyye Gümüş,^c Erbil Açar^c and Mustafa Serkan Soylu^d

^aKırıkkale University, Faculty of Arts and Sciences, Physics Department, 71450 Kırıkkale, Turkey, ^bOndokuz Mayıs University, Arts and Sciences Faculty, Department of Physics, 55139 Samsun, Turkey, ^cOndokuz Mayıs University, Arts and Sciences Faculty, Department of Chemistry, 55139 Samsun, Turkey, and ^dGiresun University, Faculty of Arts and Sciences, Department of Physics, 28100 Giresun, Turkey

Correspondence e-mail: necmismusun@gmail.com


Received 26 January 2012; accepted 31 January 2012

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.040; wR factor = 0.101; data-to-parameter ratio = 11.6.

In the title compound, $\text{C}_{17}\text{H}_{15}\text{NS}$, the benzothiophene residue and the substituted benzene ring are oriented at a dihedral angle of $61.99(7)^\circ$. An intermolecular $\text{C}-\text{H}\cdots\pi$ interaction contributes to the stability of the crystal structure.

Related literature

For the biological properties of Schiff bases, see: Barton & Ollis (1979); Layer (1963); Ingold (1969). For industrial applications of Schiff bases, see: Taggi *et al.* (2002). For chemical properties of Schiff bases, see: Aydoğın *et al.* (2001); Tanak *et al.* (2010); Ingold (1969). For related structures, see: Açar *et al.* (2010); Ceylan *et al.* (2011); Dege, Şekerci *et al.* (2006); Demirtaş *et al.* (2009); Dege, İçbudak & Adıyaman (2006, 2007); Genç *et al.* (2004); İnaç *et al.* (2012); Tecer *et al.* (2010). For the structural properties benzothiophene derivatives, see: Alarcon *et al.* (1999); Cohen *et al.* (1964); Hadjoudis *et al.* (1987); Inamoto *et al.* (2008); Köysal *et al.* (2007); Karabiyik *et al.* (2008); Kobayashi *et al.* (2009); Mlochowski & Potaczek (2009); Novopoltseva (1995); Tanak *et al.* (2010); Xu *et al.* (1994); Zhang *et al.* (2001).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{15}\text{NS}$	$V = 719.12(14) \text{ \AA}^3$
$M_r = 265.36$	$Z = 2$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
$a = 8.0446(10) \text{ \AA}$	$\mu = 0.21 \text{ mm}^{-1}$
$b = 8.8031(9) \text{ \AA}$	$T = 296 \text{ K}$
$c = 10.2809(10) \text{ \AA}$	$0.17 \times 0.15 \times 0.12 \text{ mm}$
$\beta = 98.989(11)^\circ$	

Data collection

Oxford Diffraction SuperNova (single source at offset) Eos diffractometer	Diffraction, 2007
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford)	$T_{\min} = 0.965$, $T_{\max} = 0.975$
	2788 measured reflections
	2024 independent reflections
	1722 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.101$	$\Delta\rho_{\max} = 0.15 \text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$
2024 reflections	Absolute structure: Flack (1983),
174 parameters	497 Friedel pairs
1 restraint	Flack parameter: 0.10 (11)

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C1–C6 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C9}-\text{H9}\cdots\text{Cg2}^i$	0.93	2.95	3.872 (4)	171

 Symmetry code: (i) $-x, y - \frac{1}{2}, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009) and *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *OLEX2*, *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors thank Ondokuz Mayıs University, Giresun University and Kırıkkale University, Turkey, for financial support of this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5806).

References

- Açar, A., Tanak, H. & Yavuz, M. (2010). *Mol. Phys.* **108**, 1759–1772.
 Alarcon, S. H., Pagani, D., Bacigalupo, J. & Olivieri, A. C. (1999). *J. Mol. Struct.* **475**, 233–240.
 Aydoğın, F., Öcal, N., Turgut, Z. & Yolaçan, C. (2001). *Bull. Korean Chem. Soc.* **22**, 476–480.
 Barton, D. & Ollis, W. D. (1979). *Comprehensive Organic Chemistry*, Vol 2. Oxford: Pergamon.
 Ceylan, Ü., Tanak, H., Gümüş, S. & Açar, E. (2011). *Acta Cryst.* **E67**, o2004.
 Cohen, M. D., Schmidt, G. M. J. & Flavian, S. (1964). *J. Chem. Soc.* pp. 2041–2051.
 Dege, N., İçbudak, H. & Adıyaman, E. (2006). *Acta Cryst.* **C62**, m401–m403.
 Dege, N., İçbudak, H. & Adıyaman, E. (2007). *Acta Cryst.* **C63**, m13–m15.

- Dege, N., Şekerci, M., Servi, S., Dinçer, M. & Demirbaş, Ü. (2006). *Turk. J. Chem.* **30**, 103–108.
- Demirtaş, G., Dege, N., Şekerci, M., Servi, S. & Dinçer, M. (2009). *Acta Cryst. E65*, o1668.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Genç, S., Dege, N., Yılmaz, I., Çukurovalı, A. & Dinçer, M. (2004). *Acta Cryst. E60*, e10.
- Hadjoudis, E., Vitterakis, M. & Mavridis, I. M. (1987). *Tetrahedron*, **43**, 1345–1360.
- Inaç, H., Dege, N., Gümüş, S., Açar, E. & Soylu, M. S. (2012). *Acta Cryst. E68*, o361.
- Inamoto, K., Arai, Y., Hiroya, K. & Doi, T. (2008). *Chem. Commun.* pp. 5529–5531.
- Ingold, C. K. (1969). In *Structure and Mechanism in Organic Chemistry*, 2nd ed. Ithaca, New York: Cornell University Press.
- Karabiyik, H., Ocak İskeleli, N., Petek, H., Albayrak, Ç. & Açar, E. (2008). *J. Mol. Struct.* **873**, 130–136.
- Kobayashi, K., Egara, Y., Fukamachi, S. & Konishi, H. (2009). *Tetrahedron*, **65**, 9633–9636.
- Köysal, Y., Işık, Ş. & Açar, A. (2007). *Acta Cryst. E63*, o4916.
- Layer, R. W. (1963). *Chem. Rev.* **63**, 489–510.
- Mlochowski, J. & Potaczek, P. (2009). *Phosphorus Sulfur Silicon Relat. Elem.* **184**, 1115–1123.
- Novopoltseva, O. M. (1995). Cand. Sci. (Chem.) dissertation, University of Volgograd, Russia.
- Oxford Diffraction (2007). *CrysAlis PRO*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Taggi, A. E., Hafez, A. M., Wack, H., Young, B., Ferraris, D. & Lectka, T. (2002). *J. Am. Chem. Soc.* **124**, 6626–6635.
- Tanak, H., Açar, A. & Yavuz, M. (2010). *J. Mol. Model.* **16**, 577–587.
- Tecer, E., Dege, N., Zülfikaroğlu, A., Şenyüz, N. & Batu, H. (2010). *Acta Cryst. E66*, o3369–o3370.
- Xu, X.-X., You, X.-Z., Sun, Z.-F., Wang, X. & Liu, H.-X. (1994). *Acta Cryst. C50*, 1169–1171.
- Zhang, Y., Guo, Z. J. & You, X. Z. (2001). *J. Am. Chem. Soc.* **123**, 9378–9387.

supporting information

Acta Cryst. (2012). E68, o623–o624 [doi:10.1107/S1600536812004151]

(E)-N-(1-Benzothiophen-3-ylmethylidene)-2,6-dimethylaniline**Nermin Kahveci Yağcı, Necmi Dege, Sümeyye Gümüş, Erbil Ağar and Mustafa Serkan Soylu****S1. Comment**

Benzothiophene derivatives are undoubtedly one of the most important classes of heterocycles, because some of molecules having the benzothiophene skeleton have been reported to exhibit wide variety of biological activities (Kobayashi *et al.*, 2009). Therefore, we (İnaç *et al.* 2012) and others (İnamoto *et al.*, 2008; Mlochowski & Potaczek, 2009) have recently reported efficient methods for the synthesis of benzothiophenes. However, there are only a few reports on the synthesis of 1-(benzo[*b*]thiophen-3-yl)-*N*-methylmethanamines, which may also be of potential biological importance.

Schiff bases, *i.e.*, compounds having a double C=N bond, are used as starting materials in the synthesis of important drugs, such as antibiotics and antiallergic, antiphlogistic, and antitumor substances (Barton & Ollis, 1979; Layer, 1963; Ingold 1969). On the industrial scale, they have a wide range of applications, such as dyes and pigments (Taggi *et al.*, 2002). They are also used as components of rubber compounds (Novopoltseva, 1995). Schiff base compounds display interesting photochromic and thermochromic features in the solid state and can be classified in terms of these properties (Cohen *et al.*, 1964). Photo- and thermochromism arise *via* H-atom transfer from the hydroxy O atom to the imine N atom (Hadjoudis *et al.*, 1987; Xu *et al.*, 1994). Such proton-exchanging materials can be utilized for the design of various molecular electronic devices (Alarcon *et al.*, 1999). In general, Schiff bases display two possible tautomeric forms, the phenol-imine (OH) and the keto-amine (NH) forms. Depending on the tautomers, two types of intramolecular hydrogen bonds are observed in Schiff bases: O—H \cdots N in phenol-imine (Köysal *et al.*, 2007) and N—H \cdots O in keto-amine tautomers. By means of increasing development of computational chemistry in the past decade, the research of theoretical modeling of drug design, functional material design, *etc.*, has become more mature than ever. Many important chemical and physical properties of biological and chemical systems can be predicted from the first principles by various computational techniques (Zhang *et al.*, 2001). Schiff bases have also been employed as ligands for the complexation of metal ions (Aydoğan *et al.*, 2001).

The molecular structure is not planar (Fig.1); the dihedral angle between the C10—C17 benzene and the C1—C8/S1 benzothiophene ring is 61.99 (7)°. The dihedral angle between the methylenemethanamine and benzothiophene group is 3.22 (50)°. The length of the C9=N1 double bond is 1.260 (3) Å, slightly shorter than standard 1.28 Å value of a C=N double bond and consistent with related structures (Ağar *et al.*, 2010; Ceylan *et al.* 2011; Dege, Şekerçi *et al.*, 2006; Genç *et al.* 2004; İnaç *et al.*, 2012; Tanak *et al.*, 2010; Tecer *et al.*, 2010).

The C1—S1 and C8—S1 bond distances are 1.736 (3) Å and 1.717 (3) Å, respectively. The C—S bond distances are compatible with the literature (Dege, İçbudak & Adıyaman, 2006, 2007; Demirtaş *et al.*, 2009).

The crystal structure is stabilized by an intermolecular C—H \cdots π stacking interaction (C9—H9 \cdots Cg2ⁱ = 2.95 Å) [symmetry code (i): -x, -1/2 + y, 1 - z; Cg2 is the centroids of ring C1—C6].

S2. Experimental

The compound (*E*)-1-(1-benzothiophen-3-yl)-*N*-(2,6-dimethylphenyl)methanimine was prepared by reflux a mixture of a solution containing 1-benzothiophene-3-carbaldehyde (0.026 g 0.16 mmol) in 20 ml ethanol and a solution containing 2,6-Dimethylaniline (0.0192 g 0.16 mmol) in 20 ml ethanol. The reaction mixture was stirred for 1 h under reflux. The crystals of 2(*E*)-1-(1-benzothiophen-3-yl)-*N*-(2,6-dimethylphenyl)methanimine suitable for X-ray analysis were obtained from ethylalcohol by slow evaporation (yield % 58; m.p 75–77°C).

S3. Refinement

All hydrogen atoms were positioned geometrically [$C-H=0.930$ and 0.960] and treated as riding with $U_{iso}(H)=1.2U_{eq}(C)$ or $1.5U_{eq}(C_{methyl})$.




Figure 1

The molecular structure of the title compound, showing the atom-numbering scheme and 50% probability displacement ellipsoids.

(*E*)-*N*-(1-Benzothiophen-3-ylmethylidene)-2,6-dimethylaniline

Crystal data

$C_{17}H_{15}NS$

$M_r = 265.36$

Monoclinic, $P2_1$

Hall symbol: $P\ 2_1$

$a = 8.0446$ (10) Å

$b = 8.8031$ (9) Å

$c = 10.2809$ (10) Å

$\beta = 98.989$ (11)°

$V = 719.12$ (14) Å³

$Z = 2$

$F(000) = 280$

$D_x = 1.225$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1387 reflections

$\theta = 3.5$ – 27.3 °

$\mu = 0.21$ mm⁻¹

$T = 296$ K

Prism, brown

$0.17 \times 0.15 \times 0.12$ mm

Data collection

Oxford Diffraction SuperNova (single source at offset) Eos diffractometer

Radiation source: SuperNova (Mo) X-ray

Source

Mirror monochromator

Detector resolution: 16.0454 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Oxford Diffraction, 2007)

$T_{min} = 0.965$, $T_{max} = 0.975$

2788 measured reflections

2024 independent reflections
 1722 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
 $\theta_{\text{max}} = 27.4^\circ$, $\theta_{\text{min}} = 3.5^\circ$

$h = -10 \rightarrow 5$
 $k = -5 \rightarrow 11$
 $l = -10 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.101$
 $S = 1.05$
 2024 reflections
 174 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0397P)^2 + 0.1075P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.15 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.23 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983), 497 Friedel
 pairs
 Absolute structure parameter: 0.10 (11)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.82893 (13)	0.27137 (12)	0.24201 (7)	0.0801 (3)
N1	0.9883 (3)	0.2484 (3)	0.7306 (2)	0.0511 (6)
C10	1.0962 (3)	0.1995 (3)	0.8467 (2)	0.0491 (7)
C7	0.9176 (4)	0.2495 (4)	0.4948 (2)	0.0565 (7)
C6	0.7680 (3)	0.3424 (3)	0.4760 (2)	0.0453 (6)
C11	1.0246 (4)	0.1127 (4)	0.9370 (3)	0.0567 (8)
C5	0.6838 (4)	0.4137 (4)	0.5683 (3)	0.0560 (8)
H5	0.7222	0.4028	0.6579	0.067*
C2	0.5615 (4)	0.4471 (4)	0.2995 (3)	0.0659 (9)
H2	0.5210	0.4585	0.2103	0.079*
C13	1.2877 (5)	0.1134 (5)	1.0815 (3)	0.0766 (11)
H13	1.3528	0.0846	1.1605	0.092*
C15	1.2650 (4)	0.2449 (4)	0.8730 (3)	0.0590 (8)
C1	0.7043 (4)	0.3622 (3)	0.3416 (3)	0.0551 (8)
C9	1.0162 (4)	0.2036 (4)	0.6197 (3)	0.0575 (8)
H9	1.1054	0.1370	0.6173	0.069*
C14	1.3571 (4)	0.2000 (5)	0.9925 (4)	0.0784 (12)
H14	1.4692	0.2294	1.0129	0.094*
C3	0.4813 (5)	0.5139 (5)	0.3930 (3)	0.0772 (11)

H3	0.3835	0.5696	0.3671	0.093*
C12	1.1233 (4)	0.0697 (4)	1.0539 (3)	0.0695 (9)
H12	1.0771	0.0105	1.1141	0.083*
C4	0.5443 (4)	0.4995 (5)	0.5254 (3)	0.0698 (10)
H4	0.4904	0.5494	0.5869	0.084*
C8	0.9607 (5)	0.2058 (4)	0.3779 (3)	0.0772 (11)
H8	1.0539	0.1452	0.3720	0.093*
C17	0.8439 (4)	0.0649 (6)	0.9089 (4)	0.0854 (12)
H17A	0.7741	0.1528	0.8880	0.128*
H17B	0.8135	0.0155	0.9850	0.128*
H17C	0.8282	-0.0040	0.8357	0.128*
C16	1.3420 (5)	0.3460 (6)	0.7799 (4)	0.0923 (12)
H16A	1.3583	0.2888	0.7033	0.138*
H16C	1.4485	0.3834	0.8231	0.138*
H16B	1.2681	0.4300	0.7538	0.138*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.1139 (8)	0.0853 (7)	0.0394 (4)	0.0236 (6)	0.0066 (4)	-0.0016 (5)
N1	0.0541 (12)	0.0557 (15)	0.0416 (11)	0.0048 (12)	0.0016 (10)	0.0060 (12)
C10	0.0502 (14)	0.0525 (17)	0.0424 (14)	0.0067 (14)	0.0003 (12)	-0.0035 (13)
C7	0.0698 (18)	0.0563 (19)	0.0427 (13)	0.0091 (16)	0.0063 (13)	0.0045 (15)
C6	0.0565 (15)	0.0399 (15)	0.0375 (13)	-0.0026 (13)	0.0012 (11)	0.0045 (12)
C11	0.0616 (17)	0.065 (2)	0.0417 (15)	0.0004 (16)	0.0021 (13)	0.0005 (15)
C5	0.0612 (18)	0.062 (2)	0.0426 (15)	0.0031 (16)	0.0024 (13)	0.0041 (15)
C2	0.078 (2)	0.065 (2)	0.0483 (16)	0.0040 (18)	-0.0110 (16)	0.0069 (16)
C13	0.073 (2)	0.099 (3)	0.0517 (18)	0.025 (2)	-0.0101 (17)	0.003 (2)
C15	0.0512 (16)	0.064 (2)	0.0607 (16)	-0.0009 (16)	0.0049 (14)	-0.0074 (18)
C1	0.075 (2)	0.0468 (17)	0.0405 (15)	-0.0068 (16)	0.0007 (14)	0.0026 (13)
C9	0.0655 (18)	0.058 (2)	0.0466 (15)	0.0141 (16)	0.0024 (14)	0.0058 (14)
C14	0.0481 (16)	0.100 (3)	0.082 (2)	0.0119 (19)	-0.0083 (17)	-0.025 (2)
C3	0.075 (2)	0.085 (3)	0.066 (2)	0.020 (2)	-0.0052 (18)	0.008 (2)
C12	0.083 (2)	0.074 (2)	0.0495 (18)	0.015 (2)	0.0025 (17)	0.0097 (18)
C4	0.069 (2)	0.081 (3)	0.058 (2)	0.0190 (19)	0.0062 (17)	0.0024 (18)
C8	0.098 (3)	0.081 (3)	0.0517 (18)	0.028 (2)	0.0096 (17)	0.0013 (18)
C17	0.072 (2)	0.114 (3)	0.068 (2)	-0.025 (2)	0.0046 (19)	0.016 (2)
C16	0.077 (2)	0.099 (3)	0.102 (3)	-0.019 (2)	0.020 (2)	-0.005 (2)

Geometric parameters (Å, °)

S1—C8	1.717 (3)	C13—C12	1.363 (5)
S1—C1	1.736 (3)	C13—C14	1.375 (5)
N1—C9	1.260 (3)	C13—H13	0.9300
N1—C10	1.428 (3)	C15—C14	1.390 (4)
C10—C11	1.395 (4)	C15—C16	1.509 (5)
C10—C15	1.400 (4)	C9—H9	0.9300
C7—C8	1.357 (4)	C14—H14	0.9300

C7—C6	1.442 (4)	C3—C4	1.382 (5)
C7—C9	1.456 (4)	C3—H3	0.9300
C6—C5	1.398 (4)	C12—H12	0.9300
C6—C1	1.407 (3)	C4—H4	0.9300
C11—C12	1.386 (4)	C8—H8	0.9300
C11—C17	1.497 (4)	C17—H17A	0.9600
C5—C4	1.367 (4)	C17—H17B	0.9600
C5—H5	0.9300	C17—H17C	0.9600
C2—C3	1.371 (5)	C16—H16A	0.9600
C2—C1	1.382 (4)	C16—H16C	0.9600
C2—H2	0.9300	C16—H16B	0.9600
C8—S1—C1	90.84 (15)	N1—C9—H9	117.9
C9—N1—C10	119.5 (2)	C7—C9—H9	117.9
C11—C10—C15	121.1 (2)	C13—C14—C15	122.0 (3)
C11—C10—N1	117.4 (2)	C13—C14—H14	119.0
C15—C10—N1	121.3 (3)	C15—C14—H14	119.0
C8—C7—C6	111.4 (2)	C2—C3—C4	120.7 (3)
C8—C7—C9	121.5 (3)	C2—C3—H3	119.6
C6—C7—C9	127.1 (2)	C4—C3—H3	119.6
C5—C6—C1	118.0 (3)	C13—C12—C11	120.6 (3)
C5—C6—C7	130.2 (2)	C13—C12—H12	119.7
C1—C6—C7	111.7 (2)	C11—C12—H12	119.7
C12—C11—C10	119.1 (3)	C5—C4—C3	121.6 (3)
C12—C11—C17	119.9 (3)	C5—C4—H4	119.2
C10—C11—C17	121.0 (3)	C3—C4—H4	119.2
C4—C5—C6	119.3 (3)	C7—C8—S1	114.5 (3)
C4—C5—H5	120.3	C7—C8—H8	122.7
C6—C5—H5	120.3	S1—C8—H8	122.7
C3—C2—C1	118.2 (3)	C11—C17—H17A	109.5
C3—C2—H2	120.9	C11—C17—H17B	109.5
C1—C2—H2	120.9	H17A—C17—H17B	109.5
C12—C13—C14	120.0 (3)	C11—C17—H17C	109.5
C12—C13—H13	120.0	H17A—C17—H17C	109.5
C14—C13—H13	120.0	H17B—C17—H17C	109.5
C14—C15—C10	117.2 (3)	C15—C16—H16A	109.5
C14—C15—C16	120.9 (3)	C15—C16—H16C	109.5
C10—C15—C16	121.8 (3)	H16A—C16—H16C	109.5
C2—C1—C6	122.1 (3)	C15—C16—H16B	109.5
C2—C1—S1	126.4 (2)	H16A—C16—H16B	109.5
C6—C1—S1	111.5 (2)	H16C—C16—H16B	109.5
N1—C9—C7	124.2 (3)		
C9—N1—C10—C11	-117.4 (3)	C5—C6—C1—S1	177.9 (2)
C9—N1—C10—C15	66.6 (4)	C7—C6—C1—S1	-1.2 (3)
C8—C7—C6—C5	-178.2 (3)	C8—S1—C1—C2	179.8 (3)
C9—C7—C6—C5	1.9 (6)	C8—S1—C1—C6	1.0 (2)
C8—C7—C6—C1	0.7 (4)	C10—N1—C9—C7	-178.1 (3)

C9—C7—C6—C1	-179.2 (3)	C8—C7—C9—N1	175.4 (4)
C15—C10—C11—C12	-0.7 (5)	C6—C7—C9—N1	-4.7 (5)
N1—C10—C11—C12	-176.8 (3)	C12—C13—C14—C15	-0.4 (6)
C15—C10—C11—C17	179.7 (3)	C10—C15—C14—C13	0.6 (5)
N1—C10—C11—C17	3.6 (5)	C16—C15—C14—C13	177.2 (4)
C1—C6—C5—C4	-0.2 (5)	C1—C2—C3—C4	1.5 (6)
C7—C6—C5—C4	178.7 (3)	C14—C13—C12—C11	-0.5 (6)
C11—C10—C15—C14	-0.1 (5)	C10—C11—C12—C13	1.0 (5)
N1—C10—C15—C14	175.8 (3)	C17—C11—C12—C13	-179.4 (4)
C11—C10—C15—C16	-176.6 (3)	C6—C5—C4—C3	2.0 (6)
N1—C10—C15—C16	-0.7 (5)	C2—C3—C4—C5	-2.7 (6)
C3—C2—C1—C6	0.3 (5)	C6—C7—C8—S1	0.1 (4)
C3—C2—C1—S1	-178.4 (3)	C9—C7—C8—S1	180.0 (3)
C5—C6—C1—C2	-1.0 (4)	C1—S1—C8—C7	-0.7 (3)
C7—C6—C1—C2	180.0 (3)		

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the (C1-C6) ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C9—H9 \cdots Cg2 ⁱ	0.93	2.95	3.872 (4)	171

Symmetry code: (i) $-x, y-1/2, -z+1$.