

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

rac-3-[(Anilino)(naphthalen-2-yl)methyl]thian-4-one

 Klaus Harms,^{a*} M. Saeed Abaee,^b Mohammad M. Mojtahedi^b and A. Wahid Mesbah^b

^aFachbereich Chemie, Philipps Universität Marburg, Hans Meerwein Strasse, Marburg, D-35032, Germany, and ^bDepartment of Organic Chemistry, Chemistry and Chemical Engineering Research Center of Iran, PO Box 14335-186, Tehran, Iran
Correspondence e-mail: harms@chemie.uni-marburg.de


Received 17 January 2012; accepted 10 February 2012

Key indicators: single-crystal X-ray study; $T = 193$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.033; wR factor = 0.074; data-to-parameter ratio = 14.1.

In the title compound, $\text{C}_{22}\text{H}_{21}\text{NOS}$, the thiopyranone ring adopts a chair-like conformation with the substituent in the axial position. The relative configuration of the racemic compound is $3R,7S$ according to the numbering scheme used in this publication. In the crystal packing, centrosymmetric dimers are built up *via* $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, with graph set $R_2^2(8)$.

Related literature

For the preparation and spectroscopic characterization of the title compound and a series of related compounds, see: Abaee *et al.* (2012). For the crystal structure of *rac*-3-[(3-chloroanilino)(4-chlorophenyl)methyl]thian-4-one, see: Harms *et al.* (2012). For the crystal structures of related compounds, see: Guo *et al.* (2007); Fun *et al.* (2009). For patterns in hydrogen bonding, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{21}\text{NOS}$
 $M_r = 347.46$
 Monoclinic, $P2_1/n$
 $a = 10.8049$ (10) Å
 $b = 10.5497$ (15) Å
 $c = 16.4936$ (16) Å
 $\beta = 97.141$ (8)°
 $V = 1865.5$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.18$ mm⁻¹
 $T = 193$ K
 $0.45 \times 0.45 \times 0.36$ mm

Data collection

Stoe IPDS I diffractometer
 Absorption correction: integration
 [*X-AREA* and *X-RED32* (Stoe & Cie, 2006)]
 $T_{\min} = 0.942$, $T_{\max} = 0.960$
 13547 measured reflections
 3244 independent reflections
 1939 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.074$
 $S = 0.79$
 3244 reflections
 230 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.23$ e Å⁻³
 $\Delta\rho_{\min} = -0.37$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N8}-\text{H8}\cdots\text{O1}^i$	0.926 (15)	2.121 (16)	3.0450 (18)	175.4 (13)

 Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *EXPOSE* (Stoe & Cie, 1994); cell refinement: *CELL* (Stoe & Cie, 1994); data reduction: *X-RED32* (Stoe & Cie, 2006); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2007); software used to prepare material for publication: *publCIF* (Westrip, 2010), *PLATON* (Spek, 2009), and *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2509).

References

- Abaee, M. S., Mojtahedi, M. M., Akbari, A., Mehraki, E., Mesbah, A. W. & Harms, K. (2012). *J. Heterocycl. Chem.* **49**. In the press.
 Altomare, A., Casciarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
 Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Brandenburg, K. (2007). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Fun, H.-K., Chantrapromma, S., Rai, S., Shetty, P. & Isloor, A. M. (2009). *Acta Cryst.* **E65**, o539–o540.
 Guo, Q.-X., Liu, H., Guo, Ch., Luo, S.-W., Gu, Y. & Gong, L.-Z. (2007). *J. Am. Chem. Soc.* **129**, 3790–3791.
 Harms, K., Abaee, M. S., Mojtahedi, M. M. & Mesbah, A. W. (2012). *Acta Cryst.* **E68**, o646.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Stoe & Cie (1994). *IPDS User Manual*. Stoe & Cie, Darmstadt, Germany.
 Stoe & Cie (2006). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o749 [doi:10.1107/S1600536812005983]

***rac*-3-[(Anilino)(naphthalen-2-yl)methyl]thian-4-one**

Klaus Harms, M. Saeed Abaee, Mohammad M. Mojtahedi and A. Wahid Mesbah

S1. Experimental

The title compound is an example of a series of products of an *anti*-selective three-component Mannich reaction in the thiopyran-4-one system; see Abaee *et al.* (2012) for details. Colourless crystals suitable for crystal structure determination were grown from ethyl acetate.

S2. Refinement

Data have been merged using the program *X-RED32* (Stoe & Cie, 2006). Three beamstop affected reflections (1 1 0, -1 1 1, 0 1 1) have been excluded from the data during the refinement. All C bonded H atoms were placed in geometrical positions and constrained to ride on their parent atoms with C—H distances in the range 0.95–1.00 Å. The U_{iso} values were constrained to be $1.2U_{\text{eq}}$ of the parent C atom. The position of the N bonded H atom has been refined freely with an isotropic displacement factor. The N—H bond length is 0.926 (15) Å.




Figure 1

Molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at 50% probability level. H atoms are presented as small spheres of arbitrary radius. Dotted lines indicate hydrogen bonds to the neighbouring molecule generated by crystallographic inversion symmetry. For symmetry code (i), see Table 1.

***rac*-3-[(Anilino)(naphthalen-2-yl)methyl]thian-4-one**

Crystal data

$C_{22}H_{21}NO$
 $M_r = 347.46$
 Monoclinic, $P2_1/n$
 Hall symbol: $-P 2_1/n$
 $a = 10.8049$ (10) Å
 $b = 10.5497$ (15) Å
 $c = 16.4936$ (16) Å
 $\beta = 97.141$ (8)°
 $V = 1865.5$ (4) Å³
 $Z = 4$

$F(000) = 736$
 $D_x = 1.237$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 7999 reflections
 $\theta = 2.3$ – 26.0 °
 $\mu = 0.18$ mm⁻¹
 $T = 193$ K
 Nugget, colourless
 $0.45 \times 0.45 \times 0.36$ mm

Data collection

Stoe IPDS I
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator

Detector resolution: 6.67 pixels mm⁻¹
 rotation method scans
 Absorption correction: integration
 [*X-AREA* and *X-RED32* (Stoe & Cie, 2006)]

$T_{\min} = 0.942$, $T_{\max} = 0.960$
 13547 measured reflections
 3244 independent reflections
 1939 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

$\theta_{\max} = 25.4^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -13 \rightarrow 12$
 $k = -12 \rightarrow 12$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.074$
 $S = 0.79$
 3244 reflections
 230 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0399P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.37 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.06564 (5)	0.68462 (6)	0.53056 (4)	0.0733 (2)
O1	0.38884 (11)	0.55992 (11)	0.42256 (7)	0.0481 (3)
N8	0.48932 (11)	0.69693 (12)	0.59618 (8)	0.0318 (3)
H8	0.5282 (14)	0.6206 (15)	0.5880 (10)	0.033 (4)*
C2	0.16654 (16)	0.55096 (17)	0.55576 (14)	0.0568 (6)
H2A	0.1584	0.5237	0.6123	0.068*
H2B	0.1386	0.4798	0.5188	0.068*
C3	0.30417 (14)	0.57832 (15)	0.54910 (11)	0.0359 (4)
H3	0.3517	0.4977	0.5611	0.043*
C4	0.32089 (15)	0.61845 (15)	0.46296 (11)	0.0381 (4)
C5	0.24585 (19)	0.72936 (18)	0.42754 (13)	0.0579 (6)
H5A	0.2618	0.7424	0.3703	0.070*
H5B	0.2728	0.8067	0.4588	0.070*
C6	0.1050 (2)	0.7089 (2)	0.42969 (16)	0.0808 (8)
H6A	0.0776	0.6344	0.3956	0.097*
H6B	0.0591	0.7838	0.4055	0.097*
C7	0.36003 (13)	0.67883 (14)	0.61120 (10)	0.0301 (4)
H7	0.3141	0.7603	0.5991	0.036*
C9	0.56671 (14)	0.78247 (14)	0.64329 (10)	0.0306 (4)

C10	0.52229 (16)	0.89433 (15)	0.67309 (11)	0.0414 (4)
H10	0.4354	0.9118	0.6656	0.050*
C11	0.60450 (18)	0.98055 (17)	0.71379 (12)	0.0519 (5)
H11	0.5731	1.0574	0.7333	0.062*
C12	0.73045 (18)	0.95712 (18)	0.72650 (13)	0.0542 (5)
H12	0.7860	1.0172	0.7541	0.065*
C13	0.77470 (17)	0.84535 (18)	0.69860 (12)	0.0494 (5)
H13	0.8615	0.8275	0.7078	0.059*
C14	0.69458 (15)	0.75867 (16)	0.65734 (11)	0.0396 (4)
H14	0.7268	0.6819	0.6383	0.047*
C15	0.34684 (14)	0.64065 (13)	0.69838 (10)	0.0309 (4)
C16	0.25886 (15)	0.69614 (14)	0.74005 (11)	0.0362 (4)
H16	0.2072	0.7609	0.7141	0.043*
C17	0.24295 (15)	0.65944 (15)	0.82070 (11)	0.0358 (4)
C18	0.14989 (17)	0.71312 (17)	0.86377 (12)	0.0484 (5)
H18	0.0965	0.7772	0.8386	0.058*
C19	0.1364 (2)	0.67372 (18)	0.94068 (12)	0.0561 (6)
H19	0.0730	0.7099	0.9685	0.067*
C20	0.21464 (19)	0.58064 (17)	0.97926 (12)	0.0551 (6)
H20	0.2039	0.5538	1.0329	0.066*
C21	0.30610 (18)	0.52812 (16)	0.94044 (11)	0.0457 (5)
H21	0.3597	0.4660	0.9677	0.055*
C22	0.32231 (15)	0.56508 (14)	0.85973 (11)	0.0353 (4)
C23	0.41391 (15)	0.51076 (15)	0.81611 (11)	0.0385 (4)
H23	0.4686	0.4481	0.8417	0.046*
C24	0.42500 (14)	0.54655 (14)	0.73845 (11)	0.0354 (4)
H24	0.4866	0.5076	0.7104	0.042*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0349 (2)	0.0790 (4)	0.1041 (6)	−0.0052 (3)	0.0016 (3)	−0.0435 (4)
O1	0.0546 (7)	0.0537 (7)	0.0350 (8)	0.0127 (6)	0.0011 (7)	−0.0116 (6)
N8	0.0310 (7)	0.0330 (7)	0.0325 (9)	−0.0052 (6)	0.0084 (6)	−0.0056 (6)
C2	0.0460 (11)	0.0515 (11)	0.0727 (16)	−0.0199 (9)	0.0067 (11)	−0.0201 (10)
C3	0.0378 (9)	0.0311 (8)	0.0383 (12)	−0.0046 (7)	0.0032 (8)	−0.0072 (7)
C4	0.0410 (9)	0.0359 (9)	0.0348 (12)	−0.0004 (8)	−0.0054 (9)	−0.0129 (8)
C5	0.0824 (14)	0.0510 (11)	0.0378 (13)	0.0241 (10)	−0.0030 (11)	−0.0043 (9)
C6	0.0664 (14)	0.0796 (16)	0.085 (2)	0.0312 (12)	−0.0363 (13)	−0.0339 (14)
C7	0.0305 (8)	0.0311 (8)	0.0293 (10)	−0.0030 (7)	0.0062 (7)	−0.0028 (7)
C9	0.0353 (8)	0.0343 (8)	0.0222 (10)	−0.0076 (7)	0.0031 (8)	0.0042 (7)
C10	0.0392 (9)	0.0390 (9)	0.0449 (13)	−0.0053 (8)	0.0005 (9)	−0.0058 (8)
C11	0.0577 (12)	0.0432 (10)	0.0518 (15)	−0.0085 (9)	−0.0048 (11)	−0.0117 (9)
C12	0.0549 (12)	0.0532 (12)	0.0490 (15)	−0.0191 (10)	−0.0154 (10)	−0.0005 (10)
C13	0.0380 (9)	0.0556 (12)	0.0508 (14)	−0.0110 (9)	−0.0100 (9)	0.0116 (10)
C14	0.0399 (9)	0.0402 (9)	0.0379 (12)	−0.0031 (8)	0.0022 (9)	0.0073 (8)
C15	0.0303 (8)	0.0294 (8)	0.0337 (11)	−0.0061 (7)	0.0067 (8)	−0.0040 (7)
C16	0.0391 (8)	0.0338 (9)	0.0364 (12)	0.0022 (7)	0.0077 (8)	−0.0008 (8)

C17	0.0424 (9)	0.0331 (9)	0.0336 (11)	-0.0021 (7)	0.0118 (9)	-0.0048 (7)
C18	0.0591 (11)	0.0453 (10)	0.0440 (13)	0.0128 (9)	0.0188 (10)	-0.0016 (9)
C19	0.0763 (14)	0.0537 (11)	0.0440 (14)	0.0113 (11)	0.0299 (12)	-0.0060 (10)
C20	0.0856 (15)	0.0481 (11)	0.0354 (12)	0.0026 (11)	0.0228 (12)	-0.0021 (9)
C21	0.0627 (12)	0.0381 (9)	0.0371 (12)	0.0003 (9)	0.0097 (10)	-0.0001 (8)
C22	0.0419 (9)	0.0314 (8)	0.0336 (11)	-0.0044 (7)	0.0082 (8)	-0.0027 (7)
C23	0.0382 (9)	0.0347 (9)	0.0432 (13)	0.0030 (7)	0.0078 (9)	0.0023 (8)
C24	0.0325 (8)	0.0356 (9)	0.0398 (12)	0.0012 (7)	0.0114 (8)	0.0000 (8)

Geometric parameters (Å, °)

S1—C6	1.786 (3)	C11—H11	0.9500
S1—C2	1.799 (2)	C12—C13	1.373 (3)
O1—C4	1.2190 (18)	C12—H12	0.9500
N8—C9	1.398 (2)	C13—C14	1.379 (2)
N8—C7	1.4614 (18)	C13—H13	0.9500
N8—H8	0.926 (15)	C14—H14	0.9500
C2—C3	1.532 (2)	C15—C16	1.372 (2)
C2—H2A	0.9900	C15—C24	1.413 (2)
C2—H2B	0.9900	C16—C17	1.416 (2)
C3—C4	1.515 (2)	C16—H16	0.9500
C3—C7	1.544 (2)	C17—C22	1.415 (2)
C3—H3	1.0000	C17—C18	1.419 (2)
C4—C5	1.500 (2)	C18—C19	1.360 (3)
C5—C6	1.542 (3)	C18—H18	0.9500
C5—H5A	0.9900	C19—C20	1.397 (3)
C5—H5B	0.9900	C19—H19	0.9500
C6—H6A	0.9900	C20—C21	1.361 (2)
C6—H6B	0.9900	C20—H20	0.9500
C7—C15	1.517 (2)	C21—C22	1.419 (2)
C7—H7	1.0000	C21—H21	0.9500
C9—C10	1.387 (2)	C22—C23	1.415 (2)
C9—C14	1.395 (2)	C23—C24	1.355 (2)
C10—C11	1.385 (2)	C23—H23	0.9500
C10—H10	0.9500	C24—H24	0.9500
C11—C12	1.373 (3)		
C6—S1—C2	96.89 (10)	C12—C11—C10	121.29 (18)
C9—N8—C7	120.55 (12)	C12—C11—H11	119.4
C9—N8—H8	113.1 (10)	C10—C11—H11	119.4
C7—N8—H8	111.8 (9)	C13—C12—C11	118.89 (17)
C3—C2—S1	113.67 (13)	C13—C12—H12	120.6
C3—C2—H2A	108.8	C11—C12—H12	120.6
S1—C2—H2A	108.8	C12—C13—C14	120.76 (17)
C3—C2—H2B	108.8	C12—C13—H13	119.6
S1—C2—H2B	108.8	C14—C13—H13	119.6
H2A—C2—H2B	107.7	C13—C14—C9	120.70 (16)
C4—C3—C2	110.49 (16)	C13—C14—H14	119.7

C4—C3—C7	110.36 (12)	C9—C14—H14	119.7
C2—C3—C7	112.62 (13)	C16—C15—C24	118.46 (15)
C4—C3—H3	107.7	C16—C15—C7	120.95 (14)
C2—C3—H3	107.7	C24—C15—C7	120.59 (13)
C7—C3—H3	107.7	C15—C16—C17	121.78 (15)
O1—C4—C5	121.02 (17)	C15—C16—H16	119.1
O1—C4—C3	121.42 (15)	C17—C16—H16	119.1
C5—C4—C3	117.49 (15)	C22—C17—C16	118.82 (14)
C4—C5—C6	111.66 (17)	C22—C17—C18	118.67 (16)
C4—C5—H5A	109.3	C16—C17—C18	122.51 (16)
C6—C5—H5A	109.3	C19—C18—C17	120.67 (17)
C4—C5—H5B	109.3	C19—C18—H18	119.7
C6—C5—H5B	109.3	C17—C18—H18	119.7
H5A—C5—H5B	107.9	C18—C19—C20	120.71 (17)
C5—C6—S1	113.05 (16)	C18—C19—H19	119.6
C5—C6—H6A	109.0	C20—C19—H19	119.6
S1—C6—H6A	109.0	C21—C20—C19	120.40 (18)
C5—C6—H6B	109.0	C21—C20—H20	119.8
S1—C6—H6B	109.0	C19—C20—H20	119.8
H6A—C6—H6B	107.8	C20—C21—C22	120.76 (18)
N8—C7—C15	113.57 (13)	C20—C21—H21	119.6
N8—C7—C3	106.28 (12)	C22—C21—H21	119.6
C15—C7—C3	111.80 (12)	C23—C22—C17	118.42 (15)
N8—C7—H7	108.3	C23—C22—C21	122.78 (16)
C15—C7—H7	108.3	C17—C22—C21	118.79 (15)
C3—C7—H7	108.3	C24—C23—C22	121.16 (15)
C10—C9—C14	118.27 (15)	C24—C23—H23	119.4
C10—C9—N8	122.46 (14)	C22—C23—H23	119.4
C14—C9—N8	119.17 (14)	C23—C24—C15	121.34 (14)
C11—C10—C9	120.08 (16)	C23—C24—H24	119.3
C11—C10—H10	120.0	C15—C24—H24	119.3
C9—C10—H10	120.0		
C6—S1—C2—C3	57.57 (16)	N8—C9—C14—C13	175.40 (15)
S1—C2—C3—C4	-59.40 (17)	N8—C7—C15—C16	135.46 (15)
S1—C2—C3—C7	64.52 (19)	C3—C7—C15—C16	-104.28 (16)
C2—C3—C4—O1	-121.46 (17)	N8—C7—C15—C24	-44.72 (19)
C7—C3—C4—O1	113.33 (16)	C3—C7—C15—C24	75.54 (18)
C2—C3—C4—C5	55.62 (19)	C24—C15—C16—C17	-1.5 (2)
C7—C3—C4—C5	-69.58 (19)	C7—C15—C16—C17	178.33 (14)
O1—C4—C5—C6	121.38 (19)	C15—C16—C17—C22	1.6 (2)
C3—C4—C5—C6	-55.7 (2)	C15—C16—C17—C18	-178.03 (16)
C4—C5—C6—S1	59.1 (2)	C22—C17—C18—C19	-0.7 (3)
C2—S1—C6—C5	-56.56 (16)	C16—C17—C18—C19	178.91 (18)
C9—N8—C7—C15	-56.27 (19)	C17—C18—C19—C20	0.7 (3)
C9—N8—C7—C3	-179.60 (14)	C18—C19—C20—C21	0.2 (3)
C4—C3—C7—N8	-54.62 (17)	C19—C20—C21—C22	-1.1 (3)
C2—C3—C7—N8	-178.61 (15)	C16—C17—C22—C23	-0.4 (2)

C4—C3—C7—C15	-179.05 (13)	C18—C17—C22—C23	179.22 (15)
C2—C3—C7—C15	56.96 (19)	C16—C17—C22—C21	-179.83 (15)
C7—N8—C9—C10	-34.1 (2)	C18—C17—C22—C21	-0.2 (2)
C7—N8—C9—C14	149.59 (15)	C20—C21—C22—C23	-178.28 (17)
C14—C9—C10—C11	1.6 (2)	C20—C21—C22—C17	1.1 (3)
N8—C9—C10—C11	-174.77 (16)	C17—C22—C23—C24	-0.8 (2)
C9—C10—C11—C12	-0.8 (3)	C21—C22—C23—C24	178.61 (17)
C10—C11—C12—C13	-0.5 (3)	C22—C23—C24—C15	0.9 (2)
C11—C12—C13—C14	1.0 (3)	C16—C15—C24—C23	0.3 (2)
C12—C13—C14—C9	-0.2 (3)	C7—C15—C24—C23	-179.57 (15)
C10—C9—C14—C13	-1.1 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N8—H8 \cdots O1 ⁱ	0.926 (15)	2.121 (16)	3.0450 (18)	175.4 (13)

Symmetry code: (i) $-x+1, -y+1, -z+1$.