

## N'-[3-(Hydroxyimino)butan-2-ylidene]-4-methylbenzene-1-sulfonohydrazide

Maria C. S. Bulhosa,<sup>a</sup> Vanessa Carratu Gervini,<sup>a\*</sup> Leandro Bresolin,<sup>a</sup> Aline Locatelli<sup>b</sup> and Adriano Bof de Oliveira<sup>c</sup>

<sup>a</sup>Escola de Química e Alimentos, Universidade Federal do Rio Grande, Av. Itália km 08, Campus Carreiros, 96201-900, Rio Grande, RS, Brazil, <sup>b</sup>Departamento de Química, Universidade Federal de Santa Maria, Av. Roraima, Campus, 97105-900, Santa Maria, RS, Brazil, and <sup>c</sup>Departamento de Química, Universidade Federal de Sergipe, Av. Marechal Rondon s/n, Campus, 49100-000, São Cristóvão, SE, Brazil  
Correspondence e-mail: adriano@daad-alumni.de


Received 17 January 2012; accepted 26 January 2012

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.036;  $wR$  factor = 0.112; data-to-parameter ratio = 18.5.

In the title compound,  $\text{C}_{11}\text{H}_{15}\text{N}_3\text{O}_3\text{S}$ , the  $\text{C}-\text{S}-\text{N}(\text{H})-\text{N}$  linkage is nonplanar, the torsion angle being  $75.70(12)^\circ$ . The compound has two almost planar fragments linked to the S atom: the hydrazone-derivative fragment  $[(\text{HONC}_4\text{H}_6)\text{N}-\text{N}(\text{H})-]$  and the tolyl fragment ( $\text{C}_7\text{H}_7-$ ) have maximum deviations from the mean plane through the non-H atoms of  $0.0260(10)$  and  $0.0148(14)\text{ \AA}$ , respectively. The two planar fragments make an interplanar angle of  $79.47(5)^\circ$ . In the crystal, molecules are connected through inversion centers *via* pairs of  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bonds.

### Related literature

For the synthesis and application of hydroxyimino-tosylhydrazones as complexing agents, see: Beger *et al.* (1991). For a similar structure with a tosylhydrazone derivative, see: Fonseca *et al.* (2011).



### Experimental

#### Crystal data

$\text{C}_{11}\text{H}_{15}\text{N}_3\text{O}_3\text{S}$	$\gamma = 87.688(1)^\circ$
$M_r = 269.32$	$V = 643.11(2)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 5.5740(1)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.4354(2)\text{ \AA}$	$\mu = 0.26\text{ mm}^{-1}$
$c = 11.3997(2)\text{ \AA}$	$T = 293\text{ K}$
$\alpha = 83.586(1)^\circ$	$0.55 \times 0.24 \times 0.22\text{ mm}$
$\beta = 77.453(1)^\circ$	

#### Data collection

Bruker APEXII CCD	11000 measured reflections
diffractometer	3211 independent reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2005)	2862 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.872$ , $T_{\max} = 0.946$	$R_{\text{int}} = 0.015$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.112$	$\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$
3211 reflections	
174 parameters	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}3-\text{H}8\cdots\text{O}3^{\text{i}}$	0.82 (2)	2.19 (2)	2.9830 (18)	165.0 (19)
$\text{O}1-\text{H}1\cdots\text{N}1^{\text{ii}}$	0.84 (3)	1.99 (3)	2.792 (2)	160 (2)

Symmetry codes: (i)  $-x + 2, -y, -z$ ; (ii)  $-x, -y + 1, -z$ .

Data collection: *COSMO* (Bruker, 2005); cell refinement: *SAINT*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors gratefully acknowledge Professor Dr Manfredo Hörner (Department of Chemistry, Federal University of Santa Maria, Brazil) for his help and support with the X-ray measurements, and CNPq/FAPERGS for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FY2042).

### References

- Beger, J., Siedler, F., Mühl, P. & Gloe, K. (1991). German Patent DD287027A5.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2005). *COSMO*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fonseca, A. de S., Storino, T. G., Carratu, V. S., Locatelli, A. & de Oliveira, A. B. (2011). *Acta Cryst. E*67, o3256.
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst. A*43, 920–925.

# supporting information

*Acta Cryst.* (2012). E68, o592 [doi:10.1107/S1600536812003339]

## N'-[3-(Hydroxyimino)butan-2-ylidene]-4-methylbenzene-1-sulfonohydrazide

**Maria C. S. Bulhosa, Vanessa Carratu Gervini, Leandro Bresolin, Aline Locatelli and Adriano Bof de Oliveira**

### S1. Comment


Hydrazones have a wide range of applications on inorganic chemistry. For example, sulfonylhydrazones are used as complexing agents for cobalt (II) (Beger *et al.*, 1991). As part of our study of sulfonylhydrazone derivatives, we report herein the crystal structure of an oxime-sulfonylhydrazone derivative. In the title compound (Fig. 1) the C—S—N(H)—N linkage is non-planar with the torsion angle being 75.70 (12) $^{\circ}$  and a tetrahedral environment suggests a  $sp^3$  hybridization for the S atom. The title structure contains additionally two planar fragments. The maximum deviations from the least squares planes for the hydrazone derivative fragment C1/C2/C3/C4/N1/N2/N3/O1 and for the tolyl fragment C5/C6/C7/C8/C9/C10/C11 amount to 0.0260 (10) $^{\circ}$  and for N3 and 0.0148 (14) $^{\circ}$  for C9 atoms, respectively. The dihedral angle between the two planes is 79.47 (5) $^{\circ}$ . The crystal packing is stabilized by intermolecular N—H···O (Table 1; N3—H8···O3<sup>i</sup>) and O—H···N bonds (Table 1; O1—H1···N1<sup>ii</sup>) connecting the molecules through inversion centers (Fig. 2). Symmetry codes: (i)-x+2, -y, -z; (ii)-x, -y+1, -z.

### S2. Experimental


Starting materials were commercially available and were used without further purification. The synthesis was adapted from a procedure reported previously (Beger *et al.*, 1991). The reaction of 2,3-Butanedione monoxime (5 mmol) and *p*-toluenesulfonylhydrazine (5 mmol) in ethanol (50 ml) was refluxed for 3 h. After cooling and filtering, crystals suitable for X-ray diffraction were obtained.

### S3. Refinement

H atoms attached to C atoms were positioned with idealized geometry and were refined isotropically with  $U_{\text{iso}}(\text{H})$  set to 1.2 times  $U_{\text{eq}}(\text{C})$  for the aromatic and 1.5 times  $U_{\text{eq}}(\text{C})$  for methyl H atoms using a riding model with C—H = 0.93 Å and C—H = 0.96 Å, respectively. H atoms attached to N and O atoms were located in difference Fourier maps and included in the subsequent refinement using restraints (N3—H8 = 0.82 (2) Å and O1—H1 = 0.84 (3) Å) with  $U_{\text{iso}}(\text{H})$  = 1.5 times of the  $U_{\text{eq}}(\text{N})$  and  $U_{\text{eq}}(\text{O})$ , respectively. In the last stage of refinement, they were refined freely.

**Figure 1**

The molecular structure of the title compound with labeling and displacement ellipsoids drawn at the 40% probability level.

**Figure 2**

Molecules of the title compound connected through inversion centers *via* pairs of N—H···O and O—H···N hydrogen bonds in the crystal structure. Intermolecular hydrogen bonding is indicated as dashed lines. Symmetry codes: (i)-x+2, -y, -z; (ii)-x, -y+1, -z.

### *N'*-[3-(Hydroxyimino)butan-2-ylidene]-4-methylbenzene-1-sulfonohydrazide

#### Crystal data

C<sub>11</sub>H<sub>15</sub>N<sub>3</sub>O<sub>3</sub>S  
 $M_r = 269.32$   
Triclinic,  $P\bar{1}$   
Hall symbol: -P 1  
 $a = 5.5740 (1)$  Å  
 $b = 10.4354 (2)$  Å  
 $c = 11.3997 (2)$  Å  
 $\alpha = 83.586 (1)$ °  
 $\beta = 77.453 (1)$ °

$\gamma = 87.688 (1)$ °  
 $V = 643.11 (2)$  Å<sup>3</sup>  
 $Z = 2$   
 $F(000) = 284$   
 $D_x = 1.391$  Mg m<sup>-3</sup>  
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 6049 reflections  
 $\theta = 2.6\text{--}28.3$ °  
 $\mu = 0.26$  mm<sup>-1</sup>

$T = 293\text{ K}$   
Block, colourless

#### Data collection

Bruker APEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube, Bruker  
X8 APEXII  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2005)  
 $T_{\min} = 0.872$ ,  $T_{\max} = 0.946$

$0.55 \times 0.24 \times 0.22\text{ mm}$

11000 measured reflections  
3211 independent reflections  
2862 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.015$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 1.8^\circ$   
 $h = -7 \rightarrow 6$   
 $k = -13 \rightarrow 13$   
 $l = -15 \rightarrow 15$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.112$   
 $S = 1.05$   
3211 reflections  
174 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_{\text{o}}^2) + (0.0651P)^2 + 0.1526P]$   
where  $P = (F_{\text{o}}^2 + 2F_{\text{c}}^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.27\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.94931 (6)	0.06163 (3)	0.19318 (3)	0.03670 (12)
O3	1.14412 (19)	-0.00383 (11)	0.11725 (10)	0.0465 (3)
O1	0.0184 (2)	0.53370 (13)	0.13328 (14)	0.0599 (3)
O2	1.00547 (19)	0.14436 (11)	0.27426 (10)	0.0474 (3)
N3	0.8101 (2)	0.14880 (12)	0.09796 (12)	0.0410 (3)
N2	0.6390 (2)	0.23836 (11)	0.14677 (11)	0.0383 (3)
N1	0.1759 (2)	0.44076 (12)	0.07432 (12)	0.0446 (3)
C3	0.4937 (2)	0.28939 (12)	0.08064 (12)	0.0361 (3)
C5	0.7414 (2)	-0.05473 (13)	0.27630 (13)	0.0377 (3)
C2	0.3235 (2)	0.38812 (13)	0.13806 (13)	0.0393 (3)
C8	0.4212 (3)	-0.23753 (16)	0.41661 (14)	0.0489 (4)
C6	0.8088 (3)	-0.18421 (15)	0.28051 (16)	0.0493 (4)

H9	0.9602	-0.2102	0.2366	0.059*
C9	0.3564 (3)	-0.10782 (18)	0.40921 (16)	0.0546 (4)
H11	0.2030	-0.0821	0.4513	0.066*
C10	0.5143 (3)	-0.01583 (16)	0.34077 (16)	0.0502 (4)
H12	0.4691	0.0710	0.3379	0.060*
C4	0.4871 (3)	0.25726 (17)	-0.04261 (15)	0.0496 (4)
H5	0.4554	0.1671	-0.0395	0.074*
H6	0.6424	0.2770	-0.0962	0.074*
H7	0.3591	0.3070	-0.0716	0.074*
C7	0.6482 (3)	-0.27425 (16)	0.35074 (17)	0.0554 (4)
H10	0.6934	-0.3611	0.3539	0.066*
C11	0.2476 (4)	-0.3351 (2)	0.49484 (18)	0.0665 (5)
H13	0.0895	-0.3248	0.4741	0.100*
H14	0.2317	-0.3220	0.5782	0.100*
H15	0.3110	-0.4205	0.4820	0.100*
C1	0.3317 (3)	0.41963 (17)	0.26113 (15)	0.0539 (4)
H2	0.1975	0.4771	0.2887	0.081*
H3	0.4844	0.4604	0.2586	0.081*
H4	0.3186	0.3418	0.3155	0.081*
H8	0.794 (4)	0.1114 (19)	0.0409 (19)	0.053 (5)*
H1	-0.061 (5)	0.556 (2)	0.079 (2)	0.082 (8)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.03284 (18)	0.03859 (19)	0.0401 (2)	0.00890 (12)	-0.01154 (13)	-0.00617 (13)
O3	0.0385 (5)	0.0504 (6)	0.0492 (6)	0.0159 (4)	-0.0088 (4)	-0.0071 (5)
O1	0.0532 (7)	0.0575 (7)	0.0732 (8)	0.0290 (6)	-0.0195 (6)	-0.0236 (6)
O2	0.0453 (5)	0.0482 (6)	0.0538 (6)	0.0033 (4)	-0.0191 (5)	-0.0124 (5)
N3	0.0415 (6)	0.0419 (6)	0.0407 (6)	0.0153 (5)	-0.0128 (5)	-0.0067 (5)
N2	0.0352 (5)	0.0364 (6)	0.0425 (6)	0.0083 (4)	-0.0080 (5)	-0.0040 (5)
N1	0.0374 (6)	0.0396 (6)	0.0567 (8)	0.0137 (5)	-0.0106 (5)	-0.0093 (5)
C3	0.0323 (6)	0.0333 (6)	0.0412 (7)	0.0042 (5)	-0.0067 (5)	-0.0015 (5)
C5	0.0373 (6)	0.0396 (7)	0.0382 (7)	0.0059 (5)	-0.0133 (5)	-0.0053 (5)
C2	0.0347 (6)	0.0357 (6)	0.0461 (7)	0.0045 (5)	-0.0064 (5)	-0.0045 (6)
C8	0.0541 (8)	0.0520 (9)	0.0427 (8)	-0.0057 (7)	-0.0148 (6)	-0.0042 (7)
C6	0.0467 (8)	0.0431 (8)	0.0566 (9)	0.0095 (6)	-0.0073 (7)	-0.0096 (7)
C9	0.0450 (8)	0.0599 (10)	0.0548 (9)	0.0056 (7)	-0.0030 (7)	-0.0059 (8)
C10	0.0460 (8)	0.0444 (8)	0.0568 (9)	0.0110 (6)	-0.0060 (7)	-0.0041 (7)
C4	0.0522 (8)	0.0521 (9)	0.0457 (8)	0.0189 (7)	-0.0145 (7)	-0.0094 (7)
C7	0.0620 (10)	0.0392 (8)	0.0640 (10)	0.0038 (7)	-0.0123 (8)	-0.0053 (7)
C11	0.0719 (12)	0.0650 (11)	0.0590 (11)	-0.0142 (9)	-0.0079 (9)	0.0007 (9)
C1	0.0585 (9)	0.0545 (9)	0.0503 (9)	0.0140 (7)	-0.0129 (7)	-0.0153 (7)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

S1—O2	1.4225 (11)	C8—C11	1.506 (2)
S1—O3	1.4349 (10)	C6—C7	1.383 (2)

S1—N3	1.6420 (12)	C6—H9	0.9300
S1—C5	1.7591 (14)	C9—C10	1.381 (2)
O1—N1	1.4084 (16)	C9—H11	0.9300
O1—H1	0.84 (3)	C10—H12	0.9300
N3—N2	1.3807 (16)	C4—H5	0.9600
N3—H8	0.82 (2)	C4—H6	0.9600
N2—C3	1.2843 (18)	C4—H7	0.9600
N1—C2	1.2808 (19)	C7—H10	0.9300
C3—C2	1.4806 (18)	C11—H13	0.9600
C3—C4	1.489 (2)	C11—H14	0.9600
C5—C6	1.386 (2)	C11—H15	0.9600
C5—C10	1.389 (2)	C1—H2	0.9600
C2—C1	1.487 (2)	C1—H3	0.9600
C8—C9	1.385 (2)	C1—H4	0.9600
C8—C7	1.387 (2)		
O2—S1—O3	119.84 (7)	C10—C9—C8	121.49 (15)
O2—S1—N3	107.87 (7)	C10—C9—H11	119.3
O3—S1—N3	104.20 (6)	C8—C9—H11	119.3
O2—S1—C5	108.48 (7)	C9—C10—C5	119.14 (15)
O3—S1—C5	108.09 (7)	C9—C10—H12	120.4
N3—S1—C5	107.77 (6)	C5—C10—H12	120.4
N1—O1—H1	98.5 (17)	C3—C4—H5	109.5
N2—N3—S1	116.06 (10)	C3—C4—H6	109.5
N2—N3—H8	122.1 (14)	H5—C4—H6	109.5
S1—N3—H8	113.8 (14)	C3—C4—H7	109.5
C3—N2—N3	117.27 (12)	H5—C4—H7	109.5
C2—N1—O1	112.69 (13)	H6—C4—H7	109.5
N2—C3—C2	113.63 (13)	C6—C7—C8	121.31 (15)
N2—C3—C4	125.81 (13)	C6—C7—H10	119.3
C2—C3—C4	120.56 (12)	C8—C7—H10	119.3
C6—C5—C10	120.45 (14)	C8—C11—H13	109.5
C6—C5—S1	119.79 (11)	C8—C11—H14	109.5
C10—C5—S1	119.72 (11)	H13—C11—H14	109.5
N1—C2—C3	115.11 (13)	C8—C11—H15	109.5
N1—C2—C1	124.73 (13)	H13—C11—H15	109.5
C3—C2—C1	120.15 (12)	H14—C11—H15	109.5
C9—C8—C7	118.35 (15)	C2—C1—H2	109.5
C9—C8—C11	120.18 (16)	C2—C1—H3	109.5
C7—C8—C11	121.47 (16)	H2—C1—H3	109.5
C7—C6—C5	119.24 (15)	C2—C1—H4	109.5
C7—C6—H9	120.4	H2—C1—H4	109.5
C5—C6—H9	120.4	H3—C1—H4	109.5
O2—S1—N3—N2	-41.25 (12)	N2—C3—C2—N1	-179.95 (12)
O3—S1—N3—N2	-169.63 (10)	C4—C3—C2—N1	0.2 (2)
C5—S1—N3—N2	75.70 (12)	N2—C3—C2—C1	0.0 (2)
S1—N3—N2—C3	-166.53 (10)	C4—C3—C2—C1	-179.93 (15)

N3—N2—C3—C2	−177.60 (11)	C10—C5—C6—C7	−0.7 (2)
N3—N2—C3—C4	2.3 (2)	S1—C5—C6—C7	176.99 (13)
O2—S1—C5—C6	−117.69 (13)	C7—C8—C9—C10	−1.6 (3)
O3—S1—C5—C6	13.69 (15)	C11—C8—C9—C10	178.28 (17)
N3—S1—C5—C6	125.76 (13)	C8—C9—C10—C5	1.1 (3)
O2—S1—C5—C10	60.01 (14)	C6—C5—C10—C9	0.0 (2)
O3—S1—C5—C10	−168.61 (12)	S1—C5—C10—C9	−177.65 (13)
N3—S1—C5—C10	−56.54 (14)	C5—C6—C7—C8	0.2 (3)
O1—N1—C2—C3	179.96 (12)	C9—C8—C7—C6	0.9 (3)
O1—N1—C2—C1	0.1 (2)	C11—C8—C7—C6	−178.96 (17)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N3—H8···O3 <sup>i</sup>	0.82 (2)	2.19 (2)	2.9830 (18)	165.0 (19)
O1—H1···N1 <sup>ii</sup>	0.84 (3)	1.99 (3)	2.792 (2)	160 (2)

Symmetry codes: (i)  $-x+2, -y, -z$ ; (ii)  $-x, -y+1, -z$ .