

Tris(tetrabutylammonium) hexakis(*tert*-butanethiolato- κ S)hepta- μ_3 -chlorido- μ_3 -sulfido-hexamolybdate dihydrate

Pavel A. Petrov,* Dmitry Yu. Naumov and Sergey N. Konchenko

Nikolaev Institute of Inorganic Chemistry, SB Russian Academy of Sciences, Akademian Lavrentiev prospekt 3, Novosibirsk 90, 630090, Russian Federation, and, Novosibirsk State University, Pirogov street 2, Novosibirsk 90, 630090, Russian Federation

Correspondence e-mail: panah@mail.ru


Received 1 February 2012; accepted 19 February 2012

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; H-atom completeness 98%; disorder in main residue; R factor = 0.038; wR factor = 0.125; data-to-parameter ratio = 22.6.

The octahedral cluster core of the anion in the structure of the title compound, $(\text{C}_{16}\text{H}_{36}\text{N})_3[\text{Mo}_6(\text{C}_4\text{H}_9\text{S})_6(\mu_3\text{-Cl})_7(\mu_3\text{-S})] \cdot 2\text{H}_2\text{O}$, has $\bar{3}$ site symmetry. Two $\mu_3\text{-Cl}$ atoms fully occupy positions in the cluster core, while the remaining six positions are statistically occupied by Cl and S atoms in a 1:5 ratio. The fully occupied Cl-atom positions are located on sites with 3 symmetry, and the N atom of tetrabutylammonium cation is located on a site with 2 symmetry. The structure contains also two disordered solvent water molecules, one of which is located on a threefold rotation axis and the other in a general position, both with an occupancy of 0.25. The water molecules are localized in cavities formed by the tetrabutylammonium cations and the *tert*-butanethiolate groups. The metal clusters are stacked in a cubic close packing arrangement along [001].

Related literature

For a review of octahedral halogen-bridged metal clusters, see: Prokopuk & Shryver (1998). For synthesis and structures of related halogen/chalcogen clusters, see: Abramov *et al.* (2009); Ebihara *et al.* (1988); Ebihara, Imai *et al.* (1995); Ebihara, Toriumi *et al.* (1995); Michel & McCarley (1982); Nocera & Gray (1984). For a related transformation of *t*BuS[−], see: Petrov *et al.* (2010). For synthesis and structures of related clusters with sulfur-substituted halogen atoms, see: Schoonover *et al.* (1996); Szczepura *et al.* (2008).



Experimental

Crystal data

$(\text{C}_{16}\text{H}_{36}\text{N})_3[\text{Mo}_6(\text{C}_4\text{H}_9\text{S})_6\text{Cl}_7\text{S}] \cdot 2\text{H}_2\text{O}$
 $M_r = 2154.29$
 Trigonal, $R\bar{3}c$
 $a = 18.7481$ (5) Å
 $c = 52.4233$ (12) Å
 $V = 15957.7$ (7) Å³
 $Z = 6$
 Mo $K\alpha$ radiation
 $\mu = 1.04$ mm^{−1}
 $T = 150$ K
 $0.42 \times 0.35 \times 0.23$ mm

Data collection

Bruker–Nonius X8 APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2004)
 $T_{\text{min}} = 0.670$, $T_{\text{max}} = 0.797$
 36925 measured reflections
 3637 independent reflections
 3092 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.125$
 $S = 1.14$
 3637 reflections
 161 parameters
 12 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.16$ e Å^{−3}
 $\Delta\rho_{\text{min}} = -0.76$ e Å^{−3}

Table 1

Selected bond lengths (Å).

Mo1—Mo1 ⁱ	2.6067 (4)	Mo1—Cl1	2.5054 (10)
Mo1—Mo1 ⁱⁱ	2.6328 (5)	Mo1—Cl2	2.4801 (9)
Mo1—S1	2.5158 (9)	Mo1—Cl2 ⁱⁱⁱ	2.4792 (9)
Mo1—S2 ⁱⁱⁱ	2.4792 (9)	Mo1—Cl2 ^{iv}	2.4842 (9)
Mo1—S2 ^{iv}	2.4842 (9)		

Symmetry codes: (i) $y, -x + y, -z$; (ii) $-x + y, -x, z$; (iii) $-y, x - y, z$; (iv) $x - y, x, -z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008), Mercury (Macrae *et al.*, 2006) and POV-RAY (Persistence of Vision, 2004); software used to prepare material for publication: SHELXTL.

The work was supported by RFBR (grants 09–03–38245a and 10–03–38245a) and State Contract GK-02.740.11.0628. The authors are grateful to Dr Cristian Vicent (Universitat Jaume I, Castello, Spain) for ES-MS data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2589).

References

- Abramov, P. A., Sokolov, M. N., Virovets, A. V., Peresyphina, E. V., Vicent, C. & Fedin, V. P. (2009). *J. Cluster Sci.* **20**, 83–92.
- Bruker (2004). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ebihara, M., Imai, T. & Kawamura, T. (1995). *Acta Cryst.* **C51**, 1743–1745.
- Ebihara, M., Toriumi, K. & Saito, K. (1988). *Inorg. Chem.* **27**, 13–18.
- Ebihara, M., Toriumi, K., Sasaki, Y. & Saito, K. (1995). *Gazz. Chim. Ital.* **125**, 87–95.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Michel, J. B. & McCarley, R. E. (1982). *Inorg. Chem.* **21**, 1864–1872.
- Nocera, D. G. & Gray, H. B. (1984). *J. Am. Chem. Soc.* **106**, 824–825.
- Persistence of Vision (2004). *POV-RAY* Persistence of Vision Pty. Ltd, Williamstown, Victoria, Australia.
- Petrov, P. A., Virovets, A. V., Alberola, A., Llusar, R. & Konchenko, S. N. (2010). *Dalton Trans.* **39**, 8875–8877.
- Prokopuk, N. & Shryver, D. F. (1998). *Adv. Inorg. Chem.*, Vol. 46, edited by A. G. Sykes, pp. 1–49. New York: Academic Press.
- Schoonover, J. R., Zietlow, T. C., Clark, D. L., Heppert, J. A., Chisholm, M. H., Gray, H. B., Sattelberger, A. P. & Woodruff, W. H. (1996). *Inorg. Chem.* **35**, 6606–6613.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Szczepura, L. F., Ketcham, K. A., Ooro, B. A., Edwards, J. A., Templeton, J. N., Cedeno, D. L. & Jiricitano, A. J. (2008). *Inorg. Chem.* **47**, 7271–7278.

supporting information

Acta Cryst. (2012). E68, m333–m334 [doi:10.1107/S1600536812007416]

Tris(tetrabutylammonium) hexakis(*tert*-butanethiolato- κ S)hepta- μ_3 -chlorido- μ_3 -sulfido-hexamolybdate dihydrate

Pavel A. Petrov, Dmitry Yu. Naumov and Sergey N. Konchenko

S1. Comment

The octahedral clusters of early transition metals are often regarded as precursors of functional materials with redox and/or luminescent properties. The advantage of halide-bridged clusters $[\text{Mo}_6(\mu_3\text{-X})_8\text{X}_6]^{2-}$ ($X = \text{halogen}$) is the ability of tuning the electronic structure and the properties of the cluster core by step-by-step exchange of the terminal X atoms. The exchange of the μ_3 -bridging X atoms is also possible; however, reactions of this type are less common. The halogen-chalcogen clusters with one or two chalcogen atoms (or their mixtures) were obtained in the reactions of $[\text{Mo}_6(\mu_3\text{-X})_8\text{X}_6]^{2-}$ with NaSH, NaSeH or *in situ*-generated H_2Se (Michel & McCarley, 1982; Ebihara *et al.*, 1988; Ebihara, Imai *et al.*, 1995; Ebihara, Toriumi *et al.*, 1995; Abramov *et al.*, 2009).

Recently, the reaction of $[\text{Mo}_6(\mu_3\text{-X})_8(\text{OMe})_6]^{2-}$ with excess EtSH was reported leading to the smooth substitution of the terminal methoxides to ethanethiolate groups. The latter can be further substituted by other SR groups where $R = \text{butyl}$, benzyl or 3-indolyl (Szczepura *et al.*, 2008). Our attempt was aimed to prove if the reaction of $[\text{Mo}_6(\mu_3\text{-X})_8\text{X}_6]^{2-}$ with tBuSNa would stop on the substitution of the terminal X atoms, or would result in a core rearrangement as well. Previously, tBuS^- was reported to be the source of the S^{2-} anion (see, for example: Petrov *et al.* 2010).

The presence of three tetrabutylammonium cations designates the charge of the cluster core. Keeping in mind the high oxidation potential of the $[\text{Mo}_6(\mu_3\text{-Cl})_8\text{Cl}_6]^{3-/2-}$ pair (1.53 V in MeCN *versus* SCE, Nocera & Gray, 1984) one would formulate the cluster core composition as $[\text{Mo}_6(\mu_3\text{-S})(\mu_3\text{-Cl})_7(\text{S}'\text{Bu})_6]^{3-}$. The analysis of the temperature factors of the atoms in the μ_3 -positions leads us to the conclusion that two positions are occupied with Cl atoms only, while the remaining six positions are statistically occupied with Cl and S atoms in a 1:5 ratio (Fig. 1). The presence of one S atom in the cluster core has no noticeable effect on its geometry (Schoonover *et al.*, 1996; Szczepura *et al.*, 2008).

The structure contains two disordered lattice water molecules. One is located on a threefold rotation axis, the other is located in a general position. Both have an occupancy of 0.25 and are disordered over a site with symmetry 32. These two water molecules have an $\text{O} \cdots \text{O}$ distance of 2.706 (8) Å, pointing to hydrogen-bonding interactions. The water molecules are localized in cavities formed by the Bu_4N^+ cations and tBuS groups. The water incorporated in the structure most likely originated from the starting material $(\text{Bu}_4\text{N})_2[\text{Mo}_6(\mu_3\text{-Cl})_8\text{Cl}_6]n\text{H}_2\text{O}$.

The centres of the metal clusters are arranged in a cubic close packing along [001] as stacking direction (Fig. 2).

S2. Experimental

A mixture of 128.8 mg (0.083 mmol) $(\text{Bu}_4\text{N})_2[\text{Mo}_6\text{Cl}_{14}]$ and 154.8 mg (1.38 mmol) $\text{NaS}'\text{Bu}$ (1:16.7 molar ratio) in 15 ml CH_3CN was refluxed for 5 days. The resulting brown solution was filtered to remove the white residue and left standing at 278 K. After several weeks almost black crystals were formed. The largest positive and negative residual electron densities are located 0.73 Å from atom OW3 and 0.58 Å from atom S2, respectively.

S3. Refinement

The site occupation factors of the S and Cl atoms of the disordered Cl2/S2 site were preliminary refined without any constraints giving us the ratio. Constrained occupation factors were taken into account in the final refinement cycle. The composition of the anion has been confirmed by electrospray mass-spectrometry. The signal at m/z 695.7 was assigned to the $[\text{Mo}_6(\mu_3\text{-S})(\mu_3\text{-Cl})_7(\text{S}^t\text{Bu})_6]^{2-}$ anion. The H atoms of the disordered water molecules could not be located and were excluded from refinement.




Figure 1

Molecular structure of $[\text{Mo}_6(\mu_3\text{-S})(\mu_3\text{-Cl})_7(\text{S}^t\text{Bu})_6]^{3-}$ anion. Displacement ellipsoids are plotted at the 50% probability level.




Figure 2

Packing of the structure viewed along *c* axis. O and H atoms are omitted for clarity.

Tris(tetrabutylammonium) hexakis(*tert*-butanethiolato- κ S)hepta- μ_3 -chlorido- μ_3 -sulfido-hexamolybdate dihydrate

Crystal data

(C₁₆H₃₆N)₃[Mo₆(C₄H₉S)₆Cl₇S]·2H₂O

M_r = 2154.29

Trigonal, *R* $\bar{3}c$

Hall symbol: -R 3 2" c

a = 18.7481 (5) Å

c = 52.4233 (12) Å

V = 15957.7 (7) Å³

Z = 6

F(000) = 6708

D_x = 1.345 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 9844 reflections

θ = 2.5–28.3°

μ = 1.04 mm⁻¹

T = 150 K

Prism, brown

0.42 × 0.35 × 0.23 mm

Data collection

Bruker–Nonius X8 APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 25 pixels mm⁻¹

φ scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2004)

T_{min} = 0.670, *T_{max}* = 0.797

36925 measured reflections

3637 independent reflections

3092 reflections with *I* > 2σ(*I*)

R_{int} = 0.034

θ_{max} = 26.4°, θ_{min} = 2.2°

h = -23→23

k = -23→22

l = -65→55

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.038$ $wR(F^2) = 0.125$ $S = 1.14$

3637 reflections

161 parameters

12 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0654P)^2 + 69.7786P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 1.16 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.76 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. Hydrogen atoms of water molecules are not located. One of water molecules is disordered by two positions. Hydrogen atoms of cation and anion are placed geometrically.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Mo1	0.093280 (16)	0.039723 (17)	0.020197 (5)	0.02512 (13)	
Cl1	0.0000	0.0000	0.05819 (2)	0.0298 (3)	
Cl2	0.07472 (5)	-0.10085 (5)	0.019692 (16)	0.0374 (2)	0.8333333
S1	0.21632 (6)	0.09894 (7)	0.049357 (18)	0.0458 (3)	
S2	0.07472 (5)	-0.10085 (5)	0.019692 (16)	0.0374 (2)	0.1666667
O2W	0.5911 (10)	0.3102 (13)	0.0925 (4)	0.103 (6)	0.25
N1	0.2617 (3)	0.3333	0.0833	0.0555 (12)	
C1	0.3076 (2)	0.0927 (3)	0.03926 (8)	0.0478 (9)	
C2	0.3649 (4)	0.1194 (6)	0.06188 (11)	0.107 (3)	
H2A	0.4128	0.1137	0.0578	0.161*	
H2B	0.3359	0.0848	0.0766	0.161*	
H2C	0.3831	0.1771	0.0659	0.161*	
C3	0.2844 (4)	0.0089 (4)	0.02944 (17)	0.111 (3)	
H3A	0.2454	-0.0054	0.0153	0.167*	
H3B	0.2587	-0.0316	0.0432	0.167*	
H3C	0.3339	0.0089	0.0234	0.167*	
C4	0.3539 (4)	0.1528 (5)	0.01793 (13)	0.099 (2)	
H4A	0.3731	0.2091	0.0238	0.148*	
H4B	0.3172	0.1408	0.0033	0.148*	
H4C	0.4013	0.1473	0.0129	0.148*	
C11	0.3094 (3)	0.3354 (3)	0.10717 (8)	0.0559 (11)	
H11A	0.3620	0.3883	0.1075	0.067*	
H11B	0.2773	0.3342	0.1223	0.067*	

C12	0.3281 (3)	0.2659 (3)	0.10940 (9)	0.0669 (13)	
H12A	0.2761	0.2126	0.1078	0.080*	
H12B	0.3645	0.2697	0.0952	0.080*	
C13	0.3688 (4)	0.2676 (3)	0.13419 (10)	0.0809 (16)	
H13A	0.3314	0.2620	0.1484	0.097*	
H13B	0.4195	0.3218	0.1360	0.097*	
C14	0.3909 (4)	0.2012 (4)	0.13669 (13)	0.0886 (18)	
H14A	0.3405	0.1473	0.1369	0.133*	
H14B	0.4214	0.2090	0.1526	0.133*	
H14C	0.4252	0.2040	0.1222	0.133*	
C21	0.2550 (3)	0.4108 (3)	0.08358 (10)	0.0649 (13)	
H21A	0.3108	0.4587	0.0862	0.078*	
H21B	0.2212	0.4081	0.0984	0.078*	
C22	0.2179 (5)	0.4266 (4)	0.05962 (15)	0.107 (3)	
H22A	0.2472	0.4229	0.0444	0.128*	
H22B	0.1595	0.3828	0.0582	0.128*	
C23	0.2222 (4)	0.5059 (4)	0.05953 (12)	0.0854 (17)	
H23A	0.2808	0.5494	0.0606	0.102*	
H23B	0.1942	0.5099	0.0750	0.102*	
C24	0.1854 (6)	0.5224 (5)	0.03716 (18)	0.136 (4)	
H24A	0.2026	0.5059	0.0216	0.204*	
H24B	0.2039	0.5814	0.0364	0.204*	
H24C	0.1252	0.4910	0.0386	0.204*	
O1W	0.6667	0.3333	0.0468 (7)	0.095 (8)	0.25

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mo1	0.02590 (17)	0.02668 (17)	0.02253 (19)	0.01297 (12)	-0.00049 (9)	-0.00092 (9)
Cl1	0.0327 (4)	0.0327 (4)	0.0241 (6)	0.0163 (2)	0.000	0.000
Cl2	0.0400 (4)	0.0387 (4)	0.0356 (4)	0.0212 (4)	-0.0011 (3)	0.0002 (3)
S1	0.0345 (5)	0.0652 (6)	0.0409 (5)	0.0273 (4)	-0.0101 (4)	-0.0199 (4)
S2	0.0400 (4)	0.0387 (4)	0.0356 (4)	0.0212 (4)	-0.0011 (3)	0.0002 (3)
O2W	0.049 (7)	0.116 (10)	0.122 (10)	0.024 (6)	0.007 (6)	0.007 (7)
N1	0.060 (2)	0.051 (3)	0.052 (3)	0.0254 (14)	-0.0121 (11)	-0.024 (2)
C1	0.0364 (19)	0.054 (2)	0.057 (2)	0.0256 (18)	-0.0009 (17)	-0.0042 (18)
C2	0.061 (3)	0.210 (8)	0.076 (4)	0.086 (5)	-0.021 (3)	-0.020 (4)
C3	0.061 (3)	0.066 (4)	0.219 (8)	0.040 (3)	0.019 (4)	-0.011 (4)
C4	0.055 (3)	0.126 (6)	0.100 (5)	0.034 (4)	0.019 (3)	0.031 (4)
C11	0.062 (3)	0.056 (3)	0.044 (2)	0.024 (2)	-0.0055 (19)	-0.0146 (18)
C12	0.082 (3)	0.063 (3)	0.058 (3)	0.039 (3)	-0.011 (2)	-0.015 (2)
C13	0.110 (5)	0.063 (3)	0.063 (3)	0.038 (3)	-0.013 (3)	-0.002 (2)
C14	0.100 (5)	0.088 (4)	0.082 (4)	0.050 (4)	-0.012 (3)	0.009 (3)
C21	0.061 (3)	0.057 (3)	0.075 (3)	0.028 (2)	-0.019 (2)	-0.033 (2)
C22	0.131 (6)	0.086 (4)	0.122 (5)	0.070 (4)	-0.078 (5)	-0.058 (4)
C23	0.086 (4)	0.081 (4)	0.094 (4)	0.045 (3)	-0.018 (3)	-0.026 (3)
C24	0.157 (8)	0.091 (5)	0.181 (8)	0.078 (5)	-0.083 (7)	-0.039 (5)
O1W	0.086 (8)	0.086 (8)	0.113 (12)	0.043 (4)	0.000	0.000

Geometric parameters (Å, °)

Mo1—Mo1 ⁱ	2.6067 (4)	C3—H3C	0.9800
Mo1—Mo1 ⁱⁱ	2.6067 (4)	C4—H4A	0.9800
Mo1—Mo1 ⁱⁱⁱ	2.6328 (5)	C4—H4B	0.9800
Mo1—Mo1 ^{iv}	2.6328 (5)	C4—H4C	0.9800
Mo1—S1	2.5158 (9)	C11—C12	1.515 (7)
Mo1—S2 ^{iv}	2.4792 (9)	C11—H11A	0.9900
Mo1—S2 ⁱⁱ	2.4842 (9)	C11—H11B	0.9900
Mo1—Cl1	2.5054 (10)	C12—C13	1.500 (7)
Mo1—Cl2	2.4801 (9)	C12—H12A	0.9900
Mo1—Cl2 ^{iv}	2.4792 (9)	C12—H12B	0.9900
Mo1—Cl2 ⁱⁱ	2.4842 (9)	C13—C14	1.501 (8)
Cl1—Mo1 ⁱⁱⁱ	2.5054 (10)	C13—H13A	0.9900
Cl1—Mo1 ^{iv}	2.5054 (10)	C13—H13B	0.9900
Cl2—Mo1 ⁱⁱⁱ	2.4792 (9)	C14—H14A	0.9800
Cl2—Mo1 ⁱ	2.4842 (9)	C14—H14B	0.9800
S1—C1	1.849 (4)	C14—H14C	0.9800
O2W—O2W ^v	1.22 (4)	C21—C22	1.535 (8)
N1—C21 ^v	1.520 (5)	C21—H21A	0.9900
N1—C21	1.520 (5)	C21—H21B	0.9900
N1—C11 ^v	1.525 (5)	C22—C23	1.448 (8)
N1—C11	1.525 (5)	C22—H22A	0.9900
C1—C3	1.497 (7)	C22—H22B	0.9900
C1—C2	1.508 (6)	C23—C24	1.470 (9)
C1—C4	1.515 (7)	C23—H23A	0.9900
C2—H2A	0.9800	C23—H23B	0.9900
C2—H2B	0.9800	C24—H24A	0.9800
C2—H2C	0.9800	C24—H24B	0.9800
C3—H3A	0.9800	C24—H24C	0.9800
C3—H3B	0.9800		
S2 ^{iv} —Mo1—Cl2 ^{iv}	0.00 (5)	C2—C1—C4	106.6 (5)
S2 ^{iv} —Mo1—Cl2	175.69 (3)	C3—C1—S1	111.9 (3)
Cl2 ^{iv} —Mo1—Cl2	175.69 (3)	C2—C1—S1	106.4 (3)
S2 ^{iv} —Mo1—S2 ⁱⁱ	90.61 (2)	C4—C1—S1	111.6 (4)
Cl2 ^{iv} —Mo1—S2 ⁱⁱ	90.61 (2)	C1—C2—H2A	109.5
Cl2—Mo1—S2 ⁱⁱ	90.59 (2)	C1—C2—H2B	109.5
S2 ^{iv} —Mo1—Cl2 ⁱⁱ	90.61 (2)	H2A—C2—H2B	109.5
Cl2 ^{iv} —Mo1—Cl2 ⁱⁱ	90.61 (2)	C1—C2—H2C	109.5
Cl2—Mo1—Cl2 ⁱⁱ	90.59 (2)	H2A—C2—H2C	109.5
S2 ⁱⁱ —Mo1—Cl2 ⁱⁱ	0.00 (5)	H2B—C2—H2C	109.5
S2 ^{iv} —Mo1—Cl1	89.24 (2)	C1—C3—H3A	109.5
Cl2 ^{iv} —Mo1—Cl1	89.24 (2)	C1—C3—H3B	109.5
Cl2—Mo1—Cl1	89.22 (2)	H3A—C3—H3B	109.5
S2 ⁱⁱ —Mo1—Cl1	175.32 (3)	C1—C3—H3C	109.5
Cl2 ⁱⁱ —Mo1—Cl1	175.32 (3)	H3A—C3—H3C	109.5
S2 ^{iv} —Mo1—S1	89.10 (3)	H3B—C3—H3C	109.5

Cl2 ^{iv} —Mo1—S1	89.10 (3)	C1—C4—H4A	109.5
Cl2—Mo1—S1	94.93 (3)	C1—C4—H4B	109.5
S2 ⁱⁱ —Mo1—S1	94.79 (3)	H4A—C4—H4B	109.5
Cl2 ⁱⁱ —Mo1—S1	94.79 (3)	C1—C4—H4C	109.5
Cl1—Mo1—S1	89.88 (3)	H4A—C4—H4C	109.5
S2 ^{iv} —Mo1—Mo1 ⁱ	119.05 (2)	H4B—C4—H4C	109.5
Cl2 ^{iv} —Mo1—Mo1 ⁱ	119.05 (2)	C12—C11—N1	115.2 (3)
Cl2—Mo1—Mo1 ⁱ	58.40 (2)	C12—C11—H11A	108.5
S2 ⁱⁱ —Mo1—Mo1 ⁱ	58.22 (2)	N1—C11—H11A	108.5
Cl2 ⁱⁱ —Mo1—Mo1 ⁱ	58.22 (2)	C12—C11—H11B	108.5
Cl1—Mo1—Mo1 ⁱ	117.960 (18)	N1—C11—H11B	108.5
S1—Mo1—Mo1 ⁱ	138.62 (2)	H11A—C11—H11B	107.5
S2 ^{iv} —Mo1—Mo1 ⁱⁱ	58.41 (2)	C13—C12—C11	112.5 (4)
Cl2 ^{iv} —Mo1—Mo1 ⁱⁱ	58.41 (2)	C13—C12—H12A	109.1
Cl2—Mo1—Mo1 ⁱⁱ	119.04 (2)	C11—C12—H12A	109.1
S2 ⁱⁱ —Mo1—Mo1 ⁱⁱ	58.25 (2)	C13—C12—H12B	109.1
Cl2 ⁱⁱ —Mo1—Mo1 ⁱⁱ	58.25 (2)	C11—C12—H12B	109.1
Cl1—Mo1—Mo1 ⁱⁱ	117.960 (17)	H12A—C12—H12B	107.8
S1—Mo1—Mo1 ⁱⁱ	134.36 (3)	C12—C13—C14	114.0 (5)
Mo1 ⁱ —Mo1—Mo1 ⁱⁱ	60.665 (13)	C12—C13—H13A	108.8
S2 ^{iv} —Mo1—Mo1 ⁱⁱⁱ	117.95 (2)	C14—C13—H13A	108.8
Cl2 ^{iv} —Mo1—Mo1 ⁱⁱⁱ	117.95 (2)	C12—C13—H13B	108.8
Cl2—Mo1—Mo1 ⁱⁱⁱ	57.92 (2)	C14—C13—H13B	108.8
S2 ⁱⁱ —Mo1—Mo1 ⁱⁱⁱ	117.86 (2)	H13A—C13—H13B	107.7
Cl2 ⁱⁱ —Mo1—Mo1 ⁱⁱⁱ	117.86 (2)	C13—C14—H14A	109.5
Cl1—Mo1—Mo1 ⁱⁱⁱ	58.302 (15)	C13—C14—H14B	109.5
S1—Mo1—Mo1 ⁱⁱⁱ	135.38 (3)	H14A—C14—H14B	109.5
Mo1 ⁱ —Mo1—Mo1 ⁱⁱⁱ	59.667 (7)	C13—C14—H14C	109.5
Mo1 ⁱⁱ —Mo1—Mo1 ⁱⁱⁱ	90.0	H14A—C14—H14C	109.5
S2 ^{iv} —Mo1—Mo1 ^{iv}	57.95 (2)	H14B—C14—H14C	109.5
Cl2 ^{iv} —Mo1—Mo1 ^{iv}	57.95 (2)	N1—C21—C22	116.1 (4)
Cl2—Mo1—Mo1 ^{iv}	117.91 (2)	N1—C21—H21A	108.3
S2 ⁱⁱ —Mo1—Mo1 ^{iv}	117.89 (2)	C22—C21—H21A	108.3
Cl2 ⁱⁱ —Mo1—Mo1 ^{iv}	117.89 (2)	N1—C21—H21B	108.3
Cl1—Mo1—Mo1 ^{iv}	58.302 (15)	C22—C21—H21B	108.3
S1—Mo1—Mo1 ^{iv}	131.38 (2)	H21A—C21—H21B	107.4
Mo1 ⁱ —Mo1—Mo1 ^{iv}	90.0	C23—C22—C21	113.7 (5)
Mo1 ⁱⁱ —Mo1—Mo1 ^{iv}	59.667 (7)	C23—C22—H22A	108.8
Mo1 ⁱⁱⁱ —Mo1—Mo1 ^{iv}	60.0	C21—C22—H22A	108.8
Mo1 ⁱⁱⁱ —Cl1—Mo1 ^{iv}	63.40 (3)	C23—C22—H22B	108.8
Mo1 ⁱⁱⁱ —Cl1—Mo1	63.40 (3)	C21—C22—H22B	108.8
Mo1 ^{iv} —Cl1—Mo1	63.40 (3)	H22A—C22—H22B	107.7
Mo1 ⁱⁱⁱ —Cl2—Mo1	64.13 (2)	C22—C23—C24	115.3 (5)
Mo1 ⁱⁱⁱ —Cl2—Mo1 ⁱ	63.36 (2)	C22—C23—H23A	108.4
Mo1—Cl2—Mo1 ⁱ	63.35 (2)	C24—C23—H23A	108.4
C1—S1—Mo1	118.11 (13)	C22—C23—H23B	108.4
C21 ^v —N1—C21	111.8 (5)	C24—C23—H23B	108.4
C21 ^v —N1—C11 ^v	107.2 (2)	H23A—C23—H23B	107.5

C21—N1—C11 ^v	110.3 (3)	C23—C24—H24A	109.5
C21 ^v —N1—C11	110.3 (3)	C23—C24—H24B	109.5
C21—N1—C11	107.2 (2)	H24A—C24—H24B	109.5
C11 ^v —N1—C11	110.1 (5)	C23—C24—H24C	109.5
C3—C1—C2	113.8 (5)	H24A—C24—H24C	109.5
C3—C1—C4	106.5 (5)	H24B—C24—H24C	109.5

Symmetry codes: (i) $y, -x+y, -z$; (ii) $x-y, x, -z$; (iii) $-x+y, -x, z$; (iv) $-y, x-y, z$; (v) $x-y+1/3, -y+2/3, -z+1/6$.