

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## (2E)-1-(2,6-Dichloro-3-fluorophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one

 A. S. Praveen,<sup>a</sup> Jerry P. Jasinski,<sup>b\*</sup> James A. Golen,<sup>b</sup>  
 H. S. Yathirajan<sup>a</sup> and B. Narayana<sup>c</sup>

<sup>a</sup>Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, <sup>b</sup>Department of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, and <sup>c</sup>Department of Studies in Chemistry, Mangalore University, Mangalagangotri, 574 199, India  
 Correspondence e-mail: jjasinski@keene.edu


Received 12 March 2012; accepted 19 March 2012

Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in main residue;  $R$  factor = 0.061;  $wR$  factor = 0.145; data-to-parameter ratio = 17.5.

There are two independent molecules in the asymmetric unit of the title compound,  $\text{C}_{16}\text{H}_{11}\text{Cl}_2\text{FO}_2$ . The F atom equally populates both *meta* positions of the 6-dichloro-3-fluorophenyl ring in each molecule, resulting in 0.5 occupancy for both the F and H atoms in these positions. The dihedral angle between the mean planes of the benzene rings are  $77.5$  (2) and  $89.8$  (8)° in the two molecules. In the crystal, weak  $\text{C}-\text{H}\cdots\text{F}$  and  $\text{C}-\text{H}\cdots\text{O}$  interactions involving the half-occupied H and F atoms are observed. Weak  $\pi-\pi$  stacking interactions [centroid-centroid distance =  $3.150$  (2) Å] also contribute to the crystal stability.

### Related literature

For the pharmacological importance of chalcones, see: Dominguez *et al.* (2001); Li *et al.* (1995); Mei *et al.* (2001); Sarojini *et al.* (2006). For related structures, see: Betz *et al.* (2012); Yathirajan *et al.* (2007). For standard bond lengths, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

$\text{C}_{16}\text{H}_{11}\text{Cl}_2\text{FO}_2$   
 $M_r = 325.15$

Monoclinic,  $P2_1/c$   
 $a = 11.9035$  (6) Å

$b = 10.4472$  (5) Å  
 $c = 23.7435$  (12) Å  
 $\beta = 92.296$  (4)°  
 $V = 2950.3$  (3) Å<sup>3</sup>  
 $Z = 8$

Mo  $K\alpha$  radiation  
 $\mu = 0.45$  mm<sup>-1</sup>  
 $T = 173$  K  
 $0.24 \times 0.20 \times 0.17$  mm

#### Data collection

Oxford Diffraction Xcalibur Eos Gemini diffractometer  
 Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2010)  
 $T_{\min} = 0.900$ ,  $T_{\max} = 0.927$

15257 measured reflections  
 7015 independent reflections  
 5165 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.023$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.145$   
 $S = 1.06$   
 7015 reflections  
 401 parameters

4 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.67$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.57$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C2}-\text{H2A}\cdots\text{F1A}^i$	0.95	2.79	3.657 (7)	153
$\text{C4}-\text{H4A}\cdots\text{F1}^{ii}$	0.95	2.75	3.410 (5)	127
$\text{C11}-\text{H11A}\cdots\text{O3}^{ii}$	0.95	2.56	3.451 (3)	156

Symmetry codes: (i)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

ASP thanks the University of Mysore for research facilities. JPJ acknowledges the NSF-MRI program (grant No. CHE1039027) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5846).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.  
 Betz, R., Gerber, T., Hosten, E., Praveen, A. S., Yathirajan, H. S. & Narayana, B. (2012). *Acta Cryst. E* **68**, o512.  
 Dominguez, J. N., Charris, J. E., Lobo, G., de Dominguez, N. G., Moreno, M. M., Riggione, F., Sanchez, E., Olson, J. & Rosenthal, P. J. (2001). *Eur. J. Med. Chem.* **36**, 555–560.  
 Li, R., Chen, X., Gong, B., Dominguez, J. N., Davidson, E., Kurzban, G., Miller, R. E., Nuzum, E. O. & Rosenthal, P. J. (1995). *J. Med. Chem.* **38**, 5031–5037.  
 Mei, L., Prapon, W. & Mei, L. G. (2001). *J. Med. Chem.* **44**, 4443–4452.  
 Oxford Diffraction (2010). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.  
 Sarojini, B. K., Narayana, B., Ashalatha, B. V., Indira, J. & Lobo, K. G. (2006). *J. Cryst. Growth*, **295**, 54–59.  
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
 Yathirajan, H. S., Mayekar, A. N., Narayana, B., Sarojini & Bolte, M. (2007). *Acta Cryst. E* **63**, o428–o429.

## supporting information

*Acta Cryst.* (2012). E68, o1163 [https://doi.org/10.1107/S1600536812011841]

**(2E)-1-(2,6-Dichloro-3-fluorophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one**

**A. S. Praveen, Jerry P. Jasinski, James A. Golen, H. S. Yathirajan and B. Narayana**

**S1. Comment**

Many Chalcones are known to exhibit various biological properties such as antimalarial (Li *et al.*, 1995), antifungal (Dominguez *et al.*, 2001) and antibacterial activity (Mei *et al.*, 2001). They are also finding application as organic nonlinear optical materials (NLO) for their SHG conversion efficiency (Sarojini *et al.*, 2006). Crystal structures of some related chalcones, *viz.*, (2E)-1-(2,4-dichlorophenyl)-3-(2-hydroxy-3-methoxyphenyl)prop-2-en-1-one (Yathirajan *et al.*, 2007) and (2E)-1-(2,6-dichloro-3-fluorophenyl)-3-(4-fluorophenyl)prop-2-en-1-one (Betz *et al.*, 2012) have been reported. As part of our ongoing studies on chalcones, the title compound (I), C<sub>16</sub>H<sub>11</sub>C<sub>12</sub>FO<sub>2</sub>, was synthesized and its crystal structure is reported.

In (I) two molecules crystallize in the asymmetric unit (Fig. 1). In the 2,6-dichloro-3-fluorophenyl ring, the fluorine atom equally populates both *meta* positions of the phenyl ring in each molecule (C2 & C4; C18 & C20) resulting in 0.5 occupancy for both the fluorine and hydrogen atoms (H2A & H4A; H18A & H20A) in these positions. The dihedral angle between the mean planes of the benzene rings in each molecule is 77.5 (2)° and 89.8 (8)°, respectively. Bond lengths are in normal ranges (Allen *et al.*, 1987). Crystal packing is enhanced by weak C—H···F and C—H···O intermolecular interactions (Table 1) from both half-occupied H and F atoms supporting parallel chains along the *b* axis (Fig. 2) as well as weak  $\pi$ – $\pi$  stacking interactions (Table 2).

**S2. Experimental**

To a stirred solution of 1-(2,6-dichloro-3-fluorophenyl)ethanone (1 g, 4.8 mmol) and 4-methoxybenzaldehyde (0.65 g, 4.8 mmol) in ethanol (10 ml), powdered KOH (0.40 g, 7.2 mmol) was added at 273 K. The reaction mixture was stirred at room temperature for 3 h. After completion of the reaction, the reaction mixture was poured to ice cold water and acidified with 1.5 N HCl (pH 3). The solid precipitated was filtered and dried to afford 1.4 g of the title compound, (I), in 89% yield. X-ray quality crystals were obtained by slow evaporation of a tetrahydrofuran solution (m.p.: 361–362 K).

**S3. Refinement**

All of the H atoms were placed in their calculated positions and refined using the riding model with C—H lengths of 0.95 Å (CH) or 0.98 Å (CH<sub>3</sub>). The isotropic displacement parameters for these atoms were set from 1.19 to 1.20 (CH), or 1.49 (CH<sub>3</sub>) times  $U_{eq}$  of the parent atom. Overlapping of the F atoms in the *meta* position of the phenyl ring in each molecule resulted in H2A, H4A, H18A, H20A and F1, F1A, F2, F2A being refined at 0.50 occupancy. C2—F1 and C4—F1A bond distances were fixed at 1.33 (2) Å, C18—F2 and C20—F2A bond distances were fixed at 1.33 (06) Å.




Figure 1

Molecular structure of the title compound showing the atom labeling scheme and 50% probability displacement ellipsoids for two molecules in the asymmetric unit.




Figure 2

Packing diagram of the title compound viewed along the *a* axis.

(2*E*)-1-(2,6-Dichloro-3-fluorophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one

*Crystal data*

C<sub>16</sub>H<sub>11</sub>Cl<sub>2</sub>FO<sub>2</sub>

*M<sub>r</sub>* = 325.15

Monoclinic, *P*2<sub>1</sub>/*c*

Hall symbol: -*P* 2ybc

*a* = 11.9035 (6) Å

*b* = 10.4472 (5) Å

*c* = 23.7435 (12) Å

β = 92.296 (4)°

*V* = 2950.3 (3) Å<sup>3</sup>

*Z* = 8

*F*(000) = 1328

*D<sub>x</sub>* = 1.464 Mg m<sup>-3</sup>

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 3879 reflections

θ = 3.1–30.0°

μ = 0.45 mm<sup>-1</sup>

*T* = 173 K

Block, colorless

0.24 × 0.20 × 0.17 mm

*Data collection*

Oxford Diffraction Xcalibur Eos Gemini diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: 16.1500 pixels mm<sup>-1</sup>

ω scans

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2010)

*T<sub>min</sub>* = 0.900, *T<sub>max</sub>* = 0.927

15257 measured reflections

7015 independent reflections

5165 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.023

θ<sub>max</sub> = 27.9°, θ<sub>min</sub> = 3.1°

*h* = -10→15

*k* = -13→10

*l* = -30→31

*Refinement*

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.061

*wR*(*F*<sup>2</sup>) = 0.145

*S* = 1.06

7015 reflections

401 parameters

4 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

*w* = 1/[σ<sup>2</sup>(*F<sub>o</sub>*<sup>2</sup>) + (0.042*P*)<sup>2</sup> + 3.2887*P*]

where *P* = (*F<sub>o</sub>*<sup>2</sup> + 2*F<sub>c</sub>*<sup>2</sup>)/3

(Δ/σ)<sub>max</sub> < 0.001

Δρ<sub>max</sub> = 0.67 e Å<sup>-3</sup>

Δρ<sub>min</sub> = -0.57 e Å<sup>-3</sup>

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of *F*<sup>2</sup> against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*<sup>2</sup>, conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*<sup>2</sup>. The threshold expression of *F*<sup>2</sup> > σ(*F*<sup>2</sup>) is used only for calculating *R*-factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*<sup>2</sup> are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U<sub>iso</sub></i> */ <i>U<sub>eq</sub></i>	Occ. (<1)
Cl1	0.73833 (6)	0.53784 (8)	0.11864 (4)	0.0566 (2)	

C12	0.73589 (8)	0.86623 (10)	0.29381 (4)	0.0697 (3)	
C13	0.75952 (8)	0.51754 (10)	0.41195 (6)	0.0931 (4)	
C14	0.75948 (8)	0.21696 (11)	0.22698 (4)	0.0719 (3)	
F1	0.5430 (3)	0.4344 (3)	0.16850 (18)	0.0600 (10)	0.50
F1A	0.5483 (5)	0.7210 (7)	0.3246 (2)	0.121 (2)	0.50
F2	0.9551 (3)	0.6350 (3)	0.3500 (2)	0.0771 (13)	0.50
F2A	0.9515 (4)	0.3907 (7)	0.2061 (2)	0.118 (2)	0.50
O1	0.91972 (15)	0.72821 (19)	0.18987 (9)	0.0473 (5)	
O2	0.57619 (15)	1.3317 (2)	−0.00869 (9)	0.0499 (5)	
O3	0.57623 (15)	0.3266 (2)	0.33521 (10)	0.0514 (5)	
O4	1.00724 (16)	−0.28448 (19)	0.48786 (9)	0.0499 (5)	
C1	0.6802 (2)	0.5932 (3)	0.17975 (13)	0.0418 (6)	
C2	0.5874 (2)	0.5310 (3)	0.20029 (16)	0.0554 (8)	
H2A	0.5557	0.4601	0.1804	0.066*	0.50
C3	0.5411 (3)	0.5714 (3)	0.24921 (17)	0.0611 (9)	
H3A	0.4776	0.5287	0.2634	0.073*	
C4	0.5875 (3)	0.6741 (4)	0.27727 (14)	0.0570 (9)	
H4A	0.5564	0.7018	0.3114	0.068*	0.50
C5	0.6787 (2)	0.7381 (3)	0.25700 (12)	0.0451 (7)	
C6	0.7264 (2)	0.6993 (2)	0.20724 (11)	0.0355 (5)	
C7	0.8245 (2)	0.7701 (2)	0.18319 (11)	0.0361 (6)	
C8	0.8000 (2)	0.8855 (2)	0.15059 (11)	0.0360 (5)	
H8A	0.8614	0.9377	0.1405	0.043*	
C9	0.6966 (2)	0.9225 (2)	0.13400 (11)	0.0357 (6)	
H9A	0.6362	0.8733	0.1475	0.043*	
C10	0.6665 (2)	1.0297 (2)	0.09763 (11)	0.0345 (5)	
C11	0.5534 (2)	1.0535 (3)	0.08407 (13)	0.0447 (7)	
H11A	0.4979	1.0002	0.0997	0.054*	
C12	0.5196 (2)	1.1523 (3)	0.04865 (13)	0.0470 (7)	
H12A	0.4420	1.1663	0.0400	0.056*	
C13	0.5993 (2)	1.2302 (3)	0.02593 (12)	0.0376 (6)	
C14	0.7130 (2)	1.2079 (3)	0.03816 (12)	0.0402 (6)	
H14A	0.7680	1.2607	0.0219	0.048*	
C15	0.7457 (2)	1.1101 (3)	0.07347 (12)	0.0386 (6)	
H15A	0.8235	1.0964	0.0818	0.046*	
C16	0.4637 (3)	1.3482 (3)	−0.02947 (16)	0.0602 (9)	
H16A	0.4596	1.4221	−0.0549	0.090*	
H16B	0.4148	1.3628	0.0022	0.090*	
H16C	0.4390	1.2712	−0.0500	0.090*	
C17	0.8124 (3)	0.4772 (3)	0.34698 (17)	0.0618 (9)	
C18	0.9025 (3)	0.5439 (3)	0.3250 (3)	0.0935 (18)	
H18A	0.9327	0.6150	0.3454	0.112*	0.50
C19	0.9488 (3)	0.5117 (5)	0.2757 (3)	0.104 (2)	
H19A	1.0113	0.5562	0.2616	0.125*	
C20	0.9013 (3)	0.4145 (4)	0.2488 (2)	0.0862 (15)	
H20A	0.9319	0.3914	0.2139	0.103*	0.50
C21	0.8131 (2)	0.3439 (3)	0.26580 (15)	0.0562 (9)	
C22	0.7669 (2)	0.3748 (3)	0.31718 (14)	0.0465 (7)	

C23	0.6745 (2)	0.2949 (3)	0.34163 (12)	0.0406 (6)
C24	0.7103 (2)	0.1816 (3)	0.37319 (12)	0.0400 (6)
H24A	0.6546	0.1310	0.3902	0.048*
C25	0.8174 (2)	0.1453 (2)	0.37936 (11)	0.0360 (6)
H25A	0.8709	0.1979	0.3617	0.043*
C26	0.86199 (19)	0.0345 (2)	0.40993 (11)	0.0328 (5)
C27	0.9691 (2)	-0.0094 (3)	0.39798 (11)	0.0367 (6)
H27A	1.0117	0.0353	0.3713	0.044*
C28	1.0141 (2)	-0.1161 (3)	0.42405 (12)	0.0418 (6)
H28A	1.0864	-0.1458	0.4146	0.050*
C29	0.9545 (2)	-0.1804 (2)	0.46389 (11)	0.0356 (5)
C30	0.8491 (2)	-0.1371 (3)	0.47756 (12)	0.0404 (6)
H30A	0.8083	-0.1801	0.5055	0.049*
C31	0.8035 (2)	-0.0312 (3)	0.45043 (12)	0.0396 (6)
H31A	0.7307	-0.0026	0.4596	0.048*
C32	0.9572 (3)	-0.3425 (3)	0.53478 (13)	0.0508 (7)
H32A	1.0056	-0.4118	0.5494	0.076*
H32B	0.8835	-0.3773	0.5230	0.076*
H32C	0.9479	-0.2783	0.5644	0.076*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0398 (4)	0.0599 (5)	0.0693 (5)	0.0048 (3)	-0.0089 (3)	-0.0187 (4)
C12	0.0707 (6)	0.0786 (6)	0.0601 (5)	-0.0014 (5)	0.0070 (4)	-0.0206 (5)
C13	0.0633 (6)	0.0740 (6)	0.1389 (10)	0.0131 (5)	-0.0371 (6)	-0.0454 (7)
C14	0.0612 (5)	0.0891 (7)	0.0663 (6)	0.0046 (5)	0.0130 (4)	0.0001 (5)
F1	0.0344 (17)	0.0391 (17)	0.106 (3)	-0.0165 (14)	-0.0007 (17)	-0.0054 (19)
F1A	0.111 (4)	0.154 (6)	0.102 (4)	0.008 (4)	0.048 (3)	0.035 (4)
F2	0.061 (2)	0.0393 (19)	0.129 (4)	-0.0120 (18)	-0.026 (2)	-0.012 (2)
F2A	0.077 (3)	0.191 (7)	0.086 (4)	0.000 (4)	0.016 (3)	0.056 (4)
O1	0.0286 (9)	0.0475 (11)	0.0655 (13)	0.0027 (8)	-0.0018 (8)	0.0070 (10)
O2	0.0358 (10)	0.0515 (12)	0.0622 (13)	0.0030 (9)	-0.0004 (9)	0.0201 (10)
O3	0.0288 (9)	0.0486 (11)	0.0764 (15)	0.0101 (9)	-0.0027 (9)	-0.0001 (11)
O4	0.0418 (10)	0.0481 (11)	0.0606 (13)	0.0136 (9)	0.0100 (9)	0.0170 (10)
C1	0.0287 (12)	0.0377 (14)	0.0585 (18)	0.0052 (11)	-0.0052 (11)	0.0057 (13)
C2	0.0306 (14)	0.0423 (16)	0.093 (3)	-0.0017 (12)	-0.0051 (15)	0.0131 (17)
C3	0.0391 (16)	0.056 (2)	0.089 (3)	0.0004 (15)	0.0112 (16)	0.0328 (19)
C4	0.0467 (17)	0.070 (2)	0.0554 (19)	0.0143 (16)	0.0148 (14)	0.0232 (17)
C5	0.0413 (15)	0.0473 (16)	0.0468 (16)	0.0065 (13)	-0.0002 (12)	0.0086 (13)
C6	0.0270 (12)	0.0346 (13)	0.0445 (14)	0.0044 (10)	-0.0036 (10)	0.0077 (11)
C7	0.0307 (12)	0.0364 (13)	0.0410 (14)	-0.0006 (10)	-0.0006 (10)	-0.0016 (11)
C8	0.0323 (12)	0.0354 (13)	0.0402 (14)	-0.0037 (10)	0.0011 (10)	0.0012 (11)
C9	0.0312 (12)	0.0336 (13)	0.0424 (14)	-0.0019 (10)	0.0034 (10)	0.0000 (11)
C10	0.0287 (11)	0.0339 (13)	0.0409 (14)	-0.0003 (10)	0.0013 (10)	0.0003 (11)
C11	0.0280 (12)	0.0464 (15)	0.0602 (18)	-0.0030 (11)	0.0065 (12)	0.0119 (14)
C12	0.0273 (12)	0.0516 (17)	0.0621 (19)	0.0025 (12)	0.0019 (12)	0.0139 (15)
C13	0.0318 (12)	0.0367 (13)	0.0442 (15)	0.0015 (11)	-0.0005 (10)	0.0028 (12)

C14	0.0308 (12)	0.0400 (14)	0.0499 (16)	-0.0041 (11)	0.0027 (11)	0.0063 (12)
C15	0.0274 (12)	0.0395 (14)	0.0486 (15)	-0.0014 (11)	-0.0029 (10)	0.0029 (12)
C16	0.0421 (16)	0.059 (2)	0.078 (2)	0.0068 (15)	-0.0109 (15)	0.0170 (18)
C17	0.0412 (16)	0.0400 (16)	0.102 (3)	0.0077 (13)	-0.0227 (17)	0.0070 (17)
C18	0.048 (2)	0.050 (2)	0.178 (5)	-0.0162 (18)	-0.052 (3)	0.047 (3)
C19	0.041 (2)	0.103 (4)	0.166 (5)	-0.017 (2)	-0.022 (3)	0.098 (4)
C20	0.0372 (18)	0.099 (3)	0.121 (4)	-0.001 (2)	-0.007 (2)	0.073 (3)
C21	0.0315 (14)	0.0593 (19)	0.078 (2)	0.0049 (13)	-0.0020 (14)	0.0289 (18)
C22	0.0287 (13)	0.0353 (14)	0.075 (2)	0.0052 (11)	-0.0088 (13)	0.0157 (14)
C23	0.0303 (13)	0.0363 (14)	0.0549 (17)	0.0047 (11)	-0.0011 (11)	-0.0043 (13)
C24	0.0309 (12)	0.0375 (13)	0.0518 (16)	0.0017 (11)	0.0052 (11)	0.0027 (12)
C25	0.0317 (12)	0.0324 (12)	0.0440 (15)	0.0010 (10)	0.0020 (10)	-0.0013 (11)
C26	0.0286 (11)	0.0312 (12)	0.0384 (14)	0.0008 (10)	-0.0013 (10)	-0.0047 (11)
C27	0.0290 (12)	0.0388 (13)	0.0425 (14)	0.0020 (10)	0.0054 (10)	0.0031 (12)
C28	0.0287 (12)	0.0472 (15)	0.0498 (16)	0.0081 (11)	0.0069 (11)	0.0037 (13)
C29	0.0326 (12)	0.0328 (12)	0.0412 (14)	0.0046 (10)	-0.0009 (10)	0.0014 (11)
C30	0.0361 (13)	0.0418 (14)	0.0439 (15)	0.0021 (12)	0.0094 (11)	0.0047 (12)
C31	0.0298 (12)	0.0402 (14)	0.0496 (16)	0.0057 (11)	0.0096 (11)	0.0015 (12)
C32	0.0507 (17)	0.0482 (17)	0.0536 (18)	0.0037 (14)	0.0040 (14)	0.0162 (14)

*Geometric parameters (Å, °)*

C11—C1	1.732 (3)	C14—H14A	0.9500
C12—C5	1.724 (3)	C15—H15A	0.9500
C13—C17	1.741 (4)	C16—H16A	0.9800
C14—C21	1.723 (4)	C16—H16B	0.9800
O1—C7	1.220 (3)	C16—H16C	0.9800
O2—C13	1.363 (3)	C17—C22	1.381 (4)
O2—C16	1.419 (3)	C17—C18	1.398 (6)
O3—C23	1.219 (3)	C18—C19	1.354 (7)
O4—C29	1.369 (3)	C18—H18A	0.9500
O4—C32	1.420 (3)	C19—C20	1.315 (7)
C1—C2	1.387 (4)	C19—H19A	0.9500
C1—C6	1.388 (4)	C20—C21	1.359 (5)
C2—C3	1.372 (5)	C20—H20A	0.9500
C2—H2A	0.9500	C21—C22	1.396 (5)
C3—C4	1.368 (5)	C22—C23	1.516 (4)
C3—H3A	0.9500	C23—C24	1.456 (4)
C4—C5	1.378 (4)	C24—C25	1.333 (3)
C4—H4A	0.9500	C24—H24A	0.9500
C5—C6	1.391 (4)	C25—C26	1.455 (3)
C6—C7	1.514 (3)	C25—H25A	0.9500
C7—C8	1.456 (4)	C26—C31	1.391 (4)
C8—C9	1.335 (3)	C26—C27	1.395 (3)
C8—H8A	0.9500	C27—C28	1.373 (4)
C9—C10	1.450 (4)	C27—H27A	0.9500
C9—H9A	0.9500	C28—C29	1.379 (4)
C10—C11	1.393 (3)	C28—H28A	0.9500

C10—C15	1.403 (3)	C29—C30	1.384 (3)
C11—C12	1.381 (4)	C30—C31	1.381 (4)
C11—H11A	0.9500	C30—H30A	0.9500
C12—C13	1.376 (4)	C31—H31A	0.9500
C12—H12A	0.9500	C32—H32A	0.9800
C13—C14	1.393 (3)	C32—H32B	0.9800
C14—C15	1.368 (4)	C32—H32C	0.9800
C13—O2—C16	118.0 (2)	H16B—C16—H16C	109.5
C29—O4—C32	117.7 (2)	C22—C17—C18	119.0 (4)
C2—C1—C6	120.9 (3)	C22—C17—C13	119.5 (3)
C2—C1—C11	119.3 (2)	C18—C17—C13	121.4 (3)
C6—C1—C11	119.8 (2)	C19—C18—C17	122.9 (4)
C3—C2—C1	120.5 (3)	C19—C18—H18A	118.5
C3—C2—H2A	119.8	C17—C18—H18A	118.5
C1—C2—H2A	119.8	C20—C19—C18	115.5 (4)
C4—C3—C2	119.0 (3)	C20—C19—H19A	122.3
C4—C3—H3A	120.5	C18—C19—H19A	122.3
C2—C3—H3A	120.5	C19—C20—C21	126.7 (5)
C3—C4—C5	121.3 (3)	C19—C20—H20A	116.7
C3—C4—H4A	119.3	C21—C20—H20A	116.7
C5—C4—H4A	119.3	C20—C21—C22	118.0 (4)
C4—C5—C6	120.6 (3)	C20—C21—C14	122.2 (3)
C4—C5—C12	120.0 (3)	C22—C21—C14	119.7 (2)
C6—C5—C12	119.4 (2)	C17—C22—C21	117.9 (3)
C1—C6—C5	117.7 (2)	C17—C22—C23	120.4 (3)
C1—C6—C7	120.6 (2)	C21—C22—C23	121.6 (3)
C5—C6—C7	121.7 (2)	O3—C23—C24	123.1 (3)
O1—C7—C8	122.1 (2)	O3—C23—C22	120.6 (2)
O1—C7—C6	120.2 (2)	C24—C23—C22	116.3 (2)
C8—C7—C6	117.7 (2)	C25—C24—C23	123.2 (3)
C9—C8—C7	124.0 (2)	C25—C24—H24A	118.4
C9—C8—H8A	118.0	C23—C24—H24A	118.4
C7—C8—H8A	118.0	C24—C25—C26	127.5 (2)
C8—C9—C10	127.0 (2)	C24—C25—H25A	116.2
C8—C9—H9A	116.5	C26—C25—H25A	116.2
C10—C9—H9A	116.5	C31—C26—C27	117.7 (2)
C11—C10—C15	117.2 (2)	C31—C26—C25	123.7 (2)
C11—C10—C9	119.3 (2)	C27—C26—C25	118.6 (2)
C15—C10—C9	123.5 (2)	C28—C27—C26	121.2 (2)
C12—C11—C10	122.0 (2)	C28—C27—H27A	119.4
C12—C11—H11A	119.0	C26—C27—H27A	119.4
C10—C11—H11A	119.0	C27—C28—C29	120.1 (2)
C13—C12—C11	119.5 (2)	C27—C28—H28A	119.9
C13—C12—H12A	120.3	C29—C28—H28A	119.9
C11—C12—H12A	120.3	O4—C29—C28	115.6 (2)
O2—C13—C12	124.8 (2)	O4—C29—C30	124.6 (2)
O2—C13—C14	115.3 (2)	C28—C29—C30	119.9 (2)



C12—C13—C14	119.9 (2)	C31—C30—C29	119.7 (2)
C15—C14—C13	120.2 (2)	C31—C30—H30A	120.2
C15—C14—H14A	119.9	C29—C30—H30A	120.2
C13—C14—H14A	119.9	C30—C31—C26	121.3 (2)
C14—C15—C10	121.2 (2)	C30—C31—H31A	119.3
C14—C15—H15A	119.4	C26—C31—H31A	119.3
C10—C15—H15A	119.4	O4—C32—H32A	109.5
O2—C16—H16A	109.5	O4—C32—H32B	109.5
O2—C16—H16B	109.5	H32A—C32—H32B	109.5
H16A—C16—H16B	109.5	O4—C32—H32C	109.5
O2—C16—H16C	109.5	H32A—C32—H32C	109.5
H16A—C16—H16C	109.5	H32B—C32—H32C	109.5
C6—C1—C2—C3	-1.8 (4)	C22—C17—C18—C19	-0.9 (5)
C11—C1—C2—C3	178.8 (2)	C13—C17—C18—C19	177.1 (3)
C1—C2—C3—C4	0.2 (5)	C17—C18—C19—C20	1.7 (6)
C2—C3—C4—C5	0.8 (5)	C18—C19—C20—C21	-1.0 (6)
C3—C4—C5—C6	-0.3 (4)	C19—C20—C21—C22	-0.5 (5)
C3—C4—C5—C12	-179.4 (2)	C19—C20—C21—C14	-179.1 (3)
C2—C1—C6—C5	2.2 (4)	C18—C17—C22—C21	-0.7 (4)
C11—C1—C6—C5	-178.3 (2)	C13—C17—C22—C21	-178.7 (2)
C2—C1—C6—C7	-177.3 (2)	C18—C17—C22—C23	175.9 (3)
C11—C1—C6—C7	2.2 (3)	C13—C17—C22—C23	-2.2 (4)
C4—C5—C6—C1	-1.2 (4)	C20—C21—C22—C17	1.3 (4)
C12—C5—C6—C1	177.96 (19)	C14—C21—C22—C17	180.0 (2)
C4—C5—C6—C7	178.3 (3)	C20—C21—C22—C23	-175.1 (3)
C12—C5—C6—C7	-2.5 (3)	C14—C21—C22—C23	3.5 (4)
C1—C6—C7—O1	-79.7 (3)	C17—C22—C23—O3	86.1 (4)
C5—C6—C7—O1	100.9 (3)	C21—C22—C23—O3	-97.5 (3)
C1—C6—C7—C8	98.1 (3)	C17—C22—C23—C24	-93.0 (3)
C5—C6—C7—C8	-81.3 (3)	C21—C22—C23—C24	83.4 (3)
O1—C7—C8—C9	167.9 (3)	O3—C23—C24—C25	178.7 (3)
C6—C7—C8—C9	-9.8 (4)	C22—C23—C24—C25	-2.2 (4)
C7—C8—C9—C10	-174.4 (2)	C23—C24—C25—C26	179.9 (3)
C8—C9—C10—C11	179.2 (3)	C24—C25—C26—C31	-18.7 (4)
C8—C9—C10—C15	1.2 (4)	C24—C25—C26—C27	160.9 (3)
C15—C10—C11—C12	-0.4 (4)	C31—C26—C27—C28	1.8 (4)
C9—C10—C11—C12	-178.5 (3)	C25—C26—C27—C28	-177.7 (2)
C10—C11—C12—C13	-0.1 (5)	C26—C27—C28—C29	-1.6 (4)
C16—O2—C13—C12	-9.4 (4)	C32—O4—C29—C28	170.9 (3)
C16—O2—C13—C14	170.9 (3)	C32—O4—C29—C30	-8.3 (4)
C11—C12—C13—O2	-178.8 (3)	C27—C28—C29—O4	-179.2 (2)
C11—C12—C13—C14	0.9 (5)	C27—C28—C29—C30	0.0 (4)
O2—C13—C14—C15	178.6 (3)	O4—C29—C30—C31	-179.7 (3)
C12—C13—C14—C15	-1.1 (4)	C28—C29—C30—C31	1.2 (4)
C13—C14—C15—C10	0.6 (4)	C29—C30—C31—C26	-0.8 (4)
C11—C10—C15—C14	0.1 (4)	C27—C26—C31—C30	-0.6 (4)
C9—C10—C15—C14	178.2 (3)	C25—C26—C31—C30	178.9 (3)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C2—H2A $\cdots$ F1A <sup>i</sup>	0.95	2.79	3.657 (7)	153
C4—H4A $\cdots$ F1 <sup>ii</sup>	0.95	2.75	3.410 (5)	127
C11—H11A $\cdots$ O3 <sup>ii</sup>	0.95	2.56	3.451 (3)	156

Symmetry codes: (i)  $-x+1, y-1/2, -z+1/2$ ; (ii)  $-x+1, y+1/2, -z+1/2$ .