

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,4-Dimethylpyrano[2,3-c]pyrazol-6(2H)-one

 Bilal Shahid,^a Muhammad Zia-ur-Rehman,^{b*} Muhammad Nadeem Arshad,^c Rabia Nazir^b and Ertan Şahin^d

^aDepartment of Chemistry, GC University, Lahore 54000, Pakistan, ^bApplied Chemistry Research Centre, Pakistan Council of Scientific & Industrial Research Laboratories Complex, Lahore 54600, Pakistan, ^cDepartment of Chemistry, University of Gujrat (H. H. Campus), Gujrat 57000, Pakistan, and ^dDepartment of Chemistry, Faculty of Science, Atatürk University, 25240 Erzurum, Turkey
Correspondence e-mail: rehman_pcsir@hotmail.com


Received 22 March 2012; accepted 23 March 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.041; wR factor = 0.117; data-to-parameter ratio = 16.5.

The asymmetric unit of the title compound, $\text{C}_8\text{H}_8\text{N}_2\text{O}_2$, comprises two independent molecules in both of which, all non-H atoms lie in a common plane (r.m.s. deviation = 0.014 and 0.017 Å). In the crystal, $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds connect the molecules into zigzag chains running along $[10\bar{1}]$. Weak $\text{C}-\text{H}\cdots\text{O}$ interactions connect the chains into an infinite network.

Related literature

For related structures, see: Ahmad *et al.* (2011); Ramsay & Steel (1985).



Experimental

Crystal data

$\text{C}_8\text{H}_8\text{N}_2\text{O}_2$
 $M_r = 164.16$
 Monoclinic, $P2_1/n$
 $a = 13.6219$ (3) Å
 $b = 6.8766$ (2) Å

$c = 16.2369$ (4) Å
 $\beta = 96.091$ (2)°
 $V = 1512.36$ (7) Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 0.11$ mm⁻¹
 $T = 296$ K

$0.30 \times 0.18 \times 0.11$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2007)
 $T_{\min} = 0.970$, $T_{\max} = 0.989$

16831 measured reflections
 3770 independent reflections
 2654 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.117$
 $S = 1.02$
 3770 reflections
 228 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.23$ e Å⁻³
 $\Delta\rho_{\min} = -0.16$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N4}-\text{H4N}\cdots\text{O2}^{\text{i}}$	0.91 (2)	1.91 (2)	2.7860 (17)	160.9 (17)
$\text{N2}-\text{H2N}\cdots\text{O4}^{\text{ii}}$	0.912 (19)	1.984 (19)	2.8872 (18)	170.4 (17)
$\text{C2}-\text{H2}\cdots\text{O2}^{\text{iii}}$	0.93	2.49	3.4082 (19)	171
$\text{C7}-\text{H7A}\cdots\text{O3}^{\text{iv}}$	0.96	2.54	3.458 (2)	159
$\text{C10}-\text{H10}\cdots\text{O4}^{\text{iv}}$	0.93	2.45	3.3563 (18)	164

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $x, y + 1, z$; (iii) $-x, -y, -z + 1$; (iv) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

The authors acknowledge the PCSIR Laboratories Complex, Lahore, Pakistan for the provision of facilities necessary to carry out the synthetic work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5857).

References

- Ahmad, N., Tahir, M. N., Khan, M. A., Ather, A. Q. & Khan, M. N. (2011). *Acta Cryst.* **E67**, o1021.
 Bruker (2007). *SADABS*, *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Ramsay, C. G. & Steel, P. J. (1985). *Acta Cryst.* **C41**, 135–136.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o1217 [https://doi.org/10.1107/S1600536812012779]

3,4-Dimethylpyrano[2,3-*c*]pyrazol-6(2*H*)-one

Bilal Shahid, Muhammad Zia-ur-Rehman, Muhammad Nadeem Arshad, Rabia Nazir and Ertan Şahin

S1. Comment

In continuation of our work on the synthesis and biological evaluation of various heterocyclic compounds, we, herein report the crystal structure of the title compound (**I**) Fig. 1.

Structure of the title compound is closely related to already published structures of 3,4-dimethyl-1-(2-pyridyl)pyrano(2,3-*c*)pyrazol-6 (1*H*)-one (Ramsay & Steel, 1985) and 3,4-dimethyl-1-phenylpyrano [2,3-*c*]pyrazol-6(1*H*)-one (Ahmad *et al.*, 2011). There are two asymmetric molecules per unit cell and each molecule comprises of one pyranone ring fused with pyrazole ring. N-H \cdots O hydrogen bonding interactions connect the molecules to zig-zag chains running along [1 0 $\bar{1}$]. In addition, weak intermolecular C—H \cdots O interactions connect these infinite chains to a three dimensional network (Table 1, Fig. 2).

S2. Experimental

A mixture of hydrazine hydrate (1.0 mmole), ethyl acetoacetate (2.0 mmoles) was heated for one hour at 120°C followed by cooling to room temperature. The contents were triturated with diethyl ether, filtered and the residue obtained was crystallized from acetic acid. M.p. 246°C; Yield (87%).

S3. Refinement

All H-atoms bonded to C were positioned with idealized geometry with C—H = 0.93 Å for aromatic and C—H = 0.96 Å for methyl groups and were refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ for aromatic and $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$ for methyl carbon atoms. The coordinates of the H atoms bonded to N were refined with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{N})$.




Figure 1

The labelled molecular structure of (I) showing molecules A and B with 50% displacement ellipsoids.




Figure 2

A perspective view showing hydrogen bond interactions drawn as dashed lines.

3,4-Dimethylpyrano[2,3-c]pyrazol-6(2H)-one

Crystal data

$C_8H_8N_2O_2$

$M_r = 164.16$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 13.6219\ (3)\ \text{\AA}$

$b = 6.8766\ (2)\ \text{\AA}$

$c = 16.2369\ (4)\ \text{\AA}$

$\beta = 96.091\ (2)^\circ$

$V = 1512.36\ (7)\ \text{\AA}^3$

$Z = 8$

$F(000) = 688$

$D_x = 1.443\ \text{Mg m}^{-3}$

Melting point: 518 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5051 reflections

$\theta = 2.5\text{--}27.8^\circ$

$\mu = 0.11\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Needle, yellow

$0.30 \times 0.18 \times 0.11\ \text{mm}$

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2007)

$T_{\min} = 0.970$, $T_{\max} = 0.989$

16831 measured reflections

3770 independent reflections

2654 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 1.9^\circ$

$h = -18 \rightarrow 18$

$k = -8 \rightarrow 9$

$l = -21 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.117$

$S = 1.02$

3770 reflections

228 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0553P)^2 + 0.3127P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.23\ \text{e \AA}^{-3}$

$\Delta\rho_{\min} = -0.16\ \text{e \AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick,

2008), $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0060 (10)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.14791 (7)	0.41930 (15)	0.44365 (6)	0.0399 (3)
O2	0.13549 (8)	0.13137 (18)	0.50096 (7)	0.0512 (3)
O3	0.00479 (8)	0.23381 (15)	0.19680 (7)	0.0417 (3)
O4	0.14510 (8)	0.19091 (16)	0.27386 (7)	0.0493 (3)
N1	0.14966 (10)	0.7225 (2)	0.38066 (8)	0.0445 (3)
N2	0.07651 (10)	0.8288 (2)	0.33725 (9)	0.0446 (3)
N3	-0.14217 (10)	0.2981 (2)	0.11578 (9)	0.0480 (4)
N4	-0.17201 (10)	0.4537 (2)	0.06663 (9)	0.0475 (4)
C1	0.09219 (11)	0.2599 (2)	0.46042 (9)	0.0372 (3)
C2	-0.01109 (11)	0.2563 (2)	0.43051 (9)	0.0375 (3)
H2	-0.0480	0.1485	0.4431	0.045*
C3	-0.05750 (10)	0.4004 (2)	0.38520 (9)	0.0341 (3)
C4	0.00154 (10)	0.5642 (2)	0.36819 (8)	0.0333 (3)
C5	0.10172 (10)	0.5654 (2)	0.39826 (9)	0.0351 (3)
C6	-0.01200 (11)	0.7428 (2)	0.32825 (9)	0.0375 (3)
C7	-0.10067 (12)	0.8367 (3)	0.28407 (10)	0.0487 (4)
H7A	-0.0840	0.9652	0.2671	0.073*
H7B	-0.1518	0.8446	0.3204	0.073*
H7C	-0.1236	0.7610	0.2362	0.073*
C8	-0.16479 (11)	0.3900 (3)	0.35436 (11)	0.0473 (4)
H8A	-0.1921	0.2711	0.3729	0.071*
H8B	-0.1724	0.3937	0.2949	0.071*
H8C	-0.1987	0.4986	0.3755	0.071*
C9	0.09864 (11)	0.2979 (2)	0.22481 (9)	0.0364 (3)
C10	0.13176 (10)	0.4804 (2)	0.19482 (9)	0.0368 (3)
H10	0.1960	0.5196	0.2121	0.044*
C11	0.07528 (10)	0.5986 (2)	0.14298 (9)	0.0332 (3)
C12	-0.02244 (10)	0.5304 (2)	0.11579 (8)	0.0333 (3)
C13	-0.05254 (11)	0.3506 (2)	0.14382 (9)	0.0363 (3)
C14	-0.10441 (11)	0.5942 (2)	0.06431 (9)	0.0380 (3)
C15	-0.12391 (13)	0.7747 (3)	0.01454 (11)	0.0509 (4)
H15A	-0.1839	0.7596	-0.0217	0.076*
H15B	-0.0700	0.7980	-0.0178	0.076*
H15C	-0.1304	0.8828	0.0510	0.076*
C16	0.11158 (13)	0.7903 (2)	0.11518 (11)	0.0481 (4)
H16A	0.1782	0.8107	0.1394	0.072*
H16B	0.0702	0.8923	0.1324	0.072*
H16C	0.1097	0.7911	0.0559	0.072*
H2N	0.0908 (14)	0.948 (3)	0.3166 (11)	0.058*
H4N	-0.2329 (15)	0.450 (3)	0.0380 (11)	0.058*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0295 (5)	0.0439 (6)	0.0448 (6)	0.0013 (4)	-0.0031 (4)	0.0031 (5)

O2	0.0412 (6)	0.0510 (7)	0.0590 (7)	0.0083 (5)	-0.0059 (5)	0.0141 (6)
O3	0.0400 (6)	0.0337 (5)	0.0503 (6)	-0.0062 (4)	-0.0005 (5)	0.0064 (5)
O4	0.0461 (7)	0.0435 (6)	0.0568 (7)	0.0067 (5)	-0.0016 (5)	0.0123 (5)
N1	0.0365 (7)	0.0427 (7)	0.0533 (8)	-0.0043 (6)	0.0002 (6)	0.0018 (6)
N2	0.0455 (8)	0.0362 (7)	0.0518 (8)	-0.0023 (6)	0.0037 (6)	0.0039 (6)
N3	0.0368 (7)	0.0496 (8)	0.0561 (9)	-0.0102 (6)	-0.0017 (6)	0.0001 (7)
N4	0.0332 (7)	0.0567 (9)	0.0504 (8)	-0.0009 (6)	-0.0051 (6)	-0.0039 (7)
C1	0.0344 (8)	0.0406 (8)	0.0361 (8)	0.0032 (6)	0.0014 (6)	0.0002 (6)
C2	0.0318 (7)	0.0398 (8)	0.0409 (8)	-0.0023 (6)	0.0028 (6)	0.0035 (6)
C3	0.0289 (7)	0.0405 (8)	0.0330 (7)	0.0008 (6)	0.0037 (6)	-0.0025 (6)
C4	0.0305 (7)	0.0368 (7)	0.0323 (7)	0.0015 (6)	0.0022 (6)	-0.0013 (6)
C5	0.0307 (7)	0.0387 (8)	0.0352 (8)	0.0001 (6)	0.0009 (6)	-0.0020 (6)
C6	0.0387 (8)	0.0369 (8)	0.0371 (8)	0.0023 (6)	0.0044 (6)	-0.0008 (6)
C7	0.0493 (10)	0.0450 (9)	0.0510 (10)	0.0092 (7)	0.0017 (8)	0.0084 (7)
C8	0.0304 (8)	0.0539 (10)	0.0564 (10)	-0.0017 (7)	-0.0001 (7)	0.0063 (8)
C9	0.0339 (8)	0.0348 (8)	0.0404 (8)	0.0027 (6)	0.0035 (6)	-0.0002 (6)
C10	0.0278 (7)	0.0380 (8)	0.0443 (8)	-0.0033 (6)	0.0021 (6)	-0.0009 (6)
C11	0.0333 (7)	0.0318 (7)	0.0353 (7)	-0.0013 (6)	0.0067 (6)	-0.0027 (6)
C12	0.0325 (7)	0.0327 (7)	0.0351 (8)	0.0012 (6)	0.0051 (6)	-0.0025 (6)
C13	0.0333 (8)	0.0367 (8)	0.0386 (8)	-0.0028 (6)	0.0023 (6)	-0.0015 (6)
C14	0.0351 (8)	0.0434 (8)	0.0352 (8)	0.0053 (7)	0.0022 (6)	-0.0046 (6)
C15	0.0498 (10)	0.0539 (10)	0.0477 (10)	0.0141 (8)	-0.0013 (8)	0.0043 (8)
C16	0.0486 (10)	0.0381 (9)	0.0577 (10)	-0.0075 (7)	0.0059 (8)	0.0065 (7)

Geometric parameters (Å, °)

O1—C5	1.3603 (17)	C6—C7	1.485 (2)
O1—C1	1.3768 (18)	C7—H7A	0.9600
O2—C1	1.2165 (18)	C7—H7B	0.9600
O3—C13	1.3609 (18)	C7—H7C	0.9600
O3—C9	1.3828 (18)	C8—H8A	0.9600
O4—C9	1.2122 (18)	C8—H8B	0.9600
N1—C5	1.3094 (19)	C8—H8C	0.9600
N1—N2	1.3695 (19)	C9—C10	1.436 (2)
N2—C6	1.337 (2)	C10—C11	1.350 (2)
N2—H2N	0.912 (19)	C10—H10	0.9300
N3—C13	1.3076 (19)	C11—C12	1.4361 (19)
N3—N4	1.370 (2)	C11—C16	1.495 (2)
N4—C14	1.338 (2)	C12—C14	1.393 (2)
N4—H4N	0.91 (2)	C12—C13	1.394 (2)
C1—C2	1.439 (2)	C14—C15	1.489 (2)
C2—C3	1.351 (2)	C15—H15A	0.9600
C2—H2	0.9300	C15—H15B	0.9600
C3—C4	1.428 (2)	C15—H15C	0.9600
C3—C8	1.495 (2)	C16—H16A	0.9600
C4—C6	1.392 (2)	C16—H16B	0.9600
C4—C5	1.3998 (19)	C16—H16C	0.9600

C5—O1—C1	117.54 (11)	C3—C8—H8B	109.5
C13—O3—C9	117.92 (11)	H8A—C8—H8B	109.5
C5—N1—N2	101.55 (12)	C3—C8—H8C	109.5
C6—N2—N1	114.76 (13)	H8A—C8—H8C	109.5
C6—N2—H2N	125.6 (12)	H8B—C8—H8C	109.5
N1—N2—H2N	119.7 (12)	O4—C9—O3	114.99 (13)
C13—N3—N4	101.29 (13)	O4—C9—C10	126.33 (14)
C14—N4—N3	114.69 (13)	O3—C9—C10	118.68 (13)
C14—N4—H4N	126.9 (12)	C11—C10—C9	123.95 (13)
N3—N4—H4N	118.4 (12)	C11—C10—H10	118.0
O2—C1—O1	116.08 (13)	C9—C10—H10	118.0
O2—C1—C2	124.78 (15)	C10—C11—C12	116.31 (13)
O1—C1—C2	119.13 (13)	C10—C11—C16	122.29 (14)
C3—C2—C1	123.74 (14)	C12—C11—C16	121.39 (13)
C3—C2—H2	118.1	C14—C12—C13	103.33 (13)
C1—C2—H2	118.1	C14—C12—C11	137.69 (14)
C2—C3—C4	116.33 (13)	C13—C12—C11	118.98 (13)
C2—C3—C8	122.12 (14)	N3—C13—O3	120.62 (13)
C4—C3—C8	121.55 (13)	N3—C13—C12	115.28 (14)
C6—C4—C5	103.48 (13)	O3—C13—C12	124.10 (13)
C6—C4—C3	137.45 (14)	N4—C14—C12	105.41 (13)
C5—C4—C3	119.05 (13)	N4—C14—C15	122.24 (14)
N1—C5—O1	120.99 (13)	C12—C14—C15	132.35 (15)
N1—C5—C4	114.79 (13)	C14—C15—H15A	109.5
O1—C5—C4	124.21 (13)	C14—C15—H15B	109.5
N2—C6—C4	105.43 (13)	H15A—C15—H15B	109.5
N2—C6—C7	122.57 (14)	C14—C15—H15C	109.5
C4—C6—C7	132.00 (14)	H15A—C15—H15C	109.5
C6—C7—H7A	109.5	H15B—C15—H15C	109.5
C6—C7—H7B	109.5	C11—C16—H16A	109.5
H7A—C7—H7B	109.5	C11—C16—H16B	109.5
C6—C7—H7C	109.5	H16A—C16—H16B	109.5
H7A—C7—H7C	109.5	C11—C16—H16C	109.5
H7B—C7—H7C	109.5	H16A—C16—H16C	109.5
C3—C8—H8A	109.5	H16B—C16—H16C	109.5
C5—N1—N2—C6	-0.04 (18)	C3—C4—C6—C7	-0.6 (3)
C13—N3—N4—C14	0.15 (18)	C13—O3—C9—O4	177.73 (13)
C5—O1—C1—O2	179.84 (13)	C13—O3—C9—C10	-1.88 (19)
C5—O1—C1—C2	-0.76 (19)	O4—C9—C10—C11	-176.82 (15)
O2—C1—C2—C3	-179.65 (15)	O3—C9—C10—C11	2.8 (2)
O1—C1—C2—C3	1.0 (2)	C9—C10—C11—C12	-1.7 (2)
C1—C2—C3—C4	-0.8 (2)	C9—C10—C11—C16	178.13 (14)
C1—C2—C3—C8	179.23 (14)	C10—C11—C12—C14	-179.24 (16)
C2—C3—C4—C6	-177.72 (16)	C16—C11—C12—C14	0.9 (3)
C8—C3—C4—C6	2.3 (3)	C10—C11—C12—C13	0.0 (2)
C2—C3—C4—C5	0.4 (2)	C16—C11—C12—C13	-179.86 (14)
C8—C3—C4—C5	-179.65 (14)	N4—N3—C13—O3	179.70 (13)

N2—N1—C5—O1	-178.64 (13)	N4—N3—C13—C12	-0.08 (17)
N2—N1—C5—C4	0.29 (17)	C9—O3—C13—N3	-179.50 (13)
C1—O1—C5—N1	179.23 (13)	C9—O3—C13—C12	0.3 (2)
C1—O1—C5—C4	0.4 (2)	C14—C12—C13—N3	-0.02 (17)
C6—C4—C5—N1	-0.42 (17)	C11—C12—C13—N3	-179.50 (13)
C3—C4—C5—N1	-179.09 (13)	C14—C12—C13—O3	-179.79 (13)
C6—C4—C5—O1	178.48 (13)	C11—C12—C13—O3	0.7 (2)
C3—C4—C5—O1	-0.2 (2)	N3—N4—C14—C12	-0.16 (18)
N1—N2—C6—C4	-0.21 (18)	N3—N4—C14—C15	-179.68 (14)
N1—N2—C6—C7	179.12 (14)	C13—C12—C14—N4	0.10 (15)
C5—C4—C6—N2	0.35 (15)	C11—C12—C14—N4	179.43 (16)
C3—C4—C6—N2	178.62 (16)	C13—C12—C14—C15	179.54 (16)
C5—C4—C6—C7	-178.89 (16)	C11—C12—C14—C15	-1.1 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N4—H4N \cdots O2 ⁱ	0.91 (2)	1.91 (2)	2.7860 (17)	160.9 (17)
N2—H2N \cdots O4 ⁱⁱ	0.912 (19)	1.984 (19)	2.8872 (18)	170.4 (17)
C2—H2 \cdots O2 ⁱⁱⁱ	0.93	2.49	3.4082 (19)	171
C7—H7A \cdots O3 ⁱⁱ	0.96	2.54	3.458 (2)	159
C10—H10 \cdots O4 ^{iv}	0.93	2.45	3.3563 (18)	164

Symmetry codes: (i) $x-1/2, -y+1/2, z-1/2$; (ii) $x, y+1, z$; (iii) $-x, -y, -z+1$; (iv) $-x+1/2, y+1/2, -z+1/2$.