

## 5-Nitro-1-(prop-2-yn-1-yl)-2,3-dihydro-1*H*-1,3-benzodiazol-2-one

Younes Ouzidan,<sup>a</sup> Youssef Kandri Rodi,<sup>b</sup> Hafid Zouihri,<sup>c</sup> El Mokhtar Essassi<sup>d</sup> and Seik Weng Ng<sup>e,f\*</sup>

<sup>a</sup>Laboratoire de Chimie Organique Appliquée, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, <sup>b</sup>Laboratoire de Chimie Organique Appliquée, Faculté des Sciences et Techniques, Université Sidi Mohamed Ben Abdallah, Fès, Morocco, <sup>c</sup>CNRST Division UATRS, Angle Allal Fassi/FAR, BP 8027 Hay Riad, Rabat, Morocco, <sup>d</sup>Laboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, <sup>e</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and <sup>f</sup>Chemistry Department, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia  
Correspondence e-mail: seikweng@um.edu.my


Received 23 March 2012; accepted 26 March 2012

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.045;  $wR$  factor = 0.158; data-to-parameter ratio = 14.8.

In the two independent molecules of the title compound,  $\text{C}_{10}\text{H}_7\text{N}_3\text{O}_3$ , the nitro substituent is twisted slightly with respect to the benzodiazol fused-ring system [dihedral angles = 4.9 (3) and 8.5 (1) $^\circ$ ]. The two independent molecules are disposed about a pseudo inversion center and are held together by N–H $\cdots$ O hydrogen bonds. The supramolecular dimer is essentially planar [dihedral angle between the fused rings = 2.0 (1) $^\circ$ ]. Adjacent dimers are linked by acetylene–nitro C–H $\cdots$ O interactions, generating a ribbon motif along (110).

### Related literature

For related structures, see: Ouzidan *et al.* (2011a,b,c).



### Experimental

#### Crystal data



$M_r = 217.19$

Triclinic, $P\bar{1}$	$V = 986.20 (4)\text{ \AA}^3$
$a = 7.2541 (2)\text{ \AA}$	$Z = 4$
$b = 10.0362 (2)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 14.6793 (3)\text{ \AA}$	$\mu = 0.11\text{ mm}^{-1}$
$\alpha = 100.978 (1)^\circ$	$T = 293\text{ K}$
$\beta = 92.047 (1)^\circ$	$0.27 \times 0.22 \times 0.13\text{ mm}$
$\gamma = 109.043 (1)^\circ$	

#### Data collection

Bruker APEX DUO diffractometer	2855 reflections with $I > 2\sigma(I)$
20686 measured reflections	$R_{\text{int}} = 0.032$
4506 independent reflections	

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.158$	$\Delta\rho_{\text{max}} = 0.20\text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$
4506 reflections	
305 parameters	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1–H1 $\cdots$ O4	0.96 (2)	1.81 (3)	2.766 (2)	173 (2)
N4–H4 $\cdots$ O1	0.92 (2)	1.91 (2)	2.823 (2)	172 (2)
C4–H41 $\cdots$ O3 <sup>i</sup>	0.92 (4)	2.39 (4)	3.231 (4)	153 (3)
C14–H141 $\cdots$ O6 <sup>ii</sup>	0.88 (3)	2.51 (3)	3.383 (4)	169 (3)

Symmetry codes: (i)  $x - 1, y - 1, z$ ; (ii)  $x + 1, y + 1, z$ .

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Université Mohammed V-Agdal and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12) for supporting this study

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5859).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2010). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ouzidan, Y., Kandri Rodi, Y., Essassi, E. M., El Ammari, L., Fronczek, F. R. & Venkatraman, R. (2011a). *Acta Cryst. E67*, o669.
- Ouzidan, Y., Kandri Rodi, Y., Jasinski, J. P., Butcher, R. J., Golen, J. A. & El Ammari, L. (2011b). *Acta Cryst. E67*, o1091.
- Ouzidan, Y., Kandri Rodi, Y., Saffon, N., Essassi, E. M. & Ng, S. W. (2011c). *Acta Cryst. E67*, o558.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

# supporting information

*Acta Cryst.* (2012). E68, o1240 [https://doi.org/10.1107/S1600536812013177]

## 5-Nitro-1-(prop-2-yn-1-yl)-2,3-dihydro-1*H*-1,3-benzodiazol-2-one

**Younes Ouzidan, Youssef Kandri Rodi, Hafid Zouihri, El Mokhtar Essassi and Seik Weng Ng**

### S1. Comment

Benzodiazoles are of interest owing to their pharmacological properties. When the parent compound, benzodiazol-2-one, reacts with propargyl bromide, both amino groups are alkylated to give 1,3-bis(prop-2-ynyl)-benzodiazol-2-one (Ouzidan *et al.*, 2011*b*). The presence of an electron-withdrawing nitro group allows only one amino group to be alkylated, as noted from the reactions of 5-nitrobenzodiazol-2-onezol with *n*-octyl bromide and *n*-nonyl bromide (Ouzidan *et al.*, 2011*b*, 2011*c*). In the two independent molecules of C<sub>10</sub>H<sub>7</sub>N<sub>3</sub>O<sub>3</sub> (Scheme I), the nitro substituent is slightly bent with respect to the benzodiazol fused-ring (Fig. 2). The two are disposed about a false inversion center, and are held together by N–H···O hydrogen bonds. Adjacent dimers are linked by C–H<sub>acetylene</sub>···O<sub>nitro</sub> interactions to generate a ribbon motif (Fig. 2).

### S2. Experimental


To a mixture of 5-nitro-1*H*-benzodiazol-2(3*H*)-one (0.25 g, 1.5 mmol), potassium carbonate (0.35 g, 2.5 mmol), tetra-*n*-butylammonium bromide (0.1 g, 0.2 mmol) in DMF (15 ml) was added propargyl bromide (0.14 ml, 1.6 mmol). Stirring was continued at room temperature for 6 hours. The salt was removed by filtration and the filtrate concentrated under reduced pressure. The residue was separated by chromatography on a column of silica gel with ethylacetate/hexane; yellow crystals were obtained upon evaporation of the solvent.

### S3. Refinement


All H atoms were located in a difference map. The aromatic and methylene H-atoms were placed in calculated positions (C–H 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with U(H) set to 1.2U(C).

The amino and acetylenic H-atoms were freely refined.

The (0 0 1) reflection was omitted owing to bad disagreement.

**Figure 1**

Anisotropic displacement ellipsoid plot (Barbour, 2001) of the two independent molecules of  $C_{10}H_7N_3O_3$  molecule at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

**Figure 2**

Anisotropic displacement ellipsoid plot (Barbour, 2001) of the hydrogen-bonded ribbon motif.

### 5-Nitro-1-(prop-2-yn-1-yl)-2,3-dihydro-1*H*-1,3-benzodiazol-2-one

#### Crystal data

$C_{10}H_7N_3O_3$   
 $M_r = 217.19$   
Triclinic,  $P\bar{1}$   
Hall symbol: -P 1  
 $a = 7.2541 (2)$  Å  
 $b = 10.0362 (2)$  Å  
 $c = 14.6793 (3)$  Å  
 $\alpha = 100.978 (1)$ °  
 $\beta = 92.047 (1)$ °

$\gamma = 109.043 (1)$ °  
 $V = 986.20 (4)$  Å<sup>3</sup>  
 $Z = 4$   
 $F(000) = 448$   
 $D_x = 1.463$  Mg m<sup>-3</sup>  
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 5674 reflections  
 $\theta = 2.2\text{--}28.0$ °  
 $\mu = 0.11$  mm<sup>-1</sup>

$T = 293\text{ K}$   
Prism, yellow

$0.27 \times 0.22 \times 0.13\text{ mm}$

#### Data collection

Bruker APEX DUO  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
20686 measured reflections  
4506 independent reflections

2855 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.032$   
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.2^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -13 \rightarrow 12$   
 $l = -19 \rightarrow 19$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.158$   
 $S = 1.05$   
4506 reflections  
305 parameters  
0 restraints  
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
Hydrogen site location: inferred from neighbouring sites  
H atoms treated by a mixture of independent and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0746P)^2 + 0.2589P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.24\text{ e \AA}^{-3}$

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^* / U_{\text{eq}}$
O1	0.4863 (2)	0.28214 (16)	0.32061 (10)	0.0583 (4)
O2	0.2729 (3)	0.70175 (19)	-0.06407 (12)	0.0780 (5)
O3	0.4106 (3)	0.84006 (18)	0.06741 (13)	0.0826 (6)
O4	0.7053 (2)	0.67457 (15)	0.41638 (10)	0.0556 (4)
O5	1.0136 (4)	0.2778 (2)	0.79280 (13)	0.1056 (8)
O6	0.8109 (3)	0.13338 (17)	0.67890 (13)	0.0826 (6)
N1	0.4993 (2)	0.47954 (18)	0.25677 (11)	0.0459 (4)
N2	0.3361 (2)	0.26206 (16)	0.17380 (10)	0.0438 (4)
N3	0.3367 (2)	0.72031 (19)	0.01736 (13)	0.0535 (4)
N4	0.7143 (2)	0.48027 (17)	0.47997 (11)	0.0416 (4)
N5	0.8738 (2)	0.70106 (16)	0.55973 (10)	0.0429 (4)
N6	0.9101 (3)	0.25456 (19)	0.72058 (13)	0.0583 (5)
C1	0.4468 (3)	0.3367 (2)	0.25799 (13)	0.0447 (4)
C2	0.2539 (3)	0.1058 (2)	0.14690 (15)	0.0555 (5)
H2A	0.3007	0.0644	0.1936	0.067*
H2B	0.2988	0.0734	0.0880	0.067*
C3	0.0406 (4)	0.0549 (2)	0.13742 (15)	0.0585 (6)
C4	-0.1296 (5)	0.0194 (3)	0.1295 (2)	0.0859 (9)
C5	0.3154 (3)	0.35855 (19)	0.12192 (12)	0.0391 (4)
C6	0.2169 (3)	0.3381 (2)	0.03563 (13)	0.0450 (4)
H6	0.1471	0.2461	0.0011	0.054*
C7	0.2259 (3)	0.4598 (2)	0.00241 (13)	0.0454 (4)
H7	0.1628	0.4505	-0.0559	0.054*

C8	0.3287 (3)	0.5950 (2)	0.05579 (13)	0.0418 (4)
C9	0.4281 (3)	0.6184 (2)	0.14315 (13)	0.0428 (4)
H9	0.4961	0.7107	0.1778	0.051*
C10	0.4197 (2)	0.49734 (19)	0.17501 (12)	0.0391 (4)
C11	0.7577 (3)	0.6231 (2)	0.47796 (13)	0.0432 (4)
C12	0.9419 (3)	0.8575 (2)	0.58554 (15)	0.0552 (5)
H12A	0.8899	0.8870	0.6430	0.066*
H12B	0.8916	0.8943	0.5372	0.066*
C13	1.1548 (4)	0.9205 (2)	0.59851 (15)	0.0621 (6)
C14	1.3251 (5)	0.9698 (3)	0.6118 (2)	0.0942 (10)
C15	0.9034 (3)	0.60770 (18)	0.61264 (12)	0.0381 (4)
C16	1.0082 (3)	0.6333 (2)	0.69790 (13)	0.0452 (4)
H16	1.0756	0.7267	0.7310	0.054*
C17	1.0092 (3)	0.5145 (2)	0.73252 (13)	0.0463 (5)
H17	1.0786	0.5271	0.7898	0.056*
C18	0.9065 (3)	0.3768 (2)	0.68142 (13)	0.0432 (4)
C19	0.7999 (3)	0.34865 (19)	0.59536 (12)	0.0404 (4)
H19	0.7318	0.2552	0.5625	0.048*
C20	0.8012 (2)	0.46764 (19)	0.56177 (12)	0.0371 (4)
H1	0.574 (3)	0.554 (3)	0.3093 (17)	0.073 (7)*
H4	0.645 (3)	0.409 (2)	0.4297 (16)	0.060 (6)*
H41	-0.263 (6)	-0.002 (4)	0.122 (3)	0.138 (14)*
H141	1.454 (5)	1.011 (3)	0.621 (2)	0.118 (12)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0683 (9)	0.0531 (9)	0.0505 (8)	0.0149 (7)	-0.0098 (7)	0.0175 (7)
O2	0.0971 (13)	0.0713 (11)	0.0680 (11)	0.0265 (10)	-0.0142 (9)	0.0289 (9)
O3	0.1019 (14)	0.0448 (10)	0.0931 (13)	0.0164 (9)	-0.0181 (10)	0.0160 (9)
O4	0.0672 (9)	0.0501 (8)	0.0493 (8)	0.0188 (7)	-0.0090 (7)	0.0151 (7)
O5	0.169 (2)	0.0641 (12)	0.0750 (12)	0.0367 (12)	-0.0557 (13)	0.0130 (10)
O6	0.1049 (14)	0.0406 (9)	0.0919 (13)	0.0138 (9)	-0.0253 (10)	0.0156 (9)
N1	0.0484 (9)	0.0429 (9)	0.0398 (9)	0.0081 (7)	-0.0054 (7)	0.0082 (7)
N2	0.0512 (9)	0.0392 (9)	0.0398 (8)	0.0157 (7)	0.0000 (7)	0.0060 (7)
N3	0.0492 (10)	0.0468 (10)	0.0649 (12)	0.0145 (8)	0.0006 (8)	0.0172 (9)
N4	0.0445 (9)	0.0383 (9)	0.0376 (8)	0.0107 (7)	-0.0035 (7)	0.0050 (7)
N5	0.0526 (9)	0.0351 (8)	0.0396 (8)	0.0146 (7)	0.0018 (7)	0.0055 (7)
N6	0.0762 (12)	0.0428 (10)	0.0556 (11)	0.0211 (9)	-0.0062 (9)	0.0104 (8)
C1	0.0451 (10)	0.0465 (11)	0.0416 (10)	0.0147 (8)	0.0014 (8)	0.0092 (9)
C2	0.0770 (15)	0.0389 (11)	0.0518 (12)	0.0243 (10)	0.0004 (10)	0.0051 (9)
C3	0.0795 (16)	0.0329 (10)	0.0503 (12)	0.0084 (10)	-0.0016 (11)	-0.0006 (9)
C4	0.082 (2)	0.0561 (16)	0.088 (2)	-0.0025 (14)	0.0001 (16)	-0.0130 (14)
C5	0.0418 (9)	0.0385 (10)	0.0380 (9)	0.0159 (8)	0.0051 (7)	0.0064 (8)
C6	0.0519 (11)	0.0399 (10)	0.0384 (10)	0.0142 (8)	-0.0013 (8)	0.0008 (8)
C7	0.0472 (10)	0.0484 (11)	0.0404 (10)	0.0175 (9)	-0.0019 (8)	0.0081 (8)
C8	0.0400 (9)	0.0437 (10)	0.0446 (10)	0.0156 (8)	0.0057 (8)	0.0131 (8)
C9	0.0402 (10)	0.0371 (10)	0.0460 (11)	0.0087 (7)	0.0039 (8)	0.0046 (8)

C10	0.0356 (9)	0.0422 (10)	0.0363 (9)	0.0106 (7)	0.0031 (7)	0.0059 (8)
C11	0.0428 (10)	0.0440 (11)	0.0422 (10)	0.0144 (8)	0.0033 (8)	0.0089 (8)
C12	0.0725 (14)	0.0361 (11)	0.0553 (12)	0.0177 (10)	0.0026 (10)	0.0076 (9)
C13	0.0790 (17)	0.0394 (11)	0.0541 (13)	0.0046 (11)	-0.0032 (11)	0.0067 (10)
C14	0.081 (2)	0.0721 (19)	0.096 (2)	-0.0106 (16)	-0.0107 (17)	0.0092 (16)
C15	0.0429 (9)	0.0350 (9)	0.0360 (9)	0.0137 (7)	0.0070 (7)	0.0054 (7)
C16	0.0537 (11)	0.0372 (10)	0.0379 (10)	0.0119 (8)	-0.0001 (8)	-0.0008 (8)
C17	0.0558 (11)	0.0465 (11)	0.0344 (9)	0.0180 (9)	-0.0018 (8)	0.0032 (8)
C18	0.0498 (10)	0.0393 (10)	0.0415 (10)	0.0173 (8)	0.0039 (8)	0.0075 (8)
C19	0.0432 (10)	0.0337 (9)	0.0401 (10)	0.0111 (7)	0.0020 (8)	0.0021 (7)
C20	0.0365 (9)	0.0399 (10)	0.0337 (9)	0.0138 (7)	0.0040 (7)	0.0036 (7)

*Geometric parameters (Å, °)*

O1—C1	1.226 (2)	C4—H41	0.92 (4)
O2—N3	1.224 (2)	C5—C6	1.377 (2)
O3—N3	1.219 (2)	C5—C10	1.406 (2)
O4—C11	1.229 (2)	C6—C7	1.382 (3)
O5—N6	1.215 (2)	C6—H6	0.9300
O6—N6	1.219 (2)	C7—C8	1.380 (3)
N1—C1	1.362 (2)	C7—H7	0.9300
N1—C10	1.380 (2)	C8—C9	1.389 (3)
N1—H1	0.96 (2)	C9—C10	1.367 (3)
N2—C5	1.381 (2)	C9—H9	0.9300
N2—C1	1.385 (2)	C12—C13	1.455 (3)
N2—C2	1.452 (2)	C12—H12A	0.9700
N3—C8	1.459 (2)	C12—H12B	0.9700
N4—C11	1.370 (2)	C13—C14	1.164 (4)
N4—C20	1.383 (2)	C14—H141	0.88 (3)
N4—H4	0.92 (2)	C15—C16	1.375 (2)
N5—C11	1.379 (2)	C15—C20	1.401 (2)
N5—C15	1.387 (2)	C16—C17	1.384 (3)
N5—C12	1.452 (2)	C16—H16	0.9300
N6—C18	1.457 (2)	C17—C18	1.387 (3)
C2—C3	1.455 (3)	C17—H17	0.9300
C2—H2A	0.9700	C18—C19	1.388 (3)
C2—H2B	0.9700	C19—C20	1.373 (2)
C3—C4	1.162 (4)	C19—H19	0.9300
C1—N1—C10	110.23 (16)	C7—C8—C9	123.72 (17)
C1—N1—H1	121.9 (14)	C7—C8—N3	118.08 (17)
C10—N1—H1	127.7 (14)	C9—C8—N3	118.18 (17)
C5—N2—C1	109.67 (15)	C10—C9—C8	115.85 (17)
C5—N2—C2	126.40 (16)	C10—C9—H9	122.1
C1—N2—C2	123.92 (16)	C8—C9—H9	122.1
O3—N3—O2	122.50 (18)	C9—C10—N1	131.69 (17)
O3—N3—C8	118.46 (18)	C9—C10—C5	121.39 (16)
O2—N3—C8	119.03 (18)	N1—C10—C5	106.92 (16)

C11—N4—C20	109.80 (15)	O4—C11—N4	127.80 (18)
C11—N4—H4	120.7 (13)	O4—C11—N5	125.54 (18)
C20—N4—H4	129.3 (14)	N4—C11—N5	106.66 (16)
C11—N5—C15	109.80 (15)	N5—C12—C13	112.49 (17)
C11—N5—C12	123.13 (16)	N5—C12—H12A	109.1
C15—N5—C12	126.98 (16)	C13—C12—H12A	109.1
O5—N6—O6	122.84 (18)	N5—C12—H12B	109.1
O5—N6—C18	118.59 (18)	C13—C12—H12B	109.1
O6—N6—C18	118.57 (17)	H12A—C12—H12B	107.8
O1—C1—N1	127.84 (18)	C14—C13—C12	177.7 (3)
O1—C1—N2	125.59 (18)	C13—C14—H14I	177 (2)
N1—C1—N2	106.57 (16)	C16—C15—N5	131.48 (17)
N2—C2—C3	111.72 (17)	C16—C15—C20	122.06 (16)
N2—C2—H2A	109.3	N5—C15—C20	106.46 (15)
C3—C2—H2A	109.3	C15—C16—C17	117.40 (17)
N2—C2—H2B	109.3	C15—C16—H16	121.3
C3—C2—H2B	109.3	C17—C16—H16	121.3
H2A—C2—H2B	107.9	C16—C17—C18	119.75 (17)
C4—C3—C2	177.4 (3)	C16—C17—H17	120.1
C3—C4—H4I	176 (2)	C18—C17—H17	120.1
C6—C5—N2	131.73 (17)	C19—C18—C17	123.75 (17)
C6—C5—C10	121.69 (17)	C19—C18—N6	118.16 (17)
N2—C5—C10	106.58 (15)	C17—C18—N6	118.09 (17)
C5—C6—C7	117.37 (17)	C20—C19—C18	115.76 (16)
C5—C6—H6	121.3	C20—C19—H19	122.1
C7—C6—H6	121.3	C18—C19—H19	122.1
C8—C7—C6	119.97 (17)	C19—C20—N4	131.45 (16)
C8—C7—H7	120.0	C19—C20—C15	121.28 (16)
C6—C7—H7	120.0	N4—C20—C15	107.28 (15)
C10—N1—C1—O1	-177.65 (19)	C20—N4—C11—O4	-179.86 (18)
C10—N1—C1—N2	1.7 (2)	C20—N4—C11—N5	0.1 (2)
C5—N2—C1—O1	177.68 (18)	C15—N5—C11—O4	-179.94 (18)
C2—N2—C1—O1	-1.5 (3)	C12—N5—C11—O4	3.2 (3)
C5—N2—C1—N1	-1.7 (2)	C15—N5—C11—N4	0.1 (2)
C2—N2—C1—N1	179.04 (16)	C12—N5—C11—N4	-176.74 (17)
C5—N2—C2—C3	-64.0 (2)	C11—N5—C12—C13	-120.6 (2)
C1—N2—C2—C3	115.0 (2)	C15—N5—C12—C13	63.0 (3)
C1—N2—C5—C6	-178.67 (19)	C11—N5—C15—C16	179.53 (19)
C2—N2—C5—C6	0.5 (3)	C12—N5—C15—C16	-3.7 (3)
C1—N2—C5—C10	1.1 (2)	C11—N5—C15—C20	-0.27 (19)
C2—N2—C5—C10	-179.73 (17)	C12—N5—C15—C20	176.45 (17)
N2—C5—C6—C7	-179.67 (18)	N5—C15—C16—C17	-179.89 (18)
C10—C5—C6—C7	0.6 (3)	C20—C15—C16—C17	-0.1 (3)
C5—C6—C7—C8	-0.8 (3)	C15—C16—C17—C18	-0.2 (3)
C6—C7—C8—C9	0.4 (3)	C16—C17—C18—C19	0.1 (3)
C6—C7—C8—N3	179.11 (16)	C16—C17—C18—N6	-179.55 (17)
O3—N3—C8—C7	172.97 (18)	O5—N6—C18—C19	175.1 (2)

O2—N3—C8—C7	−8.4 (3)	O6—N6—C18—C19	−4.0 (3)
O3—N3—C8—C9	−8.3 (3)	O5—N6—C18—C17	−5.2 (3)
O2—N3—C8—C9	170.30 (18)	O6—N6—C18—C17	175.7 (2)
C7—C8—C9—C10	0.1 (3)	C17—C18—C19—C20	0.2 (3)
N3—C8—C9—C10	−178.55 (15)	N6—C18—C19—C20	179.91 (16)
C8—C9—C10—N1	179.90 (18)	C18—C19—C20—N4	179.50 (18)
C8—C9—C10—C5	−0.3 (2)	C18—C19—C20—C15	−0.5 (2)
C1—N1—C10—C9	178.72 (19)	C11—N4—C20—C19	179.75 (18)
C1—N1—C10—C5	−1.1 (2)	C11—N4—C20—C15	−0.23 (19)
C6—C5—C10—C9	−0.1 (3)	C16—C15—C20—C19	0.5 (3)
N2—C5—C10—C9	−179.84 (15)	N5—C15—C20—C19	−179.68 (15)
C6—C5—C10—N1	179.78 (16)	C16—C15—C20—N4	−179.52 (16)
N2—C5—C10—N1	0.01 (19)	N5—C15—C20—N4	0.30 (18)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O4	0.96 (2)	1.81 (3)	2.766 (2)	173 (2)
N4—H4···O1	0.92 (2)	1.91 (2)	2.823 (2)	172 (2)
C4—H41···O3 <sup>i</sup>	0.92 (4)	2.39 (4)	3.231 (4)	153 (3)
C14—H141···O6 <sup>ii</sup>	0.88 (3)	2.51 (3)	3.383 (4)	169 (3)

Symmetry codes: (i)  $x-1, y-1, z$ ; (ii)  $x+1, y+1, z$ .