

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## *trans*-Bis[(2-bromophenyl)diphenylphosphane- $\kappa$ P]carbonylchlorido-rhodium(I)

Frederick P. Malan, Rehana Malgas-Enus and Reinout Meijboom\*

Research Centre for Synthesis and Catalysis, Department of Chemistry, University of Johannesburg, PO Box 524, Auckland Park 2006, Johannesburg, South Africa  
Correspondence e-mail: rmeijboom@uj.ac.za


Received 9 March 2012; accepted 22 March 2012

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in main residue;  $R$  factor = 0.027;  $wR$  factor = 0.063; data-to-parameter ratio = 18.8.

The title compound, *trans*-[RhCl(C<sub>18</sub>H<sub>14</sub>BrP)<sub>2</sub>(CO)], has a slightly disordered square-planar geometry with the Rh ion<sup>I</sup> situated on an inversion the centre and carbonyl-chloride disorder observed as a result of the crystallographic inversion symmetry. Selected geometric parameters include: Rh–P = 2.3430 (8) Å, Rh–Cl = 2.434 (3) Å, Rh–C = 1.722 (8) Å, P–Rh–P = 180.00 (3)°, P–Rh–Cl = 95.40 (7)°, 84.60 (7)° and Rh–C–O = 177.9 (8)°.

### Related literature

For background to Vaska-type complexes, see: Roodt *et al.* (2003); Lamb *et al.* (2009); Vaska & Di Luzio (1961). For related complexes, see: Burgoyne *et al.* (2010); Makhoba *et al.* (2011); Meijboom (2011); Meijboom *et al.* (2004); Otto *et al.* (2000); Otto & Roodt (2004); Chen *et al.* (1991); Kemp *et al.* (1995).



### Experimental

#### Crystal data

[RhCl(C<sub>18</sub>H<sub>14</sub>BrP)<sub>2</sub>(CO)]  
 $M_r = 848.71$   
Monoclinic,  $P2_1/n$   
 $a = 9.3250$  (5) Å  
 $b = 17.041$  (1) Å  
 $c = 10.8880$  (6) Å  
 $\beta = 111.229$  (1)°  
 $V = 1612.77$  (16) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 3.22$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.23 \times 0.13 \times 0.11$  mm

#### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2005)  
 $T_{\min} = 0.618$ ,  $T_{\max} = 0.697$   
15235 measured reflections  
4032 independent reflections  
3531 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$   
 $wR(F^2) = 0.063$   
 $S = 1.11$   
4032 reflections  
214 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.67$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.81$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008) and DIAMOND (Brandenburg & Putz (2005)); software used to prepare material for publication: SHELXTL.

Financial assistance from the South African National Research Foundation (SA NRF), the Next Generation Scholarship (NGS) in partnership with the University of Johannesburg (UJ), the Research Fund of the University of Johannesburg, TESP and SASOL is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HP2033).

### References

- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). *APEX2* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burgoyne, A. R., Meijboom, R., Muller, A. & Omondi, B. O. (2010). *Acta Cryst. E* **66**, m1380–m1381.
- Chen, Y.-J., Wang, J.-C. & Wang, Yu (1991). *Acta Cryst. C* **47**, 2441–2442.
- Kemp, G., Roodt, A. & Purcell, W. (1995). *Rhodium Express*, pp. 21–26.
- Lamb, G. W., Law, D. S., Slawin, A. M. Z. & Clarke, M. L. (2009). *Inorg. Chim. Acta*, **362**, 4263–4267.
- Makhoba, S., Muller, A., Meijboom, R. & Omondi, B. (2011). *Acta Cryst. E* **67**, m1286–m1287.
- Meijboom, R. (2011). *Acta Cryst. E* **67**, m1438.
- Meijboom, R., Muller, A. & Roodt, A. (2004). *Acta Cryst. E* **60**, m1071–m1073.
- Otto, S. & Roodt, A. (2004). *Inorg. Chim. Acta*, **357**, 1–10.
- Otto, S., Roodt, A. & Smith, J. (2000). *Inorg. Chim. Acta*, **303**, 295–299.
- Roodt, A., Otto, S. & Steyl, G. (2003). *Coord. Chem. Rev.* **245**, 121–137.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Vaska, L. & Di Luzio, J. W. (1961). *J. Am. Chem. Soc.* **83**, 2784–2785.

## supporting information

*Acta Cryst.* (2012). E68, m510 [https://doi.org/10.1107/S1600536812012421]

***trans*-Bis[(2-bromophenyl)diphenylphosphane- $\kappa$ P]carbonylchloridorhodium(I)****Frederick P. Malan, Rehana Malgas-Enus and Reinout Meijboom****S1. Comment**

A vast range of different Vaska-type complexes have been synthesized, reported and spectroscopically studied (Roodt *et al.*, 2003; Lamb *et al.*; 2009) since the synthesis and correct formulation of the original Vaska complex, *trans*-[Ir(CO)Cl(C<sub>18</sub>H<sub>15</sub>P)<sub>2</sub>], by Vaska & Di Luzio (1961). This class of symmetrical square-planar complexes (including Rh, Ir, Pd and Pt) usually crystallizes with the metal atom on a crystallographic inversion centre, resulting in a disordered packing arrangement (Chen *et al.*, 1991; Otto *et al.*, 2000; Otto & Roodt, 2004; Meijboom *et al.*, 2004). The title compound serves as yet another complex to add to the Vaska's complex range with varying Group 5 ligand systems possessing different stereoelectronic properties.

In the title compound, the Rh atom lies at the centre of a slightly distorted square-planar geometric arrangement. The Rh atom crystallizes on a centre of symmetry, a crystallographic inversion centre, and has the carbonyl and chloro- ligands disordered at a 0.5:0.5 ratio. The stereoelectronic property of the phosphane with the bromo-functionality is indicated by the smaller O1—Rh1—P1 angle (85.39 (19)°), which is translated through symmetry to the inverted side of the molecule.


Selected spectroscopic data of the current compound is comparable to other similar complexes reported previously by Roodt *et al.* (2003) and Otto & Roodt (2004). However, the interesting difference in the magnitude of  $\nu(\text{CO})$  for the solid and solution (in DCM) states of the title compound is ascribed to the packing of the molecules, which slightly distorts the Rh C—O angle. This effect was previously observed and reported for a polymorph of *trans*-[Rh(CO)Cl{PPh<sub>3</sub>}<sub>2</sub>] (Kemp *et al.*, 1995).

**S2. Experimental**

A solution of dichlorotetracarbonylchloridorhodium (0.050 g, 0.13 mmol) in acetone (3 cm<sup>3</sup>) was slowly added to a solution of the phosphane, C<sub>18</sub>H<sub>14</sub>BrP (0.176 g, 0.51 mmol) in acetone (3 cm<sup>3</sup>) at room temperature, after which the mixture was left to crystallize. Slow evaporation of the solvent afforded the title compound as yellow crystals. Spectroscopic analysis: <sup>31</sup>P{H} NMR (CDCl<sub>3</sub>, 162.0 MHz, p.p.m.): 33.1[d, <sup>1</sup>J(Rh—P) = 132.8 Hz, 2P]; IR  $\nu(\text{CO})$ : 1950.8 cm<sup>-1</sup>; (CD<sub>2</sub>Cl<sub>2</sub>)  $\nu(\text{CO})$ : 1973.2 cm<sup>-1</sup>.

**S3. Refinement**

The aromatic H atoms were placed in geometrically idealized positions (C—H = 0.93 Å) and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for all H atoms. The highest calculated residual electron density is 0.66 e.Å<sup>-3</sup> at 1.597 Å from C15, which bears no physical meaning.



**Figure 1**

The structure of the title compound, showing 50% probability displacement ellipsoids. All rings have been numbered in the same, systematic manner. H atoms are depicted by arbitrary size spheres. Hashed atoms are generated by symmetry ( $-x + 1, -y, -z + 2$ ).

***trans*-Bis[(2-bromophenyl)diphenylphosphane- $\kappa$ P]carbonylchloridorhodium(I)**

*Crystal data*

[RhCl(C<sub>18</sub>H<sub>14</sub>BrP)<sub>2</sub>(CO)]

$M_r = 848.71$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1n$

$a = 9.3250$  (5) Å

$b = 17.041$  (1) Å

$c = 10.8880$  (6) Å

$\beta = 111.229$  (1)°

$V = 1612.77$  (16) Å<sup>3</sup>

$Z = 2$

$F(000) = 840$

$D_x = 1.748$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71069$  Å

Cell parameters from 6197 reflections

$\theta = 2.4$ – $28.3$ °

$\mu = 3.22$  mm<sup>-1</sup>

$T = 100$  K

Rectangular, yellow

$0.23 \times 0.13 \times 0.11$  mm

*Data collection*

Bruker APEXII CCD  
diffractometer

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2005)

$T_{\min} = 0.618$ ,  $T_{\max} = 0.697$

15235 measured reflections

4032 independent reflections

3531 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 28.4$ °,  $\theta_{\min} = 2.3$ °

$h = -12 \rightarrow 12$

$k = -22 \rightarrow 22$

$l = -14 \rightarrow 14$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.027$  $wR(F^2) = 0.063$  $S = 1.11$ 

4032 reflections

214 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0246P)^2 + 1.6443P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.67 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.81 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Rh1	0.5	0	1	0.01378 (7)	
C1	0.5773 (7)	-0.0478 (4)	0.8993 (6)	0.0178 (11)	0.5
O1	0.6340 (10)	-0.0800 (5)	0.8343 (9)	0.0236 (17)	0.5
Cl1	0.6037 (3)	-0.06666 (16)	0.8527 (3)	0.0174 (4)	0.5
P1	0.50550 (6)	0.11189 (4)	0.87674 (5)	0.01442 (12)	
Br1	0.13942 (3)	0.061590 (16)	0.75242 (2)	0.02135 (7)	
C7	0.4960 (3)	0.09681 (15)	0.6146 (2)	0.0189 (5)	
H7	0.6006	0.1083	0.6484	0.023*	
C2	0.4152 (3)	0.09234 (14)	0.7000 (2)	0.0156 (4)	
C8	0.4165 (3)	0.20586 (14)	0.8906 (2)	0.0164 (5)	
C12	0.3697 (3)	0.30670 (16)	1.0277 (3)	0.0269 (6)	
H12	0.379	0.3248	1.1108	0.032*	
C14	0.7017 (3)	0.14283 (14)	0.9010 (2)	0.0174 (5)	
C19	0.8252 (3)	0.09824 (16)	0.9810 (2)	0.0237 (5)	
H19	0.8069	0.0527	1.02	0.028*	
C15	0.7317 (3)	0.21089 (16)	0.8434 (3)	0.0255 (5)	
H15	0.6504	0.2418	0.7908	0.031*	
C18	0.9756 (3)	0.12109 (19)	1.0032 (3)	0.0314 (6)	
H18	1.0574	0.0914	1.0582	0.038*	
C11	0.2947 (3)	0.35167 (16)	0.9183 (3)	0.0263 (6)	
H11	0.2529	0.3999	0.9275	0.032*	
C4	0.1846 (3)	0.06291 (15)	0.5096 (2)	0.0215 (5)	
H4	0.0797	0.0523	0.475	0.026*	
C6	0.4238 (3)	0.08449 (16)	0.4807 (2)	0.0231 (5)	
H6	0.4803	0.0869	0.4259	0.028*	

C17	1.0033 (3)	0.18774 (18)	0.9436 (3)	0.0326 (7)
H17	1.1039	0.2025	0.957	0.039*
C9	0.3404 (3)	0.25267 (16)	0.7805 (2)	0.0233 (5)
H9	0.3292	0.2348	0.6968	0.028*
C3	0.2589 (3)	0.07322 (14)	0.6440 (2)	0.0169 (5)
C5	0.2675 (3)	0.06855 (15)	0.4277 (2)	0.0234 (5)
H5	0.2187	0.0617	0.3374	0.028*
C13	0.4314 (3)	0.23448 (15)	1.0149 (2)	0.0220 (5)
H13	0.4831	0.2049	1.0897	0.026*
C10	0.2816 (3)	0.32529 (16)	0.7949 (3)	0.0274 (6)
H10	0.2332	0.3563	0.7213	0.033*
C16	0.8815 (3)	0.23287 (17)	0.8639 (3)	0.0306 (6)
H16	0.9005	0.278	0.8241	0.037*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Rh1	0.01221 (11)	0.01690 (13)	0.01105 (11)	−0.00151 (9)	0.00278 (9)	0.00207 (10)
C1	0.021 (3)	0.017 (3)	0.015 (3)	0.000 (2)	0.007 (2)	0.004 (2)
O1	0.024 (4)	0.027 (4)	0.026 (3)	0.008 (2)	0.017 (2)	0.001 (2)
Cl1	0.0204 (12)	0.0177 (11)	0.0171 (11)	0.0033 (8)	0.0105 (7)	0.0011 (8)
P1	0.0130 (3)	0.0175 (3)	0.0124 (3)	−0.0011 (2)	0.0041 (2)	0.0022 (2)
Br1	0.01558 (11)	0.03052 (15)	0.01759 (12)	−0.00259 (9)	0.00558 (9)	0.00252 (10)
C7	0.0188 (11)	0.0208 (12)	0.0185 (11)	0.0022 (9)	0.0085 (9)	0.0024 (9)
C2	0.0193 (11)	0.0150 (11)	0.0129 (10)	0.0016 (9)	0.0063 (9)	0.0014 (9)
C8	0.0139 (10)	0.0191 (12)	0.0164 (11)	−0.0030 (9)	0.0059 (9)	−0.0013 (9)
C12	0.0326 (14)	0.0271 (14)	0.0274 (13)	−0.0124 (11)	0.0187 (12)	−0.0121 (11)
C14	0.0161 (10)	0.0194 (12)	0.0178 (11)	−0.0036 (9)	0.0074 (9)	−0.0034 (9)
C19	0.0190 (11)	0.0281 (14)	0.0228 (12)	−0.0024 (10)	0.0062 (10)	0.0010 (11)
C15	0.0279 (13)	0.0192 (13)	0.0335 (14)	−0.0036 (10)	0.0160 (11)	−0.0032 (11)
C18	0.0156 (12)	0.0410 (17)	0.0350 (15)	−0.0016 (11)	0.0059 (11)	−0.0044 (13)
C11	0.0250 (13)	0.0190 (13)	0.0393 (15)	−0.0041 (10)	0.0171 (12)	−0.0081 (11)
C4	0.0219 (12)	0.0214 (13)	0.0171 (11)	−0.0012 (10)	0.0022 (9)	0.0012 (10)
C6	0.0304 (13)	0.0250 (14)	0.0176 (12)	0.0037 (11)	0.0132 (10)	0.0024 (10)
C17	0.0219 (13)	0.0383 (17)	0.0443 (17)	−0.0140 (12)	0.0201 (12)	−0.0196 (14)
C9	0.0255 (12)	0.0235 (13)	0.0178 (11)	0.0023 (10)	0.0041 (10)	−0.0032 (10)
C3	0.0195 (11)	0.0169 (12)	0.0151 (11)	0.0016 (9)	0.0070 (9)	0.0023 (9)
C5	0.0320 (14)	0.0247 (14)	0.0122 (11)	0.0004 (11)	0.0064 (10)	−0.0012 (10)
C13	0.0243 (12)	0.0239 (13)	0.0171 (11)	−0.0075 (10)	0.0067 (10)	−0.0039 (10)
C10	0.0265 (13)	0.0207 (13)	0.0294 (14)	0.0011 (11)	0.0033 (11)	−0.0015 (11)
C16	0.0345 (14)	0.0236 (14)	0.0440 (17)	−0.0130 (12)	0.0266 (13)	−0.0107 (12)

*Geometric parameters (Å, °)*

Rh1—C1 <sup>i</sup>	1.720 (7)	C19—C18	1.390 (3)
Rh1—C1	1.720 (7)	C19—H19	0.93
Rh1—P1	2.3429 (6)	C15—C16	1.384 (4)
Rh1—P1 <sup>i</sup>	2.3429 (6)	C15—H15	0.93

Rh1—C11 <sup>i</sup>	2.433 (3)	C18—C17	1.378 (4)
Rh1—C11	2.433 (3)	C18—H18	0.93
C1—O1	1.163 (7)	C11—C10	1.379 (4)
P1—C14	1.829 (2)	C11—H11	0.93
P1—C2	1.831 (2)	C4—C5	1.379 (4)
P1—C8	1.835 (2)	C4—C3	1.385 (3)
Br1—C3	1.904 (2)	C4—H4	0.93
C7—C6	1.383 (3)	C6—C5	1.386 (4)
C7—C2	1.394 (3)	C6—H6	0.93
C7—H7	0.93	C17—C16	1.386 (4)
C2—C3	1.399 (3)	C17—H17	0.93
C8—C13	1.397 (3)	C9—C10	1.385 (4)
C8—C9	1.400 (3)	C9—H9	0.93
C12—C11	1.375 (4)	C5—H5	0.93
C12—C13	1.387 (4)	C13—H13	0.93
C12—H12	0.93	C10—H10	0.93
C14—C19	1.391 (3)	C16—H16	0.93
C14—C15	1.394 (4)		
C1 <sup>i</sup> —Rh1—C1	180.0000 (10)	C18—C19—H19	119.7
C1 <sup>i</sup> —Rh1—P1	94.63 (19)	C14—C19—H19	119.7
C1—Rh1—P1	85.37 (19)	C16—C15—C14	120.5 (3)
C1 <sup>i</sup> —Rh1—P1 <sup>i</sup>	85.37 (19)	C16—C15—H15	119.7
C1—Rh1—P1 <sup>i</sup>	94.63 (19)	C14—C15—H15	119.7
P1—Rh1—P1 <sup>i</sup>	180	C17—C18—C19	119.9 (3)
C1 <sup>i</sup> —Rh1—C11 <sup>i</sup>	1.46 (18)	C17—C18—H18	120.1
C1—Rh1—C11 <sup>i</sup>	178.54 (19)	C19—C18—H18	120.1
P1—Rh1—C11 <sup>i</sup>	95.39 (7)	C12—C11—C10	120.0 (2)
P1 <sup>i</sup> —Rh1—C11 <sup>i</sup>	84.61 (7)	C12—C11—H11	120
C1 <sup>i</sup> —Rh1—C11	178.54 (18)	C10—C11—H11	120
C1—Rh1—C11	1.46 (19)	C5—C4—C3	119.4 (2)
P1—Rh1—C11	84.61 (7)	C5—C4—H4	120.3
P1 <sup>i</sup> —Rh1—C11	95.39 (7)	C3—C4—H4	120.3
C11 <sup>i</sup> —Rh1—C11	180.0000 (10)	C7—C6—C5	120.4 (2)
O1—C1—Rh1	177.9 (8)	C7—C6—H6	119.8
C14—P1—C2	105.04 (11)	C5—C6—H6	119.8
C14—P1—C8	101.26 (11)	C18—C17—C16	120.1 (2)
C2—P1—C8	101.29 (10)	C18—C17—H17	120
C14—P1—Rh1	112.34 (8)	C16—C17—H17	120
C2—P1—Rh1	110.92 (8)	C10—C9—C8	120.7 (2)
C8—P1—Rh1	123.96 (8)	C10—C9—H9	119.7
C6—C7—C2	121.3 (2)	C8—C9—H9	119.7
C6—C7—H7	119.3	C4—C3—C2	122.2 (2)
C2—C7—H7	119.3	C4—C3—Br1	117.41 (18)
C7—C2—C3	116.8 (2)	C2—C3—Br1	120.34 (17)
C7—C2—P1	122.46 (18)	C4—C5—C6	119.7 (2)
C3—C2—P1	120.68 (17)	C4—C5—H5	120.1
C13—C8—C9	118.2 (2)	C6—C5—H5	120.1

C13—C8—P1	119.67 (19)	C12—C13—C8	120.5 (2)
C9—C8—P1	122.05 (18)	C12—C13—H13	119.8
C11—C12—C13	120.5 (2)	C8—C13—H13	119.8
C11—C12—H12	119.8	C11—C10—C9	120.2 (3)
C13—C12—H12	119.8	C11—C10—H10	119.9
C19—C14—C15	118.7 (2)	C9—C10—H10	119.9
C19—C14—P1	119.39 (19)	C15—C16—C17	120.1 (3)
C15—C14—P1	121.86 (19)	C15—C16—H16	120
C18—C19—C14	120.7 (3)	C17—C16—H16	120
C1 <sup>i</sup> —Rh1—C1—O1	9E1 (10)	Rh1—P1—C8—C9	-141.23 (18)
P1—Rh1—C1—O1	10E1 (2)	C2—P1—C14—C19	-117.8 (2)
P1 <sup>i</sup> —Rh1—C1—O1	-8E1 (2)	C8—P1—C14—C19	137.1 (2)
Cl1 <sup>i</sup> —Rh1—C1—O1	-2E1 (3)	Rh1—P1—C14—C19	2.8 (2)
Cl1—Rh1—C1—O1	16E1 (3)	C2—P1—C14—C15	63.4 (2)
C1 <sup>i</sup> —Rh1—P1—C14	113.6 (2)	C8—P1—C14—C15	-41.7 (2)
C1—Rh1—P1—C14	-66.4 (2)	Rh1—P1—C14—C15	-175.97 (18)
P1 <sup>i</sup> —Rh1—P1—C14	-93.80 (10)	C15—C14—C19—C18	0.1 (4)
Cl1 <sup>i</sup> —Rh1—P1—C14	112.34 (10)	P1—C14—C19—C18	-178.8 (2)
Cl1—Rh1—P1—C14	-67.66 (10)	C19—C14—C15—C16	1.0 (4)
C1 <sup>i</sup> —Rh1—P1—C2	-129.2 (2)	P1—C14—C15—C16	179.8 (2)
C1—Rh1—P1—C2	50.8 (2)	C14—C19—C18—C17	-1.3 (4)
P1 <sup>i</sup> —Rh1—P1—C2	23.42 (13)	C13—C12—C11—C10	-0.5 (4)
Cl1 <sup>i</sup> —Rh1—P1—C2	-130.44 (10)	C2—C7—C6—C5	1.1 (4)
Cl1—Rh1—P1—C2	49.56 (10)	C19—C18—C17—C16	1.4 (4)
C1 <sup>i</sup> —Rh1—P1—C8	-8.5 (2)	C13—C8—C9—C10	0.0 (4)
C1—Rh1—P1—C8	171.5 (2)	P1—C8—C9—C10	-176.5 (2)
P1 <sup>i</sup> —Rh1—P1—C8	144.10 (13)	C5—C4—C3—C2	2.6 (4)
Cl1 <sup>i</sup> —Rh1—P1—C8	-9.76 (11)	C5—C4—C3—Br1	-178.16 (19)
Cl1—Rh1—P1—C8	170.24 (11)	C7—C2—C3—C4	-3.1 (4)
C6—C7—C2—C3	1.3 (4)	P1—C2—C3—C4	175.85 (19)
C6—C7—C2—P1	-177.7 (2)	C7—C2—C3—Br1	177.59 (18)
C14—P1—C2—C7	4.1 (2)	P1—C2—C3—Br1	-3.4 (3)
C8—P1—C2—C7	109.2 (2)	C3—C4—C5—C6	-0.1 (4)
Rh1—P1—C2—C7	-117.49 (19)	C7—C6—C5—C4	-1.7 (4)
C14—P1—C2—C3	-174.83 (19)	C11—C12—C13—C8	-0.9 (4)
C8—P1—C2—C3	-69.8 (2)	C9—C8—C13—C12	1.2 (4)
Rh1—P1—C2—C3	63.6 (2)	P1—C8—C13—C12	177.67 (19)
C14—P1—C8—C13	-84.6 (2)	C12—C11—C10—C9	1.6 (4)
C2—P1—C8—C13	167.38 (19)	C8—C9—C10—C11	-1.3 (4)
Rh1—P1—C8—C13	42.4 (2)	C14—C15—C16—C17	-0.9 (4)
C14—P1—C8—C9	91.8 (2)	C18—C17—C16—C15	-0.3 (4)
C2—P1—C8—C9	-16.2 (2)		

Symmetry code: (i)  $-x+1, -y, -z+2$ .