

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-(4-Chlorophenyl)-3-(morpholin-4-yl)-urea

Yu-Feng Li\* and Jin-He Jiang


 Microscale Science Institute, Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China  
 Correspondence e-mail: liyufeng8111@163.com

Received 20 March 2012; accepted 23 March 2012

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.180; data-to-parameter ratio = 18.6.

 In the title molecule,  $\text{C}_{11}\text{H}_{14}\text{ClN}_3\text{O}_2$ , the morpholine ring has a chair conformation. In the crystal, pairs of molecules are linked into inversion dimers by  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

 For the medicinal properties of related compounds, see: Yang *et al.* (1997). For a related structure, see: Li (2011).


## Experimental

### Crystal data

 $\text{C}_{11}\text{H}_{14}\text{ClN}_3\text{O}_2$   
 $M_r = 255.70$   
 Monoclinic,  $P2_1/c$   
 $a = 13.684$  (3) Å

 $b = 9.3612$  (19) Å  
 $c = 9.758$  (2) Å  
 $\beta = 94.03$  (3)°  
 $V = 1246.8$  (4) Å<sup>3</sup>
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.30$  mm<sup>-1</sup>
 $T = 293$  K  
 $0.22 \times 0.20 \times 0.19$  mm

### Data collection

 Bruker SMART CCD diffractometer  
 11894 measured reflections

 2857 independent reflections  
 1502 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.043$ 

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.180$   
 $S = 1.17$   
 2857 reflections

 154 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.20$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.29$  e Å<sup>-3</sup>
**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2A}\cdots\text{O2}^i$	0.86	2.03	2.865 (3)	162

 Symmetry code: (i)  $-x + 3, -y, -z$ .

 Data collection: *SMART* (Bruker 1997); cell refinement: *SAINTE* (Bruker 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors would like to thank the Natural Science Foundation of Shandong Province (No. ZR2010BL025).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5438).

## References

- Bruker (1997). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Li, Y.-F. (2011). *Acta Cryst.* **E67**, o1792.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yang, D., Soulier, J. L., Sicsic, S., Mathe-Allainmat, M., Bremont, B., Croci, T., Cardamone, R., Aureggi, G. & Langlois, M. (1997). *J. Med. Chem.* **40**, 608–621.

## supporting information

*Acta Cryst.* (2012). E68, o1219 [https://doi.org/10.1107/S1600536812012615]

## 1-(4-Chlorophenyl)-3-(morpholin-4-yl)urea

Yu-Feng Li and Jin-He Jiang

## S1. Comment

Some compounds which are related to the title compound have been shown to have medicinal properties (Yang *et al.*, 1997). The molecular structure of the title compound is shown in Fig. 1. The morpholine ring (C1–C4/O1/N1) has a chair conformation. The bond lengths and angles can be compared to those within a related structure (Li, 2011). In the crystal, pairs of molecules are linked into inversion dimers by N—H···O hydrogen bonds.

## S2. Experimental

A mixture of 4-aminomorpholine (0.08 mol), and (4-chlorophenyl)carbamic chloride (0.08 mol) was stirred in refluxing ethanol (18 ml) for 4 h to afford the title compound (0.064 mol, yield 80%). Colourless blocks of the title compound were obtained by recrystallization from ethanol at room temperature.

## S3. Refinement

H atoms were fixed geometrically and allowed to ride on their attached atoms, with C—H distances = 0.93–0.97 Å; N—H = 0.86 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ .




Figure 1

The structure of the title compound showing 30% probability displacement ellipsoids.

## 1-(4-Chlorophenyl)-3-(morpholin-4-yl)urea

*Crystal data*

$\text{C}_{11}\text{H}_{14}\text{ClN}_3\text{O}_2$

$M_r = 255.70$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 13.684\ (3)\ \text{\AA}$

$b = 9.3612\ (19)\ \text{\AA}$

$c = 9.758\ (2)\ \text{\AA}$

$\beta = 94.03\ (3)^\circ$

$V = 1246.8\ (4)\ \text{\AA}^3$

$Z = 4$

$F(000) = 536$

$D_x = 1.362\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1502 reflections

$\theta = 3.0\text{--}27.5^\circ$

$\mu = 0.30\ \text{mm}^{-1}$

$T = 293$  K  
Block, colorless

$0.22 \times 0.20 \times 0.19$  mm

#### Data collection

Bruker SMART CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
11894 measured reflections  
2857 independent reflections

1502 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.043$   
 $\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 3.0^\circ$   
 $h = -17 \rightarrow 17$   
 $k = -12 \rightarrow 12$   
 $l = -12 \rightarrow 12$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.180$   
 $S = 1.17$   
2857 reflections  
154 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.085P)^2 + 0.0207P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.20 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.29 \text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.92775 (5)	0.08842 (12)	0.30739 (8)	0.0988 (4)
O2	1.39233 (12)	0.0438 (2)	0.09026 (16)	0.0612 (5)
C6	1.18312 (16)	0.0784 (3)	0.0412 (2)	0.0493 (6)
N3	1.25450 (14)	0.0646 (2)	-0.0549 (2)	0.0561 (5)
H3A	1.2345	0.0708	-0.1403	0.067*
N1	1.35490 (13)	0.0024 (2)	-0.26930 (19)	0.0535 (5)
O1	1.35069 (15)	-0.0687 (2)	-0.55127 (17)	0.0697 (5)
N2	1.40369 (14)	0.0221 (3)	-0.1380 (2)	0.0656 (6)
H2A	1.4667	0.0212	-0.1292	0.079*
C5	1.35207 (17)	0.0425 (3)	-0.0260 (2)	0.0504 (6)
C8	1.1131 (2)	0.0212 (3)	0.2519 (3)	0.0674 (7)
H8A	1.1185	-0.0217	0.3381	0.081*
C11	1.09794 (16)	0.1525 (3)	-0.0004 (3)	0.0592 (6)
H11A	1.0931	0.1986	-0.0850	0.071*
C9	1.02870 (18)	0.0905 (3)	0.2072 (3)	0.0644 (7)

C10	1.02095 (18)	0.1582 (3)	0.0826 (3)	0.0672 (7)
H10A	0.9641	0.2076	0.0543	0.081*
C7	1.19009 (18)	0.0152 (3)	0.1684 (2)	0.0608 (7)
H7A	1.2474	-0.0321	0.1986	0.073*
C3	1.3409 (2)	0.0735 (3)	-0.5071 (3)	0.0734 (8)
H3B	1.2720	0.0954	-0.5020	0.088*
H3C	1.3669	0.1373	-0.5739	0.088*
C4	1.3938 (2)	0.0989 (3)	-0.3690 (3)	0.0666 (7)
H4A	1.4634	0.0822	-0.3741	0.080*
H4B	1.3847	0.1972	-0.3409	0.080*
C2	1.3146 (2)	-0.1633 (3)	-0.4542 (2)	0.0763 (8)
H2B	1.3226	-0.2609	-0.4849	0.092*
H2C	1.2451	-0.1462	-0.4478	0.092*
C1	1.3671 (2)	-0.1449 (3)	-0.3145 (3)	0.0709 (8)
H1A	1.3404	-0.2101	-0.2496	0.085*
H1B	1.4362	-0.1662	-0.3191	0.085*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0664 (5)	0.1530 (10)	0.0797 (6)	-0.0135 (5)	0.0244 (4)	-0.0336 (5)
O2	0.0541 (9)	0.0801 (13)	0.0483 (10)	0.0043 (8)	-0.0044 (8)	-0.0082 (8)
C6	0.0484 (12)	0.0545 (15)	0.0446 (12)	0.0017 (10)	0.0000 (10)	-0.0067 (10)
N3	0.0462 (10)	0.0747 (15)	0.0468 (11)	0.0059 (9)	-0.0007 (8)	-0.0015 (9)
N1	0.0532 (11)	0.0632 (14)	0.0438 (10)	0.0018 (9)	0.0020 (9)	-0.0021 (9)
O1	0.0918 (13)	0.0737 (13)	0.0449 (9)	0.0106 (10)	0.0143 (9)	0.0027 (9)
N2	0.0474 (11)	0.0996 (19)	0.0493 (11)	0.0054 (11)	-0.0004 (9)	-0.0114 (11)
C5	0.0500 (12)	0.0519 (15)	0.0488 (13)	0.0021 (10)	0.0005 (10)	-0.0025 (10)
C8	0.0644 (15)	0.090 (2)	0.0474 (13)	-0.0009 (15)	0.0025 (12)	-0.0051 (13)
C11	0.0540 (13)	0.0626 (17)	0.0605 (14)	0.0053 (12)	0.0002 (11)	0.0021 (13)
C9	0.0565 (14)	0.080 (2)	0.0576 (15)	-0.0081 (13)	0.0085 (12)	-0.0203 (14)
C10	0.0518 (13)	0.073 (2)	0.0767 (18)	0.0061 (13)	0.0022 (13)	-0.0102 (15)
C7	0.0557 (13)	0.0742 (19)	0.0519 (14)	0.0092 (13)	0.0002 (11)	-0.0006 (13)
C3	0.093 (2)	0.069 (2)	0.0590 (16)	0.0070 (16)	0.0062 (14)	0.0129 (14)
C4	0.0748 (17)	0.0613 (18)	0.0641 (16)	-0.0030 (13)	0.0083 (13)	0.0052 (13)
C2	0.111 (2)	0.0663 (19)	0.0530 (15)	-0.0077 (17)	0.0139 (15)	-0.0052 (13)
C1	0.103 (2)	0.0593 (17)	0.0501 (13)	0.0035 (16)	0.0059 (14)	0.0055 (12)

*Geometric parameters (Å, °)*

C11—C9	1.748 (3)	C8—H8A	0.9300
O2—C5	1.226 (3)	C11—C10	1.374 (3)
C6—C7	1.373 (3)	C11—H11A	0.9300
C6—C11	1.392 (3)	C9—C10	1.369 (4)
C6—N3	1.407 (3)	C10—H10A	0.9300
N3—C5	1.361 (3)	C7—H7A	0.9300
N3—H3A	0.8600	C3—C4	1.503 (4)
N1—N2	1.414 (3)	C3—H3B	0.9700

N1—C4	1.456 (3)	C3—H3C	0.9700
N1—C1	1.461 (3)	C4—H4A	0.9700
O1—C3	1.409 (4)	C4—H4B	0.9700
O1—C2	1.411 (3)	C2—C1	1.505 (4)
N2—C5	1.356 (3)	C2—H2B	0.9700
N2—H2A	0.8600	C2—H2C	0.9700
C8—C9	1.369 (4)	C1—H1A	0.9700
C8—C7	1.377 (4)	C1—H1B	0.9700
C7—C6—C11	118.8 (2)	C6—C7—C8	120.7 (2)
C7—C6—N3	123.8 (2)	C6—C7—H7A	119.7
C11—C6—N3	117.3 (2)	C8—C7—H7A	119.7
C5—N3—C6	126.35 (19)	O1—C3—C4	112.0 (2)
C5—N3—H3A	116.8	O1—C3—H3B	109.2
C6—N3—H3A	116.8	C4—C3—H3B	109.2
N2—N1—C4	110.6 (2)	O1—C3—H3C	109.2
N2—N1—C1	109.9 (2)	C4—C3—H3C	109.2
C4—N1—C1	109.2 (2)	H3B—C3—H3C	107.9
C3—O1—C2	110.0 (2)	N1—C4—C3	109.0 (2)
C5—N2—N1	120.60 (19)	N1—C4—H4A	109.9
C5—N2—H2A	119.7	C3—C4—H4A	109.9
N1—N2—H2A	119.7	N1—C4—H4B	109.9
O2—C5—N2	121.4 (2)	C3—C4—H4B	109.9
O2—C5—N3	124.1 (2)	H4A—C4—H4B	108.3
N2—C5—N3	114.4 (2)	O1—C2—C1	111.6 (2)
C9—C8—C7	119.6 (3)	O1—C2—H2B	109.3
C9—C8—H8A	120.2	C1—C2—H2B	109.3
C7—C8—H8A	120.2	O1—C2—H2C	109.3
C10—C11—C6	120.6 (2)	C1—C2—H2C	109.3
C10—C11—H11A	119.7	H2B—C2—H2C	108.0
C6—C11—H11A	119.7	N1—C1—C2	108.9 (2)
C8—C9—C10	120.9 (2)	N1—C1—H1A	109.9
C8—C9—C11	119.9 (2)	C2—C1—H1A	109.9
C10—C9—C11	119.1 (2)	N1—C1—H1B	109.9
C9—C10—C11	119.4 (2)	C2—C1—H1B	109.9
C9—C10—H10A	120.3	H1A—C1—H1B	108.3
C11—C10—H10A	120.3		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2A $\cdots$ O2 <sup>i</sup>	0.86	2.03	2.865 (3)	162

Symmetry code: (i)  $-x+3, -y, -z$ .