

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Absolute configuration of (1*S*,2*S*)-3-methyl-2-phenyl-2,3-dihydrothiazolo-[2,3-*b*]quinazolin-5-one

 Mostafa. M. Ghorab,<sup>a</sup> Mansour. S. Al-Said,<sup>a</sup> Maged. S. Abdel-Kader,<sup>b</sup> Madhukar Hemamalani<sup>c</sup> and Hoong-Kun Fun<sup>c\*‡</sup>

<sup>a</sup>Medicinal, Aromatic and Poisonous Plants Research Center, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, <sup>b</sup>Department of Pharmacognosy, College of Pharmacy, Salman Bin Abdulaziz University, PO Box 173, Al-Kharji 1194, Saudi Arabia, and <sup>c</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia  
Correspondence e-mail: hkfun@usm.my


Received 10 February 2012; accepted 24 February 2012

 Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.043;  $wR$  factor = 0.109; data-to-parameter ratio = 13.7.

The absolute structure of the molecule in the crystal of the title compound,  $\text{C}_{17}\text{H}_{14}\text{N}_2\text{OS}$ , was determined by the refinement of the Flack parameter to 0.0 (2) based on 1011 Friedel pairs. The quinazoline ring is essentially planar, with a maximum deviation of 0.037 (2) Å. The thiazole ring is distorted from planarity [maximum deviation = 0.168 (2) Å] and adopts a slightly twisted envelope conformation, with the C atom as the flap atom. The central thiazole ring makes dihedral angles of 7.01 (8) and 76.80 (10)° with the quinazoline and phenyl rings, respectively. The corresponding angle between the quinazoline and phenyl rings is 3.74 (9)°. In the crystal, there are no classical hydrogen bonds but stabilization is provided by weak  $\text{C}-\text{H}\cdots\pi$  interactions, involving the centroids of the phenyl rings.

## Related literature

For details and applications of quinazoline derivatives, see: Ghorab *et al.* (2010*a,b,c*). For related crystal structures, see: Al-Salahi *et al.* (2012); Priya *et al.* (2011); Liu *et al.* (2010). For ring conformations, see: Cremer & Pople (1975). For bond-length data, see: Allen *et al.* (1987).



‡ Alternative address: College of Pharmacy (Visiting Professor), King Saud University, PO Box 2457, Riyadh 11451, Saudi Arabia. Thomson Reuters ResearcherID: A-3561-2009.

## Experimental

## Crystal data

$\text{C}_{17}\text{H}_{14}\text{N}_2\text{OS}$   
 $M_r = 294.36$   
 Orthorhombic,  $P2_12_12_1$   
 $a = 8.4865$  (1) Å  
 $b = 10.0846$  (2) Å  
 $c = 16.8290$  (3) Å  
 $V = 1440.28$  (4) Å<sup>3</sup>  
 $Z = 4$   
 Cu  $K\alpha$  radiation  
 $\mu = 1.99$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.96 \times 0.64 \times 0.51$  mm

## Data collection

Bruker SMART APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2009)  
 $T_{\min} = 0.251$ ,  $T_{\max} = 0.431$   
 8654 measured reflections  
 2637 independent reflections  
 2521 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$   
 $wR(F^2) = 0.109$   
 $S = 1.08$   
 2637 reflections  
 192 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.22$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.23$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983), with 1011 Friedel pairs  
 Flack parameter: 0.00 (2)

Table 1

Hydrogen-bond geometry (Å, °).

$\text{Cg3}$  and  $\text{Cg4}$  are the centroids of the  $\text{C4}-\text{C9}$  and  $\text{C11}-\text{C16}$  phenyl rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C2}-\text{H2B}\cdots\text{Cg3}^i$	0.98	2.91	3.824 (2)	155
$\text{C6}-\text{H6A}\cdots\text{Cg4}^{ii}$	0.93	2.81	3.637 (3)	146

Symmetry codes: (i)  $-x - 1, y + \frac{3}{2}, -z + \frac{3}{2}$ ; (ii)  $-x + \frac{3}{2}, -y + 1, z + \frac{1}{2}$ .

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

MMG, MSA and MSA are grateful for sponsorship of the Research Center, College of Pharmacy, and the Deanship of Scientific Research, King Saud University, Riyadh, Saudi Arabia. MH and HFK thank the Malaysian Government and Universiti Sains Malaysia for Research University Grant No. 1001/PFIZIK/811160. MH also thanks Universiti Sains Malaysia for a postdoctoral research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2145).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.  
 Al-Salahi, R., Geffken, D. & Bari, A. (2012). *Acta Cryst. E* **68**, o101.  
 Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.  
 Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.  
 Ghorab, M. M., Ismail, Z. H. & Abdalla, M. (2010*a*). *Arzneim. Forsch. Drug Res.* **60**, 87–95.

- Ghorab, M. M., Ragab, F. A., Heiba, H. I., Arafa, R. K. & El-Hossary, E. M. (2010c). *Eur. J. Med. Chem.* **45**, 3677–3684.
- Ghorab, M. M., Ragab, F. A., Heiba, H. I., Youssef, H. A. & El-Gazzar, M. G. (2010b). *Bioorg. Med. Chem. Lett.*, **20**, 6316–6320.
- Liu, B., Chen, X.-B., Yang, X.-H., Pan, D.-F. & Ma, J.-K. (2010). *Acta Cryst. E66*, o2225–o2226.
- Priya, M. G. R., Srinivasan, T., Girija, K., Chandran, N. R. & Velmurugan, D. (2011). *Acta Cryst. E67*, o2310.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

## supporting information

*Acta Cryst.* (2012). E68, o927–o928 [https://doi.org/10.1107/S160053681200832X]

## Absolute configuration of (1*S*,2*S*)-3-methyl-2-phenyl-2,3-dihydrothiazolo[2,3-*b*]quinazolin-5-one

Mostafa. M. Ghorab, Mansour. S. Al-Said, Maged. S. Abdel-Kader, Madhukar Hemamalini and Hoong-Kun Fun

### S1. Comment

Quinazoline derivatives are well known as biologically active compounds. Consequently, quinazolines have been intensively studied for their interesting pharmacological properties such as anticancer activity (Ghorab *et al.*, 2010*a,b,c*). The crystal structures of 2-Ethoxy-5-methylbis[1,2,4]triazolo[1,5-*a*:4',3'-*c*] quinazoline (Al-Salahi *et al.*, 2012), 3-(4-Chlorophenyl)quinazolin-4(3*H*)-one (Priya *et al.*, 2011) and 2-Anilino-3-(2-hydroxyphenyl)quinazolin-4(3*H*)-one methanol monosolvate (Liu *et al.*, 2010) have been reported in the literature. Herein, we report the crystal structure of title compound (I).

The asymmetric unit of the title compound is shown in Fig. 1. The quinazoline (N1,N2/C3–C10) ring is essentially planar, with a maximum deviation of 0.037 (2) Å for atom C8. The thiazole (S1/N2/C7–C9) rings adopt an envelope conformation with the C2 (0.168 (2) Å) atom as the flap atom and with puckering parameter,  $Q = 0.2724$  (19) Å and  $\theta = 226.7$  (4)° (Cremer & Pople, 1975). The central thiazole (S1/N1/C1–C2,C10) ring makes dihedral angles of 7.01 (8)° and 76.80 (10)° with the quinazoline (N1,N2/C3–C10) and phenyl (C11–C16) rings, respectively. The corresponding angle between the quinazoline (N1,N2/C3–C10) and phenyl (C11–C16) rings is 73.74 (9)°. The bond lengths (Allen *et al.*, 1987) and angles are within normal ranges.

The absolute configuration of the molecule were determined by the refinement of the Flack parameter to 0.0 (2). There are two chiral centres in the molecule. From the structure presented, these centers exhibit the following chiralities: C1 = S and C2 = S.


In the crystal structure (Fig. 2), there are no classical hydrogen bonds but stabilization is provided by weak C—H $\cdots$  $\pi$  interactions (Table 1) involving the centroids of the (C4–C9) and (C11–C16) phenyl rings.

### S2. Experimental


A mixture of 2-isothiocyanatobenzoate (1.93 g, 0.01 mole) and 2-amino-1-phenylpropan-1-ol (1.51 g, 0.01 mole) in dry dimethylformamide (30 ml) containing a catalytic amount of triethylamine was refluxed for 6 h. The solid obtained was recrystallized from ethanol to give the title thiazoloquinazline derivative compound. Single crystals suitable for X-ray structural analysis were obtained by slow evaporation from ethanol at room temperature.

### S3. Refinement

All H atoms were positioned geometrically [C—H = 0.93–0.98 Å] and were refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5U_{\text{eq}}(\text{C})$ . A rotating group model was applied to the methyl groups. 1011 Friedel pairs were used to determine the absolute configuration.

**Figure 1**

The asymmetric unit of the title compound, showing 30% probability displacement ellipsoids.

**Figure 2**

The crystal packing of the title compound (I).

(1*S*,2*S*)-3-methyl-2-phenyl- 2,3-dihydrothiazolo[2,3-*b*]quinazolin-5-one

## Crystal data

C<sub>17</sub>H<sub>14</sub>N<sub>2</sub>OS $M_r = 294.36$ Orthorhombic,  $P2_12_12_1$ 

Hall symbol: P 2ac 2ab

 $a = 8.4865$  (1) Å $b = 10.0846$  (2) Å $c = 16.8290$  (3) Å $V = 1440.28$  (4) Å<sup>3</sup> $Z = 4$  $F(000) = 616$  $D_x = 1.358$  Mg m<sup>-3</sup>Cu  $K\alpha$  radiation,  $\lambda = 1.54178$  Å

Cell parameters from 4079 reflections

 $\theta = 4.4$ – $71.7^\circ$  $\mu = 1.99$  mm<sup>-1</sup> $T = 296$  K

Block, colourless

 $0.96 \times 0.64 \times 0.51$  mm

## Data collection

Bruker SMART APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\varphi$  and  $\omega$  scansAbsorption correction: multi-scan  
(*SADABS*; Bruker, 2009) $T_{\min} = 0.251$ ,  $T_{\max} = 0.431$ 

8654 measured reflections

2637 independent reflections

2521 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.028$  $\theta_{\max} = 71.9^\circ$ ,  $\theta_{\min} = 5.1^\circ$  $h = -10 \rightarrow 10$  $k = -8 \rightarrow 12$  $l = -20 \rightarrow 20$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.043$  $wR(F^2) = 0.109$  $S = 1.08$ 

2637 reflections

192 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0729P)^2 + 0.0628P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.22$  e Å<sup>-3</sup> $\Delta\rho_{\min} = -0.23$  e Å<sup>-3</sup>Extinction correction: *SHELXTL* (Sheldrick,  
2008),  $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$ 

Extinction coefficient: 0.107 (4)

Absolute structure: Flack (1983), with 1011

Friedel pairs

Absolute structure parameter: 0.00 (2)

## Special details

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.93740 (6)	0.77405 (8)	0.26159 (3)	0.0835 (3)

N1	0.84292 (17)	0.75544 (18)	0.40678 (9)	0.0558 (4)
N2	0.99073 (19)	0.57264 (18)	0.35894 (9)	0.0585 (4)
O1	0.7394 (3)	0.7625 (3)	0.53134 (11)	0.0983 (7)
C3	0.8142 (2)	0.7024 (3)	0.48182 (12)	0.0654 (5)
C10	0.9264 (2)	0.6865 (2)	0.35080 (10)	0.0540 (4)
C9	0.9715 (2)	0.51345 (19)	0.43270 (11)	0.0561 (4)
C12	0.5242 (2)	0.8021 (2)	0.28662 (15)	0.0693 (6)
H12A	0.5479	0.7496	0.3304	0.083*
C11	0.6379 (2)	0.88563 (18)	0.25601 (12)	0.0556 (4)
C4	0.8811 (2)	0.5710 (2)	0.49297 (11)	0.0598 (5)
C13	0.3752 (3)	0.7954 (3)	0.25291 (17)	0.0743 (6)
H13A	0.2993	0.7396	0.2746	0.089*
C5	0.8607 (3)	0.5028 (3)	0.56509 (14)	0.0774 (6)
H5A	0.8007	0.5405	0.6055	0.093*
C14	0.3400 (3)	0.8704 (2)	0.18807 (17)	0.0730 (6)
H14A	0.2396	0.8670	0.1660	0.088*
C16	0.6023 (3)	0.9594 (2)	0.18898 (14)	0.0633 (5)
H16A	0.6783	1.0144	0.1666	0.076*
C6	0.9290 (4)	0.3807 (3)	0.57612 (17)	0.0875 (8)
H6A	0.9131	0.3350	0.6235	0.105*
C1	0.7973 (2)	0.9010 (2)	0.29489 (15)	0.0659 (5)
H1A	0.8396	0.9878	0.2798	0.079*
C15	0.4533 (3)	0.9515 (2)	0.15507 (14)	0.0723 (6)
H15A	0.4299	1.0010	0.1100	0.087*
C8	1.0433 (3)	0.3896 (2)	0.44566 (15)	0.0744 (6)
H8A	1.1052	0.3517	0.4061	0.089*
C2	0.7980 (2)	0.8926 (2)	0.38635 (14)	0.0659 (5)
H2B	0.6911	0.9094	0.4060	0.079*
C17	0.9107 (4)	0.9910 (3)	0.4252 (3)	0.1039 (11)
H17A	0.9070	0.9806	0.4818	0.156*
H17C	0.8802	1.0797	0.4113	0.156*
H17D	1.0160	0.9746	0.4067	0.156*
C7	1.0225 (4)	0.3247 (3)	0.51644 (19)	0.0874 (8)
H7A	1.0707	0.2431	0.5248	0.105*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0541 (3)	0.1285 (5)	0.0680 (3)	0.0297 (3)	0.0168 (2)	0.0362 (3)
N1	0.0436 (7)	0.0702 (9)	0.0535 (8)	0.0037 (6)	-0.0037 (6)	-0.0028 (7)
N2	0.0529 (8)	0.0726 (10)	0.0500 (7)	0.0055 (7)	-0.0055 (6)	-0.0051 (7)
O1	0.1001 (14)	0.1344 (18)	0.0602 (9)	0.0463 (14)	0.0130 (8)	-0.0049 (10)
C3	0.0512 (9)	0.0952 (14)	0.0497 (9)	0.0092 (10)	-0.0022 (8)	-0.0043 (9)
C10	0.0393 (7)	0.0739 (11)	0.0490 (8)	0.0004 (7)	-0.0014 (7)	0.0020 (7)
C9	0.0526 (9)	0.0635 (10)	0.0522 (9)	-0.0071 (7)	-0.0092 (7)	-0.0034 (7)
C12	0.0519 (10)	0.0741 (12)	0.0820 (13)	-0.0033 (8)	-0.0106 (9)	0.0223 (10)
C11	0.0469 (8)	0.0523 (8)	0.0675 (10)	0.0055 (6)	-0.0032 (8)	0.0022 (8)
C4	0.0472 (8)	0.0815 (12)	0.0507 (9)	-0.0083 (8)	-0.0059 (7)	0.0028 (8)

C13	0.0491 (10)	0.0816 (13)	0.0923 (16)	-0.0055 (9)	-0.0084 (10)	0.0104 (12)
C5	0.0615 (12)	0.1126 (18)	0.0579 (11)	-0.0103 (12)	-0.0015 (9)	0.0107 (12)
C14	0.0527 (10)	0.0768 (13)	0.0896 (15)	0.0096 (9)	-0.0178 (11)	-0.0050 (11)
C16	0.0634 (11)	0.0560 (9)	0.0705 (11)	0.0068 (8)	-0.0005 (9)	0.0064 (8)
C6	0.0857 (16)	0.0934 (17)	0.0833 (15)	-0.0174 (14)	-0.0132 (14)	0.0307 (13)
C1	0.0460 (9)	0.0666 (10)	0.0850 (14)	-0.0021 (8)	-0.0040 (9)	0.0180 (10)
C15	0.0745 (13)	0.0697 (11)	0.0727 (12)	0.0158 (10)	-0.0151 (10)	0.0056 (9)
C8	0.0880 (15)	0.0662 (11)	0.0691 (11)	0.0015 (11)	-0.0152 (12)	-0.0065 (9)
C2	0.0505 (9)	0.0676 (11)	0.0795 (13)	0.0061 (8)	-0.0114 (9)	-0.0047 (10)
C17	0.0876 (19)	0.0806 (16)	0.143 (3)	-0.0012 (14)	-0.038 (2)	-0.0195 (18)
C7	0.107 (2)	0.0709 (13)	0.0845 (15)	-0.0052 (13)	-0.0216 (15)	0.0107 (11)

*Geometric parameters (Å, °)*

S1—C10	1.7439 (18)	C5—C6	1.374 (5)
S1—C1	1.835 (2)	C5—H5A	0.9300
N1—C10	1.368 (3)	C14—C15	1.379 (4)
N1—C3	1.393 (3)	C14—H14A	0.9300
N1—C2	1.475 (3)	C16—C15	1.390 (3)
N2—C10	1.279 (3)	C16—H16A	0.9300
N2—C9	1.387 (3)	C6—C7	1.399 (5)
O1—C3	1.211 (3)	C6—H6A	0.9300
C3—C4	1.454 (3)	C1—C2	1.542 (3)
C9—C4	1.398 (3)	C1—H1A	0.9800
C9—C8	1.406 (3)	C15—H15A	0.9300
C12—C11	1.380 (3)	C8—C7	1.371 (4)
C12—C13	1.388 (3)	C8—H8A	0.9300
C12—H12A	0.9300	C2—C17	1.526 (3)
C11—C16	1.385 (3)	C2—H2B	0.9800
C11—C1	1.511 (3)	C17—H17A	0.9600
C4—C5	1.406 (3)	C17—H17C	0.9600
C13—C14	1.361 (4)	C17—H17D	0.9600
C13—H13A	0.9300	C7—H7A	0.9300
C10—S1—C1	93.18 (10)	C11—C16—H16A	119.9
C10—N1—C3	121.33 (18)	C15—C16—H16A	119.9
C10—N1—C2	116.71 (17)	C5—C6—C7	120.3 (2)
C3—N1—C2	121.72 (18)	C5—C6—H6A	119.9
C10—N2—C9	115.61 (16)	C7—C6—H6A	119.9
O1—C3—N1	121.6 (2)	C11—C1—C2	115.45 (17)
O1—C3—C4	124.9 (2)	C11—C1—S1	112.12 (16)
N1—C3—C4	113.49 (18)	C2—C1—S1	105.33 (14)
N2—C10—N1	127.10 (17)	C11—C1—H1A	107.9
N2—C10—S1	121.59 (14)	C2—C1—H1A	107.9
N1—C10—S1	111.31 (14)	S1—C1—H1A	107.9
N2—C9—C4	122.34 (19)	C14—C15—C16	120.2 (2)
N2—C9—C8	118.0 (2)	C14—C15—H15A	119.9
C4—C9—C8	119.6 (2)	C16—C15—H15A	119.9

C11—C12—C13	120.9 (2)	C7—C8—C9	120.2 (3)
C11—C12—H12A	119.5	C7—C8—H8A	119.9
C13—C12—H12A	119.5	C9—C8—H8A	119.9
C12—C11—C16	118.68 (18)	N1—C2—C17	110.36 (19)
C12—C11—C1	121.78 (18)	N1—C2—C1	106.58 (17)
C16—C11—C1	119.52 (18)	C17—C2—C1	113.2 (2)
C9—C4—C5	119.4 (2)	N1—C2—H2B	108.9
C9—C4—C3	119.94 (18)	C17—C2—H2B	108.9
C5—C4—C3	120.6 (2)	C1—C2—H2B	108.9
C14—C13—C12	120.0 (2)	C2—C17—H17A	109.5
C14—C13—H13A	120.0	C2—C17—H17C	109.5
C12—C13—H13A	120.0	H17A—C17—H17C	109.5
C6—C5—C4	120.2 (3)	C2—C17—H17D	109.5
C6—C5—H5A	119.9	H17A—C17—H17D	109.5
C4—C5—H5A	119.9	H17C—C17—H17D	109.5
C13—C14—C15	120.0 (2)	C8—C7—C6	120.3 (3)
C13—C14—H14A	120.0	C8—C7—H7A	119.8
C15—C14—H14A	120.0	C6—C7—H7A	119.8
C11—C16—C15	120.1 (2)		
C10—N1—C3—O1	-179.4 (2)	C3—C4—C5—C6	-177.7 (2)
C2—N1—C3—O1	6.4 (3)	C12—C13—C14—C15	1.1 (4)
C10—N1—C3—C4	0.1 (3)	C12—C11—C16—C15	1.8 (3)
C2—N1—C3—C4	-174.12 (17)	C1—C11—C16—C15	-176.6 (2)
C9—N2—C10—N1	0.6 (3)	C4—C5—C6—C7	1.6 (4)
C9—N2—C10—S1	-179.34 (13)	C12—C11—C1—C2	-34.8 (3)
C3—N1—C10—N2	-2.2 (3)	C16—C11—C1—C2	143.6 (2)
C2—N1—C10—N2	172.28 (18)	C12—C11—C1—S1	85.8 (2)
C3—N1—C10—S1	177.70 (15)	C16—C11—C1—S1	-95.81 (19)
C2—N1—C10—S1	-7.8 (2)	C10—S1—C1—C11	-105.81 (15)
C1—S1—C10—N2	171.57 (16)	C10—S1—C1—C2	20.52 (15)
C1—S1—C10—N1	-8.37 (15)	C13—C14—C15—C16	-1.6 (4)
C10—N2—C9—C4	3.2 (3)	C11—C16—C15—C14	0.1 (3)
C10—N2—C9—C8	-178.35 (18)	N2—C9—C8—C7	-177.4 (2)
C13—C12—C11—C16	-2.4 (4)	C4—C9—C8—C7	1.2 (3)
C13—C12—C11—C1	176.0 (2)	C10—N1—C2—C17	-99.9 (3)
N2—C9—C4—C5	177.15 (19)	C3—N1—C2—C17	74.6 (3)
C8—C9—C4—C5	-1.3 (3)	C10—N1—C2—C1	23.4 (2)
N2—C9—C4—C3	-5.2 (3)	C3—N1—C2—C1	-162.12 (17)
C8—C9—C4—C3	176.33 (19)	C11—C1—C2—N1	97.6 (2)
O1—C3—C4—C9	-177.2 (2)	S1—C1—C2—N1	-26.69 (19)
N1—C3—C4—C9	3.3 (3)	C11—C1—C2—C17	-140.9 (2)
O1—C3—C4—C5	0.4 (4)	S1—C1—C2—C17	94.8 (2)
N1—C3—C4—C5	-179.06 (19)	C9—C8—C7—C6	0.3 (4)
C11—C12—C13—C14	0.9 (4)	C5—C6—C7—C8	-1.7 (4)
C9—C4—C5—C6	0.0 (3)		



*Hydrogen-bond geometry (Å, °)*

Cg3 and Cg4 are the centroids of the C4–C9 and C11–C16 phenyl rings, respectively.

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C2—H2 <i>B</i> $\cdots$ Cg3 <sup>i</sup>	0.98	2.91	3.824 (2)	155
C6—H6 <i>A</i> $\cdots$ Cg4 <sup>ii</sup>	0.93	2.81	3.637 (3)	146

Symmetry codes: (i)  $-x-1, y+3/2, -z+3/2$ ; (ii)  $-x+3/2, -y+1, z+1/2$ .