

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

cis-Dichloridobis(quinoline- κ N)-platinum(II) nitromethane monosolvate

Kwang Ha

 School of Applied Chemical Engineering, Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea
 Correspondence e-mail: hakwang@chonnam.ac.kr


Received 16 March 2012; accepted 22 March 2012

 Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.013$ Å; R factor = 0.030; wR factor = 0.084; data-to-parameter ratio = 15.1.

In the title compound, $[\text{PtCl}_2(\text{C}_9\text{H}_7\text{N})_2] \cdot \text{CH}_3\text{NO}_2$, the Pt^{II} cation is four-coordinated in an essentially square-planar environment by two N atoms from two quinoline ligands and two Cl^- anions. One of the nearly planar quinoline ligands [maximum deviations = 0.042 (6) and 0.018 (7) Å] is almost perpendicular to the PtCl_2N_2 unit [maximum deviation = 0.024 (3) Å], making a dihedral angle of 89.6 (1)°, whereas the other is slightly inclined to the central plane with a dihedral angle of 74.1 (1)°. The dihedral angle between the quinoline ligands is 88.3 (2)°. In the crystal, each solvent molecule is linked to the metal complex by weak intermolecular $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds.

Related literature

For the crystal structure of *cis*- $[\text{PtCl}_2(\text{quinoline})_2] \cdot 0.25\text{DMF}$ (DMF = *N,N*-dimethylformamide), see: Davies *et al.* (2001). For the crystal structure of the related Pd^{II} complex *trans*- $[\text{PdCl}_2(\text{quinoline})_2]$, see: Ha (2012).



Experimental

Crystal data

 $[\text{PtCl}_2(\text{C}_9\text{H}_7\text{N})_2] \cdot \text{CH}_3\text{NO}_2$
 $M_r = 585.35$
 Triclinic, $P\bar{1}$
 $a = 9.6204$ (5) Å
 $b = 10.3698$ (5) Å

 $c = 11.6946$ (6) Å
 $\alpha = 104.244$ (1)°
 $\beta = 101.913$ (1)°
 $\gamma = 113.834$ (1)°
 $V = 970.87$ (8) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 7.52$ mm⁻¹
 $T = 200$ K
 $0.25 \times 0.19 \times 0.13$ mm

Data collection

 Bruker SMART 1000 CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\text{min}} = 0.754$, $T_{\text{max}} = 1.000$

 6035 measured reflections
 3707 independent reflections
 3342 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.084$
 $S = 1.24$
 3707 reflections

 245 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 2.69$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.38$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Pt1—N1	2.045 (6)	Pt1—Cl1	2.2881 (18)
Pt1—N2	2.045 (6)	Pt1—Cl2	2.3019 (19)
N2—Pt1—N1	90.3 (2)	Cl1—Pt1—Cl2	91.99 (7)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C16}-\text{H16} \cdots \text{O2}^i$	0.95	2.59	3.323 (13)	134

 Symmetry code: (i) $-x, -y, -z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97.

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (grant No. 2011-0030747).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5489).

References

- Bruker (2000). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Davies, M. S., Diakos, C. I., Messerle, B. A. & Hambley, T. W. (2001). *Inorg. Chem.* **40**, 3048–3054.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Ha, K. (2012). *Acta Cryst.* **E68**, m143.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, m491 [https://doi.org/10.1107/S1600536812012469]

cis*-Dichloridobis(quinoline- κ N)platinum(II) nitromethane monosolvate*Kwang Ha****S1. Comment**

The asymmetric unit of the title compound, [PtCl₂(quinoline)₂].CH₃NO₂, contains a neutral Pt^{II} complex and a nitromethane solvent molecule (Fig. 1). In the complex, the Pt^{II} ion is four-coordinated in an essentially square-planar environment by two N atoms from two quinoline ligands and two Cl⁻ anions (Table 1). The Cl atoms are in *cis* conformation with respect to each other, like in the analogous Pt^{II} complex [PtCl₂(quinoline)₂].0.25DMF (DMF = *N,N*-dimethylformamide) (Davies *et al.*, 2001). By contrast, in the related Pd^{II} complex [PdCl₂(quinoline)₂], the Cl atoms are in *trans* conformation (Ha, 2012).

One of the nearly planar quinoline ligands [maximum deviations = 0.042 (6) Å and 0.018 (7) Å] is almost perpendicular to the PtCl₂N₂ unit [maximum deviation = 0.024 (3) Å], making dihedral angle of 89.6 (1)°, whereas the other is slightly inclined to the unit plane with a dihedral angle of 74.1 (1)°. The dihedral angle between the quinoline ligands is 88.3 (2)°. In the crystal, each solvent molecule is linked to the complex by intermolecular C—H···O hydrogen bonds (Fig. 2 and Table 2). Moreover, the complex molecules display numerous intermolecular π - π interactions between adjacent six-membered rings, the shortest ring centroid-centroid distance being 3.617 (5) Å.

S2. Experimental

To a solution of K₂PtCl₄ (0.2074 g, 0.500 mmol) in H₂O (20 ml) was added quinoline (0.1304 g, 1.010 mmol), and refluxed for 3 h. The formed precipitate was separated by filtration, washed with H₂O and EtOH, and dried at 50°C, to give a yellow powder (0.1761 g). Crystals suitable for X-ray analysis were obtained by slow evaporation from a CH₃NO₂ solution at room temperature.

S3. Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms: C—H = 0.95 Å (CH) or 0.98 Å (CH₃) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$. The highest peak (2.69 e Å⁻³) and the deepest hole (-1.38 e Å⁻³) in the difference Fourier map are located 0.86 Å and 0.75 Å, respectively, from the Pt1 atom.




Figure 1

A structure detail of the title compound, with atom numbering. Displacement ellipsoids are drawn at the 40% probability level for non-H atoms.




Figure 2

A view of the unit-cell contents of the title compound. Intermolecular C—H...O hydrogen-bond interactions are drawn with dashed lines.

cis-Dichloridobis(quinoline- κ N)platinum(II) nitromethane monosolvate

Crystal data

[PtCl₂(C₆H₇N)₂]·CH₃NO₂

$M_r = 585.35$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.6204 (5) \text{ \AA}$

$b = 10.3698 (5) \text{ \AA}$

$c = 11.6946 (6) \text{ \AA}$

$\alpha = 104.244 (1)^\circ$

$\beta = 101.913 (1)^\circ$

$\gamma = 113.834 (1)^\circ$

$V = 970.87 (8) \text{ \AA}^3$

$Z = 2$

$F(000) = 560$

$D_x = 2.002 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4362 reflections

$\theta = 2.4\text{--}26.0^\circ$

$\mu = 7.52 \text{ mm}^{-1}$

$T = 200 \text{ K}$

Block, yellow

$0.25 \times 0.19 \times 0.13 \text{ mm}$

Data collection

Bruker SMART 1000 CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2000)

$T_{\min} = 0.754$, $T_{\max} = 1.000$

6035 measured reflections
 3707 independent reflections
 3342 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

$\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 1.9^\circ$
 $h = -9 \rightarrow 11$
 $k = -12 \rightarrow 12$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.084$
 $S = 1.24$
 3707 reflections
 245 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.P)^2 + 7.8267P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 2.69 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -1.38 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pt1	0.24674 (4)	0.27007 (3)	0.45574 (3)	0.02575 (10)
Cl1	0.0985 (2)	0.3934 (2)	0.48336 (18)	0.0326 (4)
Cl2	0.2977 (3)	0.2747 (3)	0.65828 (18)	0.0395 (5)
N1	0.3854 (7)	0.1676 (7)	0.4262 (6)	0.0245 (13)
N2	0.2011 (7)	0.2682 (7)	0.2765 (6)	0.0252 (13)
C1	0.5394 (9)	0.2527 (9)	0.4458 (7)	0.0284 (16)
H1	0.5840	0.3595	0.4829	0.034*
C2	0.6416 (10)	0.1950 (9)	0.4153 (7)	0.0327 (18)
H2	0.7528	0.2614	0.4333	0.039*
C3	0.5792 (10)	0.0430 (9)	0.3596 (7)	0.0310 (17)
H3	0.6456	0.0013	0.3368	0.037*
C4	0.4121 (9)	-0.0533 (9)	0.3356 (7)	0.0279 (16)
C5	0.3400 (10)	-0.2136 (9)	0.2791 (7)	0.0345 (18)
H5	0.4021	-0.2590	0.2533	0.041*
C6	0.1834 (10)	-0.3010 (9)	0.2622 (7)	0.0358 (19)
H6	0.1349	-0.4078	0.2213	0.043*
C7	0.0899 (10)	-0.2368 (9)	0.3042 (7)	0.0331 (18)
H7	-0.0189	-0.3013	0.2944	0.040*
C8	0.1541 (9)	-0.0833 (8)	0.3586 (6)	0.0274 (16)
H8	0.0909	-0.0407	0.3869	0.033*
C9	0.3187 (8)	0.0128 (8)	0.3725 (6)	0.0218 (14)

C10	0.3191 (11)	0.3682 (9)	0.2522 (8)	0.0331 (18)
H10	0.4228	0.4291	0.3160	0.040*
C11	0.2975 (11)	0.3873 (9)	0.1381 (8)	0.041 (2)
H11	0.3854	0.4592	0.1243	0.049*
C12	0.1504 (12)	0.3031 (9)	0.0463 (8)	0.044 (2)
H12	0.1338	0.3173	-0.0316	0.053*
C13	0.0218 (10)	0.1942 (9)	0.0665 (7)	0.0333 (18)
C14	-0.1344 (12)	0.1003 (11)	-0.0233 (8)	0.044 (2)
H14	-0.1561	0.1111	-0.1024	0.052*
C15	-0.2533 (12)	-0.0033 (11)	-0.0024 (8)	0.049 (2)
H15	-0.3571	-0.0651	-0.0660	0.059*
C16	-0.2242 (10)	-0.0207 (10)	0.1153 (8)	0.040 (2)
H16	-0.3084	-0.0954	0.1300	0.048*
C17	-0.0761 (10)	0.0692 (9)	0.2076 (7)	0.0343 (18)
H17	-0.0582	0.0585	0.2869	0.041*
C18	0.0508 (9)	0.1783 (8)	0.1850 (7)	0.0275 (16)
O1	0.8350 (9)	0.4264 (8)	0.0474 (7)	0.067 (2)
O2	0.5859 (11)	0.3279 (13)	0.0137 (9)	0.112 (4)
N3	0.7193 (10)	0.3685 (8)	0.0788 (7)	0.0438 (18)
C19	0.7469 (14)	0.3484 (11)	0.1983 (9)	0.061 (3)
H19A	0.7237	0.4160	0.2566	0.091*
H19B	0.8600	0.3726	0.2334	0.091*
H19C	0.6756	0.2430	0.1857	0.091*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pt1	0.02316 (16)	0.02821 (16)	0.02450 (16)	0.01381 (13)	0.00675 (12)	0.00619 (12)
C11	0.0309 (10)	0.0313 (10)	0.0369 (10)	0.0193 (9)	0.0113 (8)	0.0075 (8)
C12	0.0361 (11)	0.0639 (14)	0.0288 (10)	0.0318 (11)	0.0133 (9)	0.0177 (10)
N1	0.020 (3)	0.033 (3)	0.029 (3)	0.014 (3)	0.016 (3)	0.015 (3)
N2	0.026 (3)	0.025 (3)	0.028 (3)	0.018 (3)	0.007 (3)	0.008 (3)
C1	0.017 (4)	0.032 (4)	0.033 (4)	0.010 (3)	0.005 (3)	0.014 (3)
C2	0.032 (4)	0.038 (5)	0.032 (4)	0.017 (4)	0.008 (4)	0.021 (4)
C3	0.039 (5)	0.042 (5)	0.030 (4)	0.027 (4)	0.017 (4)	0.022 (4)
C4	0.032 (4)	0.040 (4)	0.028 (4)	0.027 (4)	0.012 (3)	0.020 (3)
C5	0.037 (5)	0.038 (5)	0.035 (4)	0.025 (4)	0.011 (4)	0.013 (4)
C6	0.038 (5)	0.029 (4)	0.031 (4)	0.015 (4)	0.006 (4)	0.003 (3)
C7	0.028 (4)	0.026 (4)	0.039 (5)	0.012 (3)	0.006 (4)	0.010 (3)
C8	0.025 (4)	0.031 (4)	0.022 (4)	0.012 (3)	0.003 (3)	0.009 (3)
C9	0.020 (4)	0.026 (4)	0.018 (3)	0.012 (3)	0.002 (3)	0.007 (3)
C10	0.044 (5)	0.033 (4)	0.039 (5)	0.023 (4)	0.024 (4)	0.024 (4)
C11	0.046 (5)	0.028 (4)	0.047 (5)	0.016 (4)	0.017 (4)	0.013 (4)
C12	0.069 (7)	0.033 (5)	0.029 (4)	0.025 (5)	0.017 (5)	0.008 (4)
C13	0.040 (5)	0.032 (4)	0.025 (4)	0.022 (4)	0.001 (4)	0.005 (3)
C14	0.052 (6)	0.053 (6)	0.030 (5)	0.031 (5)	0.010 (4)	0.014 (4)
C15	0.039 (5)	0.061 (6)	0.031 (5)	0.026 (5)	-0.005 (4)	0.002 (4)
C16	0.026 (4)	0.044 (5)	0.034 (5)	0.011 (4)	0.002 (4)	0.003 (4)

C17	0.039 (5)	0.040 (5)	0.023 (4)	0.023 (4)	0.009 (4)	0.007 (3)
C18	0.031 (4)	0.027 (4)	0.026 (4)	0.019 (3)	0.005 (3)	0.007 (3)
O1	0.062 (5)	0.061 (5)	0.064 (5)	0.013 (4)	0.025 (4)	0.028 (4)
O2	0.051 (6)	0.168 (10)	0.072 (6)	0.023 (6)	0.001 (5)	0.043 (7)
N3	0.042 (5)	0.034 (4)	0.027 (4)	0.004 (4)	0.000 (3)	0.001 (3)
C19	0.071 (8)	0.049 (6)	0.041 (6)	0.013 (6)	0.010 (5)	0.017 (5)

Geometric parameters (Å, °)

Pt1—N1	2.045 (6)	C8—H8	0.9500
Pt1—N2	2.045 (6)	C10—C11	1.383 (11)
Pt1—C11	2.2881 (18)	C10—H10	0.9500
Pt1—C12	2.3019 (19)	C11—C12	1.353 (13)
N1—C1	1.315 (9)	C11—H11	0.9500
N1—C9	1.375 (9)	C12—C13	1.409 (12)
N2—C10	1.331 (10)	C12—H12	0.9500
N2—C18	1.373 (9)	C13—C14	1.404 (12)
C1—C2	1.403 (11)	C13—C18	1.420 (10)
C1—H1	0.9500	C14—C15	1.332 (13)
C2—C3	1.355 (11)	C14—H14	0.9500
C2—H2	0.9500	C15—C16	1.418 (12)
C3—C4	1.428 (11)	C15—H15	0.9500
C3—H3	0.9500	C16—C17	1.364 (11)
C4—C9	1.405 (10)	C16—H16	0.9500
C4—C5	1.420 (11)	C17—C18	1.416 (11)
C5—C6	1.348 (12)	C17—H17	0.9500
C5—H5	0.9500	O1—N3	1.213 (10)
C6—C7	1.413 (11)	O2—N3	1.185 (11)
C6—H6	0.9500	N3—C19	1.450 (11)
C7—C8	1.366 (10)	C19—H19A	0.9800
C7—H7	0.9500	C19—H19B	0.9800
C8—C9	1.438 (10)	C19—H19C	0.9800
N2—Pt1—N1	90.3 (2)	C4—C9—C8	119.4 (7)
N2—Pt1—C11	87.40 (17)	N2—C10—C11	122.9 (8)
N1—Pt1—C11	176.81 (17)	N2—C10—H10	118.5
N2—Pt1—C12	179.33 (17)	C11—C10—H10	118.5
N1—Pt1—C12	90.30 (17)	C12—C11—C10	119.6 (8)
C11—Pt1—C12	91.99 (7)	C12—C11—H11	120.2
C1—N1—C9	118.8 (6)	C10—C11—H11	120.2
C1—N1—Pt1	118.8 (5)	C11—C12—C13	119.8 (8)
C9—N1—Pt1	121.9 (5)	C11—C12—H12	120.1
C10—N2—C18	119.4 (7)	C13—C12—H12	120.1
C10—N2—Pt1	117.8 (5)	C14—C13—C12	124.0 (8)
C18—N2—Pt1	122.5 (5)	C14—C13—C18	117.6 (8)
N1—C1—C2	123.9 (7)	C12—C13—C18	118.3 (7)
N1—C1—H1	118.1	C15—C14—C13	122.9 (8)
C2—C1—H1	118.1	C15—C14—H14	118.6

C3—C2—C1	118.9 (8)	C13—C14—H14	118.6
C3—C2—H2	120.5	C14—C15—C16	119.6 (8)
C1—C2—H2	120.5	C14—C15—H15	120.2
C2—C3—C4	118.9 (7)	C16—C15—H15	120.2
C2—C3—H3	120.5	C17—C16—C15	120.4 (9)
C4—C3—H3	120.5	C17—C16—H16	119.8
C9—C4—C5	119.6 (7)	C15—C16—H16	119.8
C9—C4—C3	118.7 (7)	C16—C17—C18	120.0 (8)
C5—C4—C3	121.7 (7)	C16—C17—H17	120.0
C6—C5—C4	119.9 (7)	C18—C17—H17	120.0
C6—C5—H5	120.0	N2—C18—C17	120.6 (7)
C4—C5—H5	120.0	N2—C18—C13	119.9 (7)
C5—C6—C7	121.2 (7)	C17—C18—C13	119.4 (7)
C5—C6—H6	119.4	O2—N3—O1	121.4 (9)
C7—C6—H6	119.4	O2—N3—C19	120.1 (9)
C8—C7—C6	120.8 (8)	O1—N3—C19	118.5 (8)
C8—C7—H7	119.6	N3—C19—H19A	109.5
C6—C7—H7	119.6	N3—C19—H19B	109.5
C7—C8—C9	119.0 (7)	H19A—C19—H19B	109.5
C7—C8—H8	120.5	N3—C19—H19C	109.5
C9—C8—H8	120.5	H19A—C19—H19C	109.5
N1—C9—C4	120.6 (7)	H19B—C19—H19C	109.5
N1—C9—C8	120.0 (6)		
N2—Pt1—N1—C1	86.5 (6)	C5—C4—C9—C8	3.3 (10)
Cl2—Pt1—N1—C1	-93.2 (5)	C3—C4—C9—C8	-174.8 (6)
N2—Pt1—N1—C9	-85.6 (5)	C7—C8—C9—N1	179.6 (7)
Cl2—Pt1—N1—C9	94.7 (5)	C7—C8—C9—C4	-3.1 (10)
N1—Pt1—N2—C10	-75.1 (5)	C18—N2—C10—C11	0.5 (11)
Cl1—Pt1—N2—C10	102.7 (5)	Pt1—N2—C10—C11	-173.6 (6)
N1—Pt1—N2—C18	111.0 (5)	N2—C10—C11—C12	0.8 (12)
Cl1—Pt1—N2—C18	-71.3 (5)	C10—C11—C12—C13	-1.6 (13)
C9—N1—C1—C2	-0.1 (11)	C11—C12—C13—C14	-179.2 (8)
Pt1—N1—C1—C2	-172.4 (6)	C11—C12—C13—C18	1.1 (12)
N1—C1—C2—C3	1.6 (11)	C12—C13—C14—C15	179.1 (9)
C1—C2—C3—C4	-0.9 (11)	C18—C13—C14—C15	-1.2 (13)
C2—C3—C4—C9	-1.0 (10)	C13—C14—C15—C16	0.5 (14)
C2—C3—C4—C5	-179.1 (7)	C14—C15—C16—C17	0.9 (14)
C9—C4—C5—C6	-0.5 (11)	C15—C16—C17—C18	-1.5 (13)
C3—C4—C5—C6	177.6 (7)	C10—N2—C18—C17	178.9 (7)
C4—C5—C6—C7	-2.5 (12)	Pt1—N2—C18—C17	-7.2 (9)
C5—C6—C7—C8	2.7 (12)	C10—N2—C18—C13	-1.0 (10)
C6—C7—C8—C9	0.2 (11)	Pt1—N2—C18—C13	172.8 (5)
C1—N1—C9—C4	-1.9 (10)	C16—C17—C18—N2	-179.2 (7)
Pt1—N1—C9—C4	170.2 (5)	C16—C17—C18—C13	0.7 (11)
C1—N1—C9—C8	175.3 (6)	C14—C13—C18—N2	-179.5 (7)
Pt1—N1—C9—C8	-12.6 (9)	C12—C13—C18—N2	0.2 (11)
C5—C4—C9—N1	-179.4 (7)	C14—C13—C18—C17	0.6 (11)

C3—C4—C9—N1	2.5 (10)	C12—C13—C18—C17	-179.7 (7)
-------------	----------	-----------------	------------

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C16—H16···O2 ⁱ	0.95	2.59	3.323 (13)	134

Symmetry code: (i) $-x, -y, -z$.