

N'-(*E*)-4-Benzyl-2-hydroxybenzylidene]-4-nitrobenzohydrazide monohydrate

Bibitha Joseph,^a M. Sithambaresan^{b*} and
M. R. Prathapachandra Kurup^a

^aDepartment of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, India, and ^bDepartment of Chemistry, Faculty of Science, Eastern University, Sri Lanka, Chenkalady, Sri Lanka
Correspondence e-mail: eesans@yahoo.com


Received 19 March 2012; accepted 9 April 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.030; wR factor = 0.088; data-to-parameter ratio = 5.9.

The title compound, $C_{21}H_{17}N_3O_5 \cdot H_2O$, exists in the keto form with an *E* conformation with respect to the azomethine double bond. The twist angles between the aromatic rings are in the range $4.67(10)$ – $17.54(10)^\circ$. A water molecule of solvation is present in the lattice. A conventional intramolecular O—H···N hydrogen bond increases the rigidity of the molecule. Intermolecular O—H···O, N—H···O and C—H···O hydrogen-bonding interactions establish a supramolecular linkage among the molecules in the crystal structure. There are also C—H···π interactions present.

Related literature

For the biological and other applications of carbohydrazides, see: Lakshmi *et al.* (2011); Grande *et al.* (2007); Naseema *et al.* (2010). For the synthesis, see: Emmanuel *et al.* (2011). For related structures of carbohydrazides, see: Fun *et al.* (2008). For the keto form, see: Bakir & Brown (2002).



Experimental

Crystal data

$C_{21}H_{17}N_3O_5 \cdot H_2O$
 $M_r = 409.39$
Monoclinic, Pc
 $a = 4.6275(7)\text{ \AA}$
 $b = 6.5332(11)\text{ \AA}$
 $c = 31.856(5)\text{ \AA}$
 $\beta = 92.417(4)^\circ$
 $V = 962.2(3)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.11\text{ mm}^{-1}$

$T = 296\text{ K}$
 $0.30 \times 0.28 \times 0.25\text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2004)
 $T_{\min} = 0.969$, $T_{\max} = 0.974$

7388 measured reflections
1713 independent reflections
1663 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.088$
 $S = 1.05$
1713 reflections
288 parameters
7 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.16\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

Cg is the centroid of the C1–C6 ring

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4O···O1S ⁱ	0.86 (2)	2.57 (4)	3.058 (3)	117 (4)
O4—H4O···N3	0.86 (2)	1.93 (3)	2.670 (2)	143 (4)
N2—H2N···O1S	0.86 (2)	2.01 (2)	2.855 (3)	166 (3)
O1S—H2S···O4 ⁱⁱ	0.87 (2)	2.01 (2)	2.860 (3)	165 (4)
O1S—H1S···O3 ⁱⁱⁱ	0.88 (2)	1.80 (2)	2.676 (3)	175 (5)
C14—H14···O1S	0.93	2.37	3.184 (2)	146
C21—H21···O1S	0.93	2.43	3.325 (3)	161
C7—H7B···Cg ^{iv}	0.97	2.69	3.472 (2)	138

Symmetry codes: (i) $x, y + 1, z$; (ii) $x - 1, y - 1, z$; (iii) $x, y - 1, z$; (iv) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2010); software used to prepare material for publication: *SHELXL97* and *publCIF* (Westrip, 2010).

BJ is grateful to the Council for Scientific and Industrial Research, New Delhi, India, for the award of a Senior Research Fellowship. The authors also thank the Sophisticated Analytical Instrument Facility, Cochin University of Science & Technology, Kochi-22, for providing the single-crystal X-ray diffraction data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2537).

References

- Bakir, M. & Brown, O. (2002). *J. Mol. Struct.* **609**, 129–136.
- Brandenburg, K. (2010). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *APEX2*, *SAINT* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Emmanuel, J., Sithambaresan, M. & Kurup, M. R. P. (2011). *Acta Cryst. E67*, o3267.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fun, H.-K., Patil, P. S., Jebas, S. R., Sujith, K. V. & Kalluraya, B. (2008). *Acta Cryst. E64*, o1594–o1595.
- Grande, F., Aiello, F., Grazia, O. D., Brizzi, A., Garofalo, A. & Neamati, N. (2007). *Bioorg. Med. Chem.* **15**, 288–294.

organic compounds

- Lakshmi, B., Avaji, P. G., Shivananda, K. N., Nagella, P., Manohar, S. H. & Mahendra, K. N. (2011). *Polyhedron*, **30**, 1507–1515.
- Naseema, K., Sujith, K. V., Manjunatha, K. B., Kalluraya, B., Umesh, G. & Rao, V. (2010). *Opt. Laser Technol.* **42**, 741–748.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o1421–o1422 [doi:10.1107/S1600536812015401]

N'-[*(E*)-4-Benzylxy-2-hydroxybenzylidene]-4-nitrobenzohydrazide monohydrate

Bibitha Joseph, M. Sithambaresan and M. R. Prathapachandra Kurup

S1. Comment

There is growing interest in the structural features of carbohydrazides as they show a wide range of biological activities, with potential uses in antibacterial, antifungal and anticancer studies (Lakshmi *et al.*, 2011; Grande *et al.*, 2007).

Hydrazones and their metal complexes have found applications in chemical processes like non linear optics, sensors *etc.* (Naseema *et al.*, 2010).

The title compound *N'*-{*(E*)-[4-(benzylxy)-2-hydroxyphenyl]methylidene}-4-nitrobenzohydrazide hydrate is found to exist in the *E* configuration with respect to N3=C14 bond. A perspective view of the molecular structure of the title compound, along with the atom-labeling is shown in Fig. 1. The bond length of C15=O3 [1.229 (2) Å] shows a significant double-bond character (Fun *et al.*, 2008) indicating that the molecule exists in the keto form in the solid state (Bakir & Brown, 2002) and the dihedral angles between the aromatic rings are in the range of 4.67 (10)–17.54 (10)°.

The lattice water molecule plays an essential role in packing of the molecules forming conventional and non-conventional hydrogen bonds between the carbohydrazide and water molecules (Fig. 2). A C–H···π interaction is also observed in the crystal structure between one of the H atoms attached to the C7 carbon atom and the phenyl ring of the adjacent molecule in the crystal system (Fig. 3). Two types of very weak π–π interactions also present with a shortest centroid-centroid distance of 4.9302 (14) Å. In crystal packing, the parallel arrangement of the molecules along *a* axis is shown in Fig. 4.


S2. Experimental

The title compound was prepared by adapting a reported procedure (Emmanuel *et al.*, 2011) by refluxing a mixture of methanolic solutions of 4-nitrobenzohydrazide (0.181 g, 1 mmol) and 4-(benzylxy)-2-hydroxybenzaldehyde (0.228 g, 1 mmol) for 4 h. The formed crystals were collected, washed with few drops of methanol and dried over P₄O₁₀ *in vacuo*. Single crystals of the title compound suitable for X-ray analysis were obtained by slow evaporation from its methanolic solution.


S3. Refinement

All H atoms on C were placed in calculated positions, guided by difference maps, with C—H bond distances 0.93–0.97 Å. H atoms were assigned as $U_{\text{iso}}=1.2 U_{\text{eq}}$. N2—H2N and O4—H4O (0.86 Å) H atoms were located from difference maps and restrained using *DFIX* instructions. The O1S—H1S and O1S—H2S (0.86 Å) H atoms of the water molecule is also located from difference maps and restrained using *DFIX* and *DANG* instructions.


In the absence of significant anomalous scattering effects Friedel pairs have been merged.

**Figure 1**


ORTEP view of the molecular structure of the title compound, along with the atom-labelling, drawn with 50% probability displacement ellipsoids.

**Figure 2**

A view of the conventional and non-conventional hydrogen bonding interactions

**Figure 3**

Graphical representation showing C–H···π interaction in the crystal structure of $C_{21}H_{17}N_3O_5 \cdot H_2O$.

**Figure 4**

Packing arrangement of molecules along a axis.

N'-(E)-4-Benzyl-2-hydroxybenzylidene]-4-nitrobenzohydrazide monohydrate*Crystal data* $M_r = 409.39$ Monoclinic, Pc

Hall symbol: P -2yc

 $a = 4.6275 (7) \text{ \AA}$ $b = 6.5332 (11) \text{ \AA}$ $c = 31.856 (5) \text{ \AA}$ $\beta = 92.417 (4)^\circ$ $V = 962.2 (3) \text{ \AA}^3$ $Z = 2$ $F(000) = 428.0$ $D_x = 1.413 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4698 reflections

 $\theta = 3.1\text{--}28.2^\circ$ $\mu = 0.11 \text{ mm}^{-1}$ $T = 296 \text{ K}$

Block, yellow

 $0.30 \times 0.28 \times 0.25 \text{ mm}$ *Data collection*

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω and φ scanAbsorption correction: multi-scan
(*SADABS*; Bruker, 2004) $T_{\min} = 0.969$, $T_{\max} = 0.974$

7388 measured reflections

1713 independent reflections

1663 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.020$ $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 3.1^\circ$ $h = -3 \rightarrow 5$ $k = -7 \rightarrow 7$ $l = -37 \rightarrow 37$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.030$ $wR(F^2) = 0.088$ $S = 1.05$

1713 reflections

288 parameters

7 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0687P)^2 + 0.0484P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.011$ $\Delta\rho_{\max} = 0.14 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.16 \text{ e \AA}^{-3}$ Extinction correction: *SHELXL97* (Sheldrick, 2008), $Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.021 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.5580 (7)	0.1667 (5)	0.47190 (9)	0.0986 (10)
O2	-0.5671 (7)	0.4470 (5)	0.50699 (9)	0.0960 (9)
O3	0.3371 (5)	0.9071 (3)	0.37071 (7)	0.0673 (6)

O4	0.8452 (4)	1.0067 (3)	0.26751 (6)	0.0537 (5)
O5	1.4446 (4)	0.8132 (2)	0.15654 (5)	0.0428 (4)
O1S	0.3303 (5)	0.1942 (3)	0.31076 (7)	0.0596 (5)
N1	-0.4868 (6)	0.3450 (5)	0.47795 (8)	0.0623 (7)
N2	0.3751 (4)	0.6173 (3)	0.33257 (6)	0.0380 (4)
N3	0.5600 (4)	0.7064 (3)	0.30472 (6)	0.0370 (4)
C1	1.8112 (6)	0.8824 (4)	0.09200 (9)	0.0507 (6)
H1	1.7528	0.7502	0.0983	0.061*
C2	2.0020 (6)	0.9151 (5)	0.05996 (9)	0.0593 (8)
H2	2.0703	0.8044	0.0449	0.071*
C3	2.0894 (6)	1.1089 (5)	0.05048 (9)	0.0589 (8)
H3	2.2186	1.1292	0.0293	0.071*
C4	1.9873 (6)	1.2742 (5)	0.07214 (9)	0.0600 (7)
H4	2.0454	1.4061	0.0654	0.072*
C5	1.7976 (6)	1.2430 (4)	0.10394 (9)	0.0508 (6)
H5	1.7282	1.3546	0.1186	0.061*
C6	1.7094 (4)	1.0458 (4)	0.11423 (7)	0.0377 (5)
C7	1.5050 (5)	1.0238 (4)	0.14926 (7)	0.0391 (5)
H7A	1.5901	1.0846	0.1747	0.047*
H7B	1.3264	1.0957	0.1421	0.047*
C8	1.2534 (4)	0.7701 (4)	0.18637 (7)	0.0355 (5)
C9	1.1417 (5)	0.9148 (3)	0.21297 (7)	0.0371 (5)
H9	1.1963	1.0513	0.2108	0.044*
C10	0.9484 (4)	0.8566 (4)	0.24291 (7)	0.0345 (5)
C11	0.8636 (4)	0.6529 (4)	0.24699 (7)	0.0346 (5)
C12	0.9782 (5)	0.5107 (4)	0.21932 (8)	0.0430 (5)
H12	0.9233	0.3742	0.2212	0.052*
C13	1.1688 (5)	0.5665 (4)	0.18961 (8)	0.0434 (5)
H13	1.2415	0.4686	0.1717	0.052*
C14	0.6665 (5)	0.5843 (4)	0.27812 (7)	0.0388 (5)
H14	0.6156	0.4467	0.2788	0.047*
C15	0.2747 (5)	0.7263 (4)	0.36431 (8)	0.0405 (5)
C16	0.0762 (5)	0.6172 (4)	0.39327 (7)	0.0382 (5)
C17	-0.0090 (6)	0.7250 (4)	0.42798 (8)	0.0522 (6)
H17	0.0583	0.8576	0.4326	0.063*
C18	-0.1943 (6)	0.6370 (5)	0.45600 (8)	0.0567 (7)
H18	-0.2526	0.7092	0.4793	0.068*
C19	-0.2898 (5)	0.4403 (4)	0.44837 (7)	0.0452 (6)
C20	-0.2096 (6)	0.3294 (4)	0.41418 (9)	0.0524 (6)
H20	-0.2769	0.1967	0.4098	0.063*
C21	-0.0266 (6)	0.4198 (4)	0.38649 (8)	0.0494 (6)
H21	0.0285	0.3474	0.3630	0.059*
H1S	0.325 (10)	0.105 (6)	0.3313 (11)	0.096 (14)*
H2S	0.169 (6)	0.160 (6)	0.2975 (12)	0.099 (14)*
H2N	0.336 (6)	0.489 (3)	0.3287 (9)	0.046 (7)*
H4O	0.730 (7)	0.958 (7)	0.2853 (11)	0.090 (13)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.123 (2)	0.097 (2)	0.0791 (17)	-0.0492 (18)	0.0467 (15)	-0.0001 (15)
O2	0.1081 (19)	0.108 (2)	0.0771 (16)	-0.0053 (17)	0.0631 (14)	-0.0006 (15)
O3	0.1030 (17)	0.0439 (11)	0.0571 (11)	-0.0205 (10)	0.0283 (10)	-0.0053 (9)
O4	0.0609 (11)	0.0409 (10)	0.0623 (11)	-0.0109 (8)	0.0384 (9)	-0.0097 (8)
O5	0.0446 (8)	0.0385 (9)	0.0470 (9)	0.0009 (8)	0.0217 (7)	0.0040 (7)
O1S	0.0790 (14)	0.0383 (9)	0.0638 (12)	-0.0155 (9)	0.0283 (10)	-0.0050 (9)
N1	0.0571 (14)	0.0820 (19)	0.0491 (12)	-0.0043 (13)	0.0172 (11)	0.0143 (13)
N2	0.0434 (10)	0.0318 (10)	0.0400 (10)	-0.0066 (8)	0.0150 (8)	0.0050 (8)
N3	0.0357 (10)	0.0364 (10)	0.0398 (10)	-0.0075 (8)	0.0127 (7)	0.0060 (8)
C1	0.0523 (14)	0.0453 (14)	0.0560 (15)	-0.0009 (12)	0.0202 (12)	0.0008 (11)
C2	0.0609 (17)	0.065 (2)	0.0547 (15)	0.0085 (14)	0.0296 (13)	-0.0029 (13)
C3	0.0500 (14)	0.081 (2)	0.0477 (14)	0.0040 (14)	0.0227 (11)	0.0143 (15)
C4	0.0598 (16)	0.0594 (17)	0.0624 (17)	-0.0058 (14)	0.0212 (13)	0.0237 (15)
C5	0.0517 (14)	0.0444 (14)	0.0579 (14)	0.0025 (11)	0.0195 (11)	0.0093 (12)
C6	0.0303 (10)	0.0455 (12)	0.0378 (11)	0.0008 (10)	0.0089 (9)	0.0078 (10)
C7	0.0378 (11)	0.0382 (12)	0.0420 (12)	0.0016 (10)	0.0118 (9)	0.0024 (10)
C8	0.0334 (11)	0.0391 (12)	0.0346 (10)	0.0020 (9)	0.0097 (8)	0.0037 (9)
C9	0.0390 (11)	0.0304 (11)	0.0426 (12)	-0.0034 (9)	0.0106 (9)	0.0016 (9)
C10	0.0353 (11)	0.0327 (12)	0.0362 (11)	-0.0026 (9)	0.0100 (8)	-0.0011 (9)
C11	0.0312 (10)	0.0356 (12)	0.0374 (11)	-0.0039 (9)	0.0066 (9)	0.0019 (9)
C12	0.0457 (12)	0.0300 (11)	0.0542 (14)	-0.0021 (10)	0.0139 (10)	0.0002 (10)
C13	0.0454 (12)	0.0365 (12)	0.0497 (12)	0.0018 (10)	0.0174 (10)	-0.0052 (10)
C14	0.0382 (11)	0.0338 (12)	0.0452 (12)	-0.0049 (9)	0.0102 (10)	0.0073 (9)
C15	0.0459 (12)	0.0373 (12)	0.0389 (11)	-0.0050 (10)	0.0095 (9)	0.0028 (10)
C16	0.0393 (11)	0.0419 (12)	0.0341 (11)	0.0032 (10)	0.0082 (9)	0.0054 (9)
C17	0.0611 (16)	0.0511 (15)	0.0455 (13)	-0.0088 (13)	0.0140 (11)	-0.0051 (12)
C18	0.0648 (16)	0.0659 (17)	0.0409 (13)	-0.0006 (14)	0.0201 (11)	-0.0074 (13)
C19	0.0388 (12)	0.0614 (17)	0.0362 (12)	0.0002 (11)	0.0100 (9)	0.0102 (11)
C20	0.0569 (15)	0.0443 (14)	0.0575 (15)	-0.0059 (12)	0.0198 (12)	0.0053 (12)
C21	0.0580 (15)	0.0452 (14)	0.0468 (13)	-0.0061 (12)	0.0232 (11)	-0.0028 (11)

Geometric parameters (\AA , $^\circ$)

O1—N1	1.224 (4)	C6—C7	1.500 (3)
O2—N1	1.211 (4)	C7—H7A	0.9700
O3—C15	1.231 (3)	C7—H7B	0.9700
O4—C10	1.355 (3)	C8—C9	1.384 (3)
O4—H4O	0.86 (2)	C8—C13	1.392 (4)
O5—C8	1.355 (3)	C9—C10	1.388 (3)
O5—C7	1.425 (3)	C9—H9	0.9300
O1S—H1S	0.88 (2)	C10—C11	1.395 (3)
O1S—H2S	0.873 (19)	C11—C12	1.400 (3)
N1—C19	1.476 (3)	C11—C14	1.446 (3)
N2—C15	1.336 (3)	C12—C13	1.370 (3)
N2—N3	1.386 (3)	C12—H12	0.9300

N2—H2N	0.863 (19)	C13—H13	0.9300
N3—C14	1.277 (3)	C14—H14	0.9300
C1—C6	1.375 (4)	C15—C16	1.508 (3)
C1—C2	1.394 (4)	C16—C17	1.383 (4)
C1—H1	0.9300	C16—C21	1.388 (4)
C2—C3	1.367 (5)	C17—C18	1.388 (4)
C2—H2	0.9300	C17—H17	0.9300
C3—C4	1.376 (5)	C18—C19	1.377 (4)
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.383 (4)	C19—C20	1.372 (4)
C4—H4	0.9300	C20—C21	1.381 (4)
C5—C6	1.394 (3)	C20—H20	0.9300
C5—H5	0.9300	C21—H21	0.9300
C10—O4—H4O	111 (3)	C8—C9—H9	119.9
C8—O5—C7	116.93 (17)	C10—C9—H9	119.9
H1S—O1S—H2S	98 (3)	O4—C10—C9	116.9 (2)
O2—N1—O1	123.7 (3)	O4—C10—C11	122.0 (2)
O2—N1—C19	118.2 (3)	C9—C10—C11	121.2 (2)
O1—N1—C19	118.0 (3)	C10—C11—C12	117.30 (19)
C15—N2—N3	120.28 (19)	C10—C11—C14	123.2 (2)
C15—N2—H2N	123 (2)	C12—C11—C14	119.5 (2)
N3—N2—H2N	117 (2)	C13—C12—C11	122.0 (2)
C14—N3—N2	115.35 (19)	C13—C12—H12	119.0
C6—C1—C2	120.0 (3)	C11—C12—H12	119.0
C6—C1—H1	120.0	C12—C13—C8	119.8 (2)
C2—C1—H1	120.0	C12—C13—H13	120.1
C3—C2—C1	120.4 (3)	C8—C13—H13	120.1
C3—C2—H2	119.8	N3—C14—C11	122.3 (2)
C1—C2—H2	119.8	N3—C14—H14	118.9
C2—C3—C4	120.3 (2)	C11—C14—H14	118.9
C2—C3—H3	119.8	O3—C15—N2	123.4 (2)
C4—C3—H3	119.8	O3—C15—C16	119.8 (2)
C3—C4—C5	119.6 (3)	N2—C15—C16	116.8 (2)
C3—C4—H4	120.2	C17—C16—C21	119.4 (2)
C5—C4—H4	120.2	C17—C16—C15	116.8 (2)
C4—C5—C6	120.6 (3)	C21—C16—C15	123.9 (2)
C4—C5—H5	119.7	C16—C17—C18	120.5 (3)
C6—C5—H5	119.7	C16—C17—H17	119.7
C1—C6—C5	119.1 (2)	C18—C17—H17	119.7
C1—C6—C7	123.3 (2)	C19—C18—C17	118.4 (2)
C5—C6—C7	117.6 (2)	C19—C18—H18	120.8
O5—C7—C6	110.37 (19)	C17—C18—H18	120.8
O5—C7—H7A	109.6	C20—C19—C18	122.5 (2)
C6—C7—H7A	109.6	C20—C19—N1	118.6 (2)
O5—C7—H7B	109.6	C18—C19—N1	118.9 (2)
C6—C7—H7B	109.6	C19—C20—C21	118.4 (2)
H7A—C7—H7B	108.1	C19—C20—H20	120.8

O5—C8—C9	124.0 (2)	C21—C20—H20	120.8
O5—C8—C13	116.4 (2)	C20—C21—C16	120.9 (2)
C9—C8—C13	119.6 (2)	C20—C21—H21	119.6
C8—C9—C10	120.1 (2)	C16—C21—H21	119.6
C15—N2—N3—C14	173.2 (2)	O5—C8—C13—C12	179.4 (2)
C6—C1—C2—C3	0.1 (5)	C9—C8—C13—C12	-0.6 (4)
C1—C2—C3—C4	-0.8 (5)	N2—N3—C14—C11	-179.85 (19)
C2—C3—C4—C5	0.7 (5)	C10—C11—C14—N3	-1.3 (3)
C3—C4—C5—C6	0.1 (4)	C12—C11—C14—N3	178.5 (2)
C2—C1—C6—C5	0.6 (4)	N3—N2—C15—O3	0.1 (4)
C2—C1—C6—C7	-179.7 (3)	N3—N2—C15—C16	-179.66 (19)
C4—C5—C6—C1	-0.7 (4)	O3—C15—C16—C17	-4.8 (4)
C4—C5—C6—C7	179.6 (2)	N2—C15—C16—C17	175.0 (2)
C8—O5—C7—C6	-177.51 (18)	O3—C15—C16—C21	174.1 (3)
C1—C6—C7—O5	2.1 (3)	N2—C15—C16—C21	-6.2 (3)
C5—C6—C7—O5	-178.1 (2)	C21—C16—C17—C18	0.3 (4)
C7—O5—C8—C9	-7.7 (3)	C15—C16—C17—C18	179.2 (2)
C7—O5—C8—C13	172.3 (2)	C16—C17—C18—C19	0.2 (4)
O5—C8—C9—C10	-179.4 (2)	C17—C18—C19—C20	-0.3 (4)
C13—C8—C9—C10	0.5 (3)	C17—C18—C19—N1	180.0 (2)
C8—C9—C10—O4	-179.2 (2)	O2—N1—C19—C20	-176.7 (3)
C8—C9—C10—C11	0.2 (3)	O1—N1—C19—C20	3.9 (4)
O4—C10—C11—C12	178.5 (2)	O2—N1—C19—C18	3.0 (4)
C9—C10—C11—C12	-0.9 (3)	O1—N1—C19—C18	-176.4 (3)
O4—C10—C11—C14	-1.6 (3)	C18—C19—C20—C21	-0.1 (4)
C9—C10—C11—C14	179.0 (2)	N1—C19—C20—C21	179.6 (2)
C10—C11—C12—C13	0.8 (3)	C19—C20—C21—C16	0.6 (4)
C14—C11—C12—C13	-179.1 (2)	C17—C16—C21—C20	-0.7 (4)
C11—C12—C13—C8	-0.1 (4)	C15—C16—C21—C20	-179.6 (2)

Hydrogen-bond geometry (Å, °)

Cg is the centroid of the C1—C6 ring

D—H···A	D—H	H···A	D···A	D—H···A
O4—H4O···O1 <i>S</i> ⁱ	0.86 (2)	2.57 (4)	3.058 (3)	117 (4)
O4—H4 <i>O</i> ···N3	0.86 (2)	1.93 (3)	2.670 (2)	143 (4)
N2—H2 <i>N</i> ···O1 <i>S</i>	0.86 (2)	2.01 (2)	2.855 (3)	166 (3)
O1 <i>S</i> —H2 <i>S</i> ···O4 ⁱⁱ	0.87 (2)	2.01 (2)	2.860 (3)	165 (4)
O1 <i>S</i> —H1 <i>S</i> ···O3 ⁱⁱⁱ	0.88 (2)	1.80 (2)	2.676 (3)	175 (5)
C14—H14···O1 <i>S</i>	0.93	2.37	3.184 (2)	146
C21—H21···O1 <i>S</i>	0.93	2.43	3.325 (3)	161
C7—H7 <i>B</i> ···Cg ^{iv}	0.97	2.69	3.472 (2)	138

Symmetry codes: (i) $x, y+1, z$; (ii) $x-1, y-1, z$; (iii) $x, y-1, z$; (iv) $x-1, y, z$.