

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(3-methylanilinium) hexachlorido-stannate(IV) dihydrate

Ming-Liang Liu

Ordered Matter Science Research Center, Southeast University, Nanjing 211189, People's Republic of China

Correspondence e-mail: jgsdxlm@163.com


Received 11 April 2012; accepted 20 April 2012

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.030; wR factor = 0.096; data-to-parameter ratio = 19.0.

In the title compound, $(\text{C}_7\text{H}_{10}\text{N})_2[\text{SnCl}_6] \cdot 2\text{H}_2\text{O}$, the Sn^{IV} atom lies on a site with symmetry $2/m$. One of the Cl atoms lies on a mirror plane and the 3-methylanilinium cation is also situated on a mirror plane. The water molecule is located on a twofold rotation axis. The H atoms of the methyl and ammonium groups and the solvent water molecule are disordered by symmetry. In the crystal, $\text{N}-\text{H} \cdots \text{Cl}$, $\text{O}-\text{H} \cdots \text{Cl}$ and $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds connect the organic cations, the inorganic octahedrally shaped anions and the water molecules.

Related literature

For background to ferroelectric metal-organic complexes, see: Zhang *et al.* (2009, 2010). For related structures, see: Liu (2011a,b,c).



Experimental

Crystal data

 $(\text{C}_7\text{H}_{10}\text{N})_2[\text{SnCl}_6] \cdot 2\text{H}_2\text{O}$
 $M_r = 583.74$

 Monoclinic, $C2/m$
 $a = 20.467$ (4) Å
 $b = 7.1699$ (14) Å
 $c = 7.7569$ (16) Å
 $\beta = 93.83$ (3)°
 $V = 1135.8$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.84$ mm⁻¹
 $T = 293$ K
 $0.36 \times 0.32 \times 0.28$ mm

Data collection

 Rigaku Mercury2 CCD diffractometer
 Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $T_{\text{min}} = 0.963$, $T_{\text{max}} = 0.971$

 5833 measured reflections
 1405 independent reflections
 1370 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.096$
 $S = 0.92$
 1405 reflections

 74 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.73$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

| $D-H \cdots A$ | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|---|-------|--------------|--------------|----------------|
| $\text{N1}-\text{H1A} \cdots \text{Cl2}^{\text{i}}$ | 0.89 | 2.59 | 3.476 (4) | 171 |
| $\text{N1}-\text{H1B} \cdots \text{O1}^{\text{ii}}$ | 0.89 | 1.93 | 2.809 (5) | 170 |
| $\text{N1}-\text{H1C} \cdots \text{Cl1}^{\text{iii}}$ | 0.89 | 2.75 | 3.5883 (7) | 157 |
| $\text{O1}-\text{H1D} \cdots \text{Cl2}$ | 0.85 | 2.44 | 3.228 (2) | 154 |

 Symmetry codes: (i) $x + \frac{1}{2}, y + \frac{1}{2}, z - 1$; (ii) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The author thanks an anonymous advisor from Ordered Matter Science Research Centre, Southeast University, for great help in the revision of this paper.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2538).

References

- Liu, M.-L. (2011a). *Acta Cryst.* **E67**, m1622.
 Liu, M.-L. (2011b). *Acta Cryst.* **E67**, m1812.
 Liu, M.-L. (2011c). *Acta Cryst.* **E67**, m1827.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zhang, W., Chen, L.-Z., Xiong, R.-G., Nakamura, T. & Huang, S.-P. (2009). *J. Am. Chem. Soc.* **131**, 12544–12545.
 Zhang, W., Ye, H.-Y., Cai, H.-L., Ge, J.-Z., Xiong, R.-G. & Huang, S.-P. (2010). *J. Am. Chem. Soc.* **132**, 7300–7302.

supporting information

Acta Cryst. (2012). E68, m681 [doi:10.1107/S1600536812017618]

Bis(3-methylanilinium) hexachloridostannate(IV) dihydrate**Ming-Liang Liu****S1. Comment**

Recently much attention has been devoted to metal-organic compounds due to the tunability of their special structural features and their interesting physical properties (Zhang *et al.*, 2009, 2010). As a continuation of our researches (Liu, 2011a,b,c), the title compound has been synthesized and its crystal structure is herein reported.

In the title compound, the Sn^{IV} atom lies on a 2/m symmetry site, and is coordinated by six Cl atoms (Fig. 1). One of the Cl atoms lies on a mirror plane and the 3-methylanilinium cation is also situated on a mirror plane. The water molecule is located on a twofold rotation axis. The H atoms of the methyl and amidogen groups and the water molecule are disordered induced by symmetry. N—H···Cl, O—H···Cl and N—H···O hydrogen-bonding interactions connect the [SnCl₆]²⁻ anions, the 3-methylanilinium cations and the water molecules (Table 1). The non-H atoms of the 3-methylanilinium cation are coplanar. The average Sn—Cl bond distances range from 2.4260 (13) to 2.4384 (9) Å and the *cis* Cl—Sn—Cl angles range from 88.78 (5) to 91.22 (5)°.


S2. Experimental

3-Methylbenzenamine (3.21 g, 0.03 mol) was dissolved in 30 ml ethanol, to which hydrochloric acid (1.1 g, 0.03 mol) was then added. Stannous chloride (2.25 g, 0.01 mol) was dissolved in 20 ml ethanol, to which was added hydrochloric acid, then mixed with the above solution without any precipitation under stirring at ambient temperature. Single crystals suitable for X-ray structure analysis were obtained by slow evaporation after 4 days in air.

The dielectric constant of the compound as a function of temperature indicates that the permittivity is basically temperature-independent [$\epsilon = C/(T-T_0)$], suggesting that this compound is not ferroelectric or there may be no distinct phase transition occurring within the measured temperature (below the melting point).

S3. Refinement

H atoms on C atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 (aromatic), 0.96 (methyl) and N—H = 0.89 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. Water H atoms were located from a difference Fourier map and refined as riding atoms, with O—H = 0.85 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

**Figure 1**

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms on C7, N1 and O1 are disordered over two sets of sites. [Symmetry codes: (A) $-x, -y, 2-z$; (B) $-x, y, 2-z$; (C) $x, -y, z$.]

Bis(3-methylanilinium) hexachloridostannate(IV) dihydrate

Crystal data

$(C_7H_{10}N)_2[SnCl_6] \cdot 2H_2O$

$M_r = 583.74$

Monoclinic, $C2/m$

Hall symbol: $-C 2y$

$a = 20.467 (4) \text{ \AA}$

$b = 7.1699 (14) \text{ \AA}$

$c = 7.7569 (16) \text{ \AA}$

$\beta = 93.83 (3)^\circ$

$V = 1135.8 (4) \text{ \AA}^3$

$Z = 2$

$F(000) = 580$

$D_x = 1.707 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1370 reflections

$\theta = 3.4\text{--}25.0^\circ$

$\mu = 1.84 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colourless

$0.36 \times 0.32 \times 0.28 \text{ mm}$

Data collection

Rigaku Mercury2 CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.963, T_{\max} = 0.971$

5833 measured reflections

1405 independent reflections

1370 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.0^\circ$

$h = -26 \rightarrow 25$

$k = -9 \rightarrow 9$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.096$

$S = 0.92$

1405 reflections

74 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0749P)^2 + 1.7826P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.73 \text{ e } \text{Å}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001x Fc^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.041 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ | Occ. (<1) |
|-----|--------------|--------------|--------------|----------------------------------|-----------|
| N1 | 0.4203 (2) | 1.0000 | 0.2314 (5) | 0.0565 (10) | |
| H1A | 0.4253 | 0.9464 | 0.1296 | 0.068* | 0.50 |
| H1B | 0.4429 | 0.9367 | 0.3140 | 0.068* | 0.50 |
| H1C | 0.4349 | 1.1169 | 0.2294 | 0.068* | 0.50 |
| C1 | 0.3335 (2) | 1.0000 | 0.4366 (5) | 0.0416 (8) | |
| H1 | 0.3661 | 1.0000 | 0.5261 | 0.050* | |
| C2 | 0.3501 (2) | 1.0000 | 0.2664 (5) | 0.0392 (8) | |
| C3 | 0.3029 (2) | 1.0000 | 0.1309 (5) | 0.0537 (11) | |
| H3 | 0.3147 | 1.0000 | 0.0171 | 0.064* | |
| C4 | 0.2382 (2) | 1.0000 | 0.1673 (6) | 0.0618 (13) | |
| H4 | 0.2058 | 1.0000 | 0.0774 | 0.074* | |
| C5 | 0.2208 (2) | 1.0000 | 0.3365 (7) | 0.0542 (11) | |
| H5 | 0.1767 | 1.0000 | 0.3588 | 0.065* | |
| C6 | 0.2677 (2) | 1.0000 | 0.4732 (5) | 0.0429 (8) | |
| C7 | 0.2496 (3) | 1.0000 | 0.6588 (7) | 0.0622 (13) | |
| H7A | 0.2819 | 1.0687 | 0.7285 | 0.075* | 0.50 |
| H7B | 0.2481 | 0.8739 | 0.7000 | 0.075* | 0.50 |
| H7C | 0.2075 | 1.0574 | 0.6659 | 0.075* | 0.50 |
| Sn1 | 0.0000 | 0.0000 | 1.0000 | 0.0370 (2) | |
| Cl1 | 0.07220 (7) | 0.0000 | 0.76437 (18) | 0.0617 (3) | |
| Cl2 | -0.06652 (4) | 0.24302 (12) | 0.85527 (11) | 0.0565 (3) | |
| O1 | 0.0000 | 0.3385 (11) | 0.5000 | 0.136 (3) | |
| H1D | -0.0216 | 0.3511 | 0.5891 | 0.203* | 0.50 |
| H1E | 0.0390 | 0.3062 | 0.5304 | 0.203* | 0.50 |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|-------------|-----------|-------------|----------|--------------|----------|
| N1 | 0.0412 (19) | 0.081 (3) | 0.0478 (19) | 0.000 | 0.0093 (15) | 0.000 |
| C1 | 0.0394 (19) | 0.050 (2) | 0.0354 (17) | 0.000 | -0.0004 (14) | 0.000 |
| C2 | 0.0346 (18) | 0.046 (2) | 0.0368 (17) | 0.000 | 0.0041 (14) | 0.000 |
| C3 | 0.054 (3) | 0.072 (3) | 0.0344 (18) | 0.000 | -0.0035 (17) | 0.000 |

| | | | | | | |
|-----|------------|-------------|-------------|------------|--------------|------------|
| C4 | 0.045 (2) | 0.085 (4) | 0.053 (2) | 0.000 | -0.0157 (19) | 0.000 |
| C5 | 0.035 (2) | 0.063 (3) | 0.064 (3) | 0.000 | 0.0028 (18) | 0.000 |
| C6 | 0.047 (2) | 0.0370 (19) | 0.0458 (19) | 0.000 | 0.0089 (16) | 0.000 |
| C7 | 0.070 (3) | 0.065 (3) | 0.054 (2) | 0.000 | 0.025 (2) | 0.000 |
| Sn1 | 0.0290 (2) | 0.0291 (2) | 0.0536 (3) | 0.000 | 0.00791 (15) | 0.000 |
| Cl1 | 0.0550 (7) | 0.0597 (7) | 0.0742 (7) | 0.000 | 0.0323 (6) | 0.000 |
| Cl2 | 0.0503 (4) | 0.0477 (4) | 0.0712 (5) | 0.0134 (3) | 0.0023 (3) | 0.0088 (4) |
| O1 | 0.127 (5) | 0.173 (7) | 0.105 (4) | 0.000 | -0.009 (4) | 0.000 |

Geometric parameters (Å, °)

| | | | |
|------------|-----------|---|-------------|
| N1—C2 | 1.480 (5) | C5—H5 | 0.9300 |
| N1—H1A | 0.8900 | C6—C7 | 1.511 (6) |
| N1—H1B | 0.8899 | C7—H7A | 0.9602 |
| N1—H1C | 0.8901 | C7—H7B | 0.9600 |
| C1—C2 | 1.385 (5) | C7—H7C | 0.9600 |
| C1—C6 | 1.395 (6) | Sn1—Cl1 ⁱ | 2.4260 (13) |
| C1—H1 | 0.9300 | Sn1—Cl1 | 2.4260 (13) |
| C2—C3 | 1.380 (6) | Sn1—Cl2 ⁱⁱ | 2.4384 (9) |
| C3—C4 | 1.372 (7) | Sn1—Cl2 ⁱⁱⁱ | 2.4384 (9) |
| C3—H3 | 0.9300 | Sn1—Cl2 | 2.4384 (9) |
| C4—C5 | 1.383 (7) | Sn1—Cl2 ⁱ | 2.4384 (9) |
| C4—H4 | 0.9300 | O1—H1D | 0.8500 |
| C5—C6 | 1.383 (7) | O1—H1E | 0.8499 |
| C2—N1—H1A | 109.5 | C1—C6—C7 | 119.7 (4) |
| C2—N1—H1B | 109.4 | C6—C7—H7A | 109.5 |
| H1A—N1—H1B | 109.5 | C6—C7—H7B | 109.5 |
| C2—N1—H1C | 109.5 | H7A—C7—H7B | 109.4 |
| H1A—N1—H1C | 109.5 | C6—C7—H7C | 109.4 |
| H1B—N1—H1C | 109.5 | H7A—C7—H7C | 109.5 |
| C2—C1—C6 | 119.7 (4) | H7B—C7—H7C | 109.5 |
| C2—C1—H1 | 120.2 | Cl1 ⁱ —Sn1—Cl1 | 180.0 |
| C6—C1—H1 | 120.2 | Cl1 ⁱ —Sn1—Cl2 ⁱⁱ | 89.85 (4) |
| C3—C2—C1 | 121.5 (4) | Cl1—Sn1—Cl2 ⁱⁱ | 90.15 (3) |
| C3—C2—N1 | 119.9 (4) | Cl1 ⁱ —Sn1—Cl2 ⁱⁱⁱ | 90.15 (4) |
| C1—C2—N1 | 118.5 (4) | Cl1—Sn1—Cl2 ⁱⁱⁱ | 89.85 (4) |
| C2—C3—C4 | 118.6 (4) | Cl2 ⁱⁱ —Sn1—Cl2 ⁱⁱⁱ | 180.0 |
| C2—C3—H3 | 120.7 | Cl1 ⁱ —Sn1—Cl2 | 89.85 (3) |
| C4—C3—H3 | 120.7 | Cl1—Sn1—Cl2 | 90.15 (3) |
| C5—C4—C3 | 120.7 (4) | Cl2 ⁱⁱ —Sn1—Cl2 | 91.22 (5) |
| C5—C4—H4 | 119.7 | Cl2 ⁱⁱⁱ —Sn1—Cl2 | 88.78 (5) |
| C3—C4—H4 | 119.7 | Cl1 ⁱ —Sn1—Cl2 ⁱ | 90.15 (3) |
| C6—C5—C4 | 121.1 (4) | Cl1—Sn1—Cl2 ⁱ | 89.85 (3) |
| C6—C5—H5 | 119.5 | Cl2 ⁱⁱ —Sn1—Cl2 ⁱ | 88.78 (5) |
| C4—C5—H5 | 119.5 | Cl2 ⁱⁱⁱ —Sn1—Cl2 ⁱ | 91.22 (5) |
| C5—C6—C1 | 118.4 (4) | Cl2—Sn1—Cl2 ⁱ | 180.0 |
| C5—C6—C7 | 121.9 (4) | H1D—O1—H1E | 109.5 |

| | | | |
|-------------|-------|-------------|-------|
| C6—C1—C2—C3 | 0.0 | C3—C4—C5—C6 | 0.0 |
| C6—C1—C2—N1 | 180.0 | C4—C5—C6—C1 | 0.0 |
| C1—C2—C3—C4 | 0.0 | C4—C5—C6—C7 | 180.0 |
| N1—C2—C3—C4 | 180.0 | C2—C1—C6—C5 | 0.0 |
| C2—C3—C4—C5 | 0.0 | C2—C1—C6—C7 | 180.0 |

Symmetry codes: (i) $-x, -y, -z+2$; (ii) $x, -y, z$; (iii) $-x, y, -z+2$.

Hydrogen-bond geometry (Å, °)

| <i>D—H...A</i> | <i>D—H</i> | <i>H...A</i> | <i>D...A</i> | <i>D—H...A</i> |
|----------------------------|------------|--------------|--------------|----------------|
| N1—H1A...Cl2 ^{iv} | 0.89 | 2.59 | 3.476 (4) | 171 |
| N1—H1B...O1 ^v | 0.89 | 1.93 | 2.809 (5) | 170 |
| N1—H1C...Cl1 ^{vi} | 0.89 | 2.75 | 3.5883 (7) | 157 |
| O1—H1D...Cl2 | 0.85 | 2.44 | 3.228 (2) | 154 |

Symmetry codes: (iv) $x+1/2, y+1/2, z-1$; (v) $x+1/2, y+1/2, z$; (vi) $-x+1/2, -y+3/2, -z+1$.