

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(Anthracen-1-yl)pyrrolidine-2,5-dione

Sanaz Khorasani and Manuel A. Fernandes*

 Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, PO Wits 2050, Johannesburg, South Africa
 Correspondence e-mail: manuel.fernandes@wits.ac.za


Received 2 March 2012; accepted 8 April 2012

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.041; wR factor = 0.088; data-to-parameter ratio = 8.8.

In the molecular structure of title compound, $\text{C}_{18}\text{H}_{13}\text{NO}_2$, the succinimide ring is orientated away from the plane of the anthracene moiety by $71.94(4)^\circ$. The crystal structure features three different types of intermolecular interactions, *viz.* $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ bonds. Molecules along the b axis stack on each other as a result of $\pi-\pi$ interactions which have a centroid-centroid distance of $3.6780(15)$ Å, while $\text{C}-\text{H}\cdots\pi$ interactions are present between neighbouring stacks. Also, acting between the stacks are the $\text{C}-\text{H}\cdots\text{O}$ interactions between the aromatic H atoms of the anthracene and the O atoms of the succinimide.

Related literature

For studies of regio- and stereo-selectivity of substituted anthracenes in Diels-Alder reactions, see: Singh & Ningombom (2010); Alston *et al.* (1979); Meek *et al.* (1960); Kaplan & Conroy (1963); Verma & Singh (1977). For a study involving NMR experiments, see: Hubbard *et al.* (1992).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{13}\text{NO}_2$	$V = 1321.98(13)$ Å ³
$M_r = 275.29$	$Z = 4$
Orthorhombic, $Pna2_1$	Mo $K\alpha$ radiation
$a = 18.4179(9)$ Å	$\mu = 0.09$ mm ⁻¹
$b = 5.7697(4)$ Å	$T = 173$ K
$c = 12.4403(6)$ Å	$0.49 \times 0.15 \times 0.10$ mm

Data collection

Bruker APEXII CCD diffractometer	1667 independent reflections
10655 measured reflections	1258 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.081$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	1 restraint
$wR(F^2) = 0.088$	H-atom parameters constrained
$S = 0.95$	$\Delta\rho_{\text{max}} = 0.17$ e Å ⁻³
1667 reflections	$\Delta\rho_{\text{min}} = -0.19$ e Å ⁻³
190 parameters	

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C7-C12 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C2}-\text{H2}\cdots\text{O2}^{\text{i}}$	0.95	2.38	3.234 (3)	149
$\text{C6}-\text{H6}\cdots\text{O1}^{\text{ii}}$	0.95	2.49	3.357 (3)	152
$\text{C9}-\text{H9}\cdots\text{O2}^{\text{iii}}$	0.95	2.53	3.465 (3)	167
$\text{C13}-\text{H13}\cdots\text{O1}^{\text{iv}}$	0.95	2.70	3.471 (3)	139
$\text{C17}-\text{H17A}\cdots\text{Cg2}^{\text{v}}$	0.99	2.92	3.709 (3)	138

 Symmetry codes: (i) $x, y+1, z$; (ii) $-x, -y+2, z+\frac{1}{2}$; (iii) $x+\frac{1}{2}, -y+\frac{1}{2}, z$; (iv) $x, y-1, z$; (v) $x+\frac{1}{2}, -y+\frac{3}{2}, z$.

Data collection: APEX2 (Bruker 2005); cell refinement: SAINT (Bruker 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and SCHAKAL99 (Keller, 1999); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON (Spek, 2009).

This work was supported by the National Research Foundation, Pretoria (NRF, GUN 77122) and the University of the Witwatersrand.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2341).

References

- Alston, P. V., Ottenbrite, R. M. & Newby, J. (1979). *J. Org. Chem.* **44**, 4939–4943.
- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hubbard, J. L., Carl, J. M. III, Anderson, G. D. & Rankin, G. O. (1992). *J. Heterocycl. Chem.* **29**, 719–721.
- Kaplan, F. & Conroy, H. (1963). *J. Org. Chem.* **28**, 1593–1596.
- Keller, E. (1999). SCHAKAL99. University of Freiberg, Germany.
- Meek, J. S., Wilgus, D. R. & Dann, J. R. (1960). *J. Am. Chem. Soc.* **82**, 2566–2569.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Singh, M. D. & Ningombom, A. (2010). *Indian J. Chem. Sect. B*, **49**, 789–794.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Verma, S. M. & Singh, M. D. (1977). *J. Org. Chem.* **42**, 3736–3740.

supporting information

Acta Cryst. (2012). E68, o1503 [doi:10.1107/S160053681201536X]

1-(Anthracen-1-yl)pyrrolidine-2,5-dione

Sanaz Khorasani and Manuel A. Fernandes

S1. Comment

The compound anthracene has been known for a long time and its properties have been extensively studied. The regio- and stereo-selectivity of substituted anthracenes in Diels-Alder reactions have been investigated and reported (Alston *et al.*, 1979; Meek *et al.*, 1960; Kaplan & Conroy, 1963; Verma & Singh, 1977; Singh & Ningombom, 2010). A study of the title compound and 1-succinimidonaphthalene involving synthesis, NMR experiments and molecular mechanics has been reported by Hubbard *et al.* (1992).

Both the anthracene and succinimide moieties are planar but are tilted with respect to each other at an angle of $71.94(4)^\circ$ (Fig. 1). Two anthracene bond lengths – C1—C14 [1.430(3)Å] and C5—C14 [1.443(3)Å] – are significantly longer than the 1.39Å typical of aromatic rings. As a consequence the rings containing these have been flagged as having larger than average C6-ring C—C bond lengths by *PLATON* (Spek, 2009), suggesting that the succinimide group has a significant effect on the charge distribution within the anthracene ring. The crystal structure contains three different types of intermolecular interactions, these include C—H \cdots O, C—H \cdots π and π – π interactions (Fig. 2). The π – π interaction occurs over a Cg1 \cdots Cg2 distance of 3.678(2)Å between the rings defined by C1-C5/C14 (Cg1) and C7-C12 (Cg2). This leads to the stacking of molecules along *b* axis. Geometrical details of the C—H \cdots π and C—H \cdots O interactions are given in the Table 1.

S2. Experimental

The title compound was synthesized with very low yield (a few crystals) by reaction of 1-aminoanthracene (0.200 g, 1 mmol) with succinic anhydride (0.107 g, 1 mmol) in the presence of dioxane as a solvent (3 ml) by stirring at room temperature for a few hours. Thionyl chloride (3 ml) in dioxane (2 ml) was then slowly added to the reaction mixture at room temperature. The mixture was then kept at 323 K for 12 h, followed by neutralization of excess thionyl chloride by pouring the mixture into a beaker containing ice. This mixture was then filtered yielding a dark brown material, which after recrystallization by slow evaporation from chloroform yielded a few crystals suitable for analysis by X-ray diffraction.

S3. Refinement

All H atoms were positioned geometrically, and allowed to ride on their parent atoms, with C—H bond lengths of 0.95Å for aromatic H or 0.99Å for methylene H and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The 1449 Friedel pairs were merged during structure refinement.





Figure 1

The molecular structure of the title compound, showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.

**Figure 2**

C–H \cdots O, C–H \cdots π and π – π interactions in the structure of the title compound. The C–H \cdots π and π – π interactions are respectively indicated by dollar (\$) or hash (#) symbols.

1-(Anthracen-1-yl)pyrrolidine-2,5-dione

Crystal data

$C_{18}H_{13}NO_2$

$M_r = 275.29$

Orthorhombic, $Pna2_1$

Hall symbol: P 2c -2n

$a = 18.4179$ (9) Å

$b = 5.7697$ (4) Å

$c = 12.4403$ (6) Å

$V = 1321.98$ (13) Å³

$Z = 4$

$F(000) = 576$

$D_x = 1.383$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2466 reflections

$\theta = 2.8$ – 26.7°

$\mu = 0.09$ mm⁻¹

$T = 173$ K

Plate, brown

$0.49 \times 0.15 \times 0.10$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ - and ω -scans

10655 measured reflections

1667 independent reflections

1258 reflections with $I > 2\sigma(I)$

$R_{int} = 0.081$

$\theta_{max} = 28.0^\circ$, $\theta_{min} = 2.2^\circ$

$h = -24 \rightarrow 24$

$k = -7 \rightarrow 7$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.088$
 $S = 0.95$
 1667 reflections
 190 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0468P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.12512 (12)	1.0166 (5)	0.33781 (19)	0.0300 (6)
C2	-0.13405 (14)	1.2022 (5)	0.40282 (19)	0.0348 (7)
H2	-0.1762	1.2958	0.3959	0.042*
C3	-0.08086 (14)	1.2588 (5)	0.4816 (2)	0.0369 (7)
H3	-0.0870	1.3914	0.5259	0.044*
C4	-0.02145 (13)	1.1223 (5)	0.4929 (2)	0.0373 (7)
H4	0.0131	1.1585	0.5471	0.045*
C5	-0.00952 (13)	0.9262 (5)	0.42571 (18)	0.0313 (6)
C6	0.05236 (13)	0.7881 (5)	0.43475 (19)	0.0340 (7)
H6	0.0869	0.8231	0.4891	0.041*
C7	0.06517 (13)	0.6010 (5)	0.3669 (2)	0.0331 (6)
C8	0.12738 (13)	0.4548 (5)	0.3779 (2)	0.0393 (7)
H8	0.1618	0.4860	0.4329	0.047*
C9	0.13797 (14)	0.2722 (6)	0.3111 (2)	0.0420 (7)
H9	0.1795	0.1764	0.3200	0.050*
C10	0.08711 (14)	0.2228 (6)	0.2274 (2)	0.0429 (7)
H10	0.0952	0.0961	0.1801	0.051*
C11	0.02709 (13)	0.3575 (5)	0.2156 (2)	0.0351 (7)
H11	-0.0065	0.3229	0.1599	0.042*
C12	0.01350 (12)	0.5483 (5)	0.28448 (18)	0.0300 (6)
C13	-0.04932 (12)	0.6848 (5)	0.27515 (19)	0.0298 (6)
H13	-0.0837	0.6496	0.2206	0.036*
C14	-0.06241 (12)	0.8710 (5)	0.34406 (18)	0.0280 (6)
C15	-0.19237 (13)	1.1049 (5)	0.16742 (19)	0.0324 (6)
C16	-0.24985 (14)	0.9892 (5)	0.1008 (2)	0.0409 (7)

H16A	-0.2302	0.9440	0.0298	0.049*
H16B	-0.2917	1.0943	0.0898	0.049*
C17	-0.27266 (14)	0.7749 (5)	0.1650 (2)	0.0394 (7)
H17A	-0.3250	0.7813	0.1828	0.047*
H17B	-0.2630	0.6316	0.1236	0.047*
C18	-0.22734 (12)	0.7812 (5)	0.2656 (2)	0.0328 (6)
N1	-0.17925 (10)	0.9652 (4)	0.25776 (16)	0.0301 (5)
O1	-0.16111 (10)	1.2847 (4)	0.14866 (14)	0.0409 (5)
O2	-0.23048 (9)	0.6519 (4)	0.34224 (15)	0.0415 (5)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0272 (12)	0.0376 (16)	0.0252 (11)	-0.0043 (11)	0.0015 (9)	0.0022 (13)
C2	0.0324 (13)	0.0423 (18)	0.0298 (13)	0.0020 (12)	0.0038 (10)	0.0024 (14)
C3	0.0414 (15)	0.0417 (18)	0.0278 (12)	-0.0020 (13)	0.0027 (11)	-0.0064 (13)
C4	0.0360 (14)	0.051 (2)	0.0253 (12)	-0.0075 (13)	0.0000 (11)	-0.0014 (14)
C5	0.0294 (12)	0.0421 (17)	0.0223 (11)	-0.0043 (12)	0.0000 (10)	0.0022 (12)
C6	0.0275 (12)	0.0484 (19)	0.0260 (12)	-0.0051 (12)	-0.0050 (10)	0.0036 (14)
C7	0.0280 (12)	0.0402 (18)	0.0312 (13)	-0.0038 (12)	0.0001 (10)	0.0069 (13)
C8	0.0262 (12)	0.050 (2)	0.0419 (14)	-0.0011 (13)	-0.0026 (11)	0.0111 (15)
C9	0.0290 (13)	0.0429 (19)	0.0541 (17)	0.0058 (13)	0.0047 (13)	0.0080 (16)
C10	0.0365 (15)	0.0420 (19)	0.0503 (17)	0.0007 (14)	0.0099 (13)	-0.0003 (17)
C11	0.0320 (14)	0.0402 (18)	0.0333 (13)	-0.0010 (13)	0.0013 (10)	-0.0016 (13)
C12	0.0260 (12)	0.0353 (17)	0.0287 (12)	-0.0020 (11)	0.0030 (9)	0.0051 (13)
C13	0.0258 (11)	0.0403 (18)	0.0235 (11)	-0.0061 (11)	0.0000 (9)	0.0032 (13)
C14	0.0261 (11)	0.0361 (16)	0.0217 (11)	-0.0054 (11)	0.0019 (9)	0.0022 (12)
C15	0.0279 (12)	0.0426 (17)	0.0267 (12)	0.0034 (12)	0.0039 (9)	0.0004 (13)
C16	0.0381 (13)	0.052 (2)	0.0326 (12)	-0.0054 (14)	-0.0045 (10)	0.0022 (14)
C17	0.0344 (13)	0.0445 (18)	0.0393 (14)	-0.0035 (12)	-0.0058 (11)	0.0011 (14)
C18	0.0260 (12)	0.0367 (16)	0.0357 (13)	0.0043 (11)	0.0009 (10)	0.0031 (14)
N1	0.0257 (9)	0.0377 (13)	0.0269 (9)	0.0010 (9)	-0.0007 (8)	0.0015 (10)
O1	0.0431 (10)	0.0482 (13)	0.0313 (9)	-0.0103 (10)	0.0025 (8)	0.0046 (9)
O2	0.0349 (9)	0.0453 (12)	0.0443 (11)	-0.0037 (9)	-0.0026 (8)	0.0134 (10)

Geometric parameters (Å, °)

C1—C2	1.352 (4)	C10—C11	1.359 (4)
C1—C14	1.430 (3)	C10—H10	0.9500
C1—N1	1.440 (3)	C11—C12	1.418 (4)
C2—C3	1.423 (4)	C11—H11	0.9500
C2—H2	0.9500	C12—C13	1.405 (3)
C3—C4	1.355 (4)	C13—C14	1.395 (3)
C3—H3	0.9500	C13—H13	0.9500
C4—C5	1.424 (4)	C15—O1	1.209 (3)
C4—H4	0.9500	C15—N1	1.404 (3)
C5—C6	1.395 (4)	C15—C16	1.501 (4)
C5—C14	1.443 (3)	C16—C17	1.531 (4)

C6—C7	1.390 (4)	C16—H16A	0.9900
C6—H6	0.9500	C16—H16B	0.9900
C7—C8	1.429 (4)	C17—C18	1.505 (3)
C7—C12	1.432 (3)	C17—H17A	0.9900
C8—C9	1.357 (4)	C17—H17B	0.9900
C8—H8	0.9500	C18—O2	1.211 (3)
C9—C10	1.430 (4)	C18—N1	1.386 (3)
C9—H9	0.9500		
C2—C1—C14	122.1 (2)	C12—C11—H11	119.3
C2—C1—N1	119.5 (2)	C13—C12—C11	122.1 (2)
C14—C1—N1	118.4 (2)	C13—C12—C7	119.2 (2)
C1—C2—C3	120.6 (3)	C11—C12—C7	118.7 (2)
C1—C2—H2	119.7	C14—C13—C12	121.6 (2)
C3—C2—H2	119.7	C14—C13—H13	119.2
C4—C3—C2	119.6 (3)	C12—C13—H13	119.2
C4—C3—H3	120.2	C13—C14—C1	123.9 (2)
C2—C3—H3	120.2	C13—C14—C5	119.1 (2)
C3—C4—C5	121.7 (2)	C1—C14—C5	117.0 (2)
C3—C4—H4	119.2	O1—C15—N1	124.4 (2)
C5—C4—H4	119.2	O1—C15—C16	127.7 (2)
C6—C5—C4	122.2 (2)	N1—C15—C16	107.9 (2)
C6—C5—C14	118.8 (2)	C15—C16—C17	105.4 (2)
C4—C5—C14	119.0 (2)	C15—C16—H16A	110.7
C7—C6—C5	122.2 (2)	C17—C16—H16A	110.7
C7—C6—H6	118.9	C15—C16—H16B	110.7
C5—C6—H6	118.9	C17—C16—H16B	110.7
C6—C7—C8	122.4 (2)	H16A—C16—H16B	108.8
C6—C7—C12	119.1 (2)	C18—C17—C16	105.2 (2)
C8—C7—C12	118.4 (3)	C18—C17—H17A	110.7
C9—C8—C7	121.0 (3)	C16—C17—H17A	110.7
C9—C8—H8	119.5	C18—C17—H17B	110.7
C7—C8—H8	119.5	C16—C17—H17B	110.7
C8—C9—C10	120.5 (3)	H17A—C17—H17B	108.8
C8—C9—H9	119.8	O2—C18—N1	123.9 (2)
C10—C9—H9	119.8	O2—C18—C17	127.8 (2)
C11—C10—C9	119.8 (3)	N1—C18—C17	108.3 (2)
C11—C10—H10	120.1	C18—N1—C15	112.7 (2)
C9—C10—H10	120.1	C18—N1—C1	123.5 (2)
C10—C11—C12	121.5 (3)	C15—N1—C1	123.7 (2)
C10—C11—H11	119.3		
C14—C1—C2—C3	-0.6 (4)	C2—C1—C14—C13	-176.7 (2)
N1—C1—C2—C3	-179.1 (2)	N1—C1—C14—C13	1.8 (3)
C1—C2—C3—C4	-1.4 (4)	C2—C1—C14—C5	1.9 (3)
C2—C3—C4—C5	1.9 (4)	N1—C1—C14—C5	-179.5 (2)
C3—C4—C5—C6	178.3 (3)	C6—C5—C14—C13	-1.4 (3)
C3—C4—C5—C14	-0.4 (4)	C4—C5—C14—C13	177.3 (2)

C4—C5—C6—C7	-177.9 (2)	C6—C5—C14—C1	179.8 (2)
C14—C5—C6—C7	0.8 (4)	C4—C5—C14—C1	-1.4 (3)
C5—C6—C7—C8	-178.1 (3)	O1—C15—C16—C17	-177.3 (2)
C5—C6—C7—C12	0.4 (4)	N1—C15—C16—C17	3.5 (3)
C6—C7—C8—C9	179.4 (3)	C15—C16—C17—C18	0.6 (3)
C12—C7—C8—C9	0.9 (4)	C16—C17—C18—O2	175.4 (3)
C7—C8—C9—C10	0.4 (4)	C16—C17—C18—N1	-4.5 (3)
C8—C9—C10—C11	-1.0 (4)	O2—C18—N1—C15	-172.7 (2)
C9—C10—C11—C12	0.3 (4)	C17—C18—N1—C15	7.2 (3)
C10—C11—C12—C13	-178.2 (3)	O2—C18—N1—C1	3.1 (4)
C10—C11—C12—C7	1.1 (4)	C17—C18—N1—C1	-177.0 (2)
C6—C7—C12—C13	-0.9 (3)	O1—C15—N1—C18	173.9 (2)
C8—C7—C12—C13	177.6 (2)	C16—C15—N1—C18	-6.9 (3)
C6—C7—C12—C11	179.9 (2)	O1—C15—N1—C1	-1.8 (4)
C8—C7—C12—C11	-1.6 (4)	C16—C15—N1—C1	177.4 (2)
C11—C12—C13—C14	179.4 (2)	C2—C1—N1—C18	-107.2 (3)
C7—C12—C13—C14	0.2 (3)	C14—C1—N1—C18	74.2 (3)
C12—C13—C14—C1	179.6 (2)	C2—C1—N1—C15	68.1 (3)
C12—C13—C14—C5	0.9 (3)	C14—C1—N1—C15	-110.5 (3)

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C7—C12 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C2—H2 \cdots O2 ⁱ	0.95	2.38	3.234 (3)	149
C6—H6 \cdots O1 ⁱⁱ	0.95	2.49	3.357 (3)	152
C9—H9 \cdots O2 ⁱⁱⁱ	0.95	2.53	3.465 (3)	167
C13—H13 \cdots O1 ^{iv}	0.95	2.70	3.471 (3)	139
C17—H17A \cdots Cg2 ^v	0.99	2.92	3.709 (3)	138

Symmetry codes: (i) $x, y+1, z$; (ii) $-x, -y+2, z+1/2$; (iii) $x+1/2, -y+1/2, z$; (iv) $x, y-1, z$; (v) $x+1/2, -y+3/2, z$.