

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(dicyanamido- κ N)tetrakis(pyridine- κ N)nickel(II)Susanne Wöhlert,^{a*} Mario Wriedt,^b Inke Jess^a and Christian Näther^a^aInstitut für Anorganische Chemie, Christian-Albrechts-Universität Kiel, Max-Eyth-Strasse 2, 24118 Kiel, Germany, and ^bDepartment of Chemistry, Texas A&M University, College Station, Texas 77843, USA

Correspondence e-mail: swoehlert@ac.uni-kiel.de


Received 20 April 2012; accepted 2 May 2012

Key indicators: single-crystal X-ray study; $T = 170$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.036; wR factor = 0.092; data-to-parameter ratio = 16.9.

In the crystal structure of the title compound, $[\text{Ni}(\text{C}_2\text{N}_3)_2(\text{C}_5\text{H}_5\text{N})_4]$, the Ni^{II} cations are coordinated by four pyridine ligands and two dicyanamide anions into discrete complexes. The shortest $\text{Ni}\cdots\text{Ni}$ separation is 8.1068 (10) Å. The structure is pseudo-centrosymmetric and can also be refined in the space group $C2/c$ in which both anionic ligands are strongly disordered and the refinement leads to significantly poorer reliability factors.

Related literature

For related structures, see: Boeckmann & Näther (2010, 2011); Wriedt & Näther (2011); Wu *et al.* (2004). For a description of the Cambridge Structural Database, see: Allen (2002).



Experimental

Crystal data

 $[\text{Ni}(\text{C}_2\text{N}_3)_2(\text{C}_5\text{H}_5\text{N})_4]$
 $M_r = 507.21$

 Monoclinic, Cc
 $a = 13.0439$ (6) Å
 $b = 12.8557$ (8) Å
 $c = 15.1294$ (7) Å
 $\beta = 110.191$ (5)°

 $V = 2381.1$ (2) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.85$ mm⁻¹
 $T = 170$ K
 $0.05 \times 0.04 \times 0.04$ mm

Data collection

 Stoe IPDS-1 diffractometer
 Absorption correction: numerical
 (X -SHAPE and X -RED32;
 Stoe & Cie, 2008)
 $T_{\text{min}} = 0.911$, $T_{\text{max}} = 0.973$

 11154 measured reflections
 5386 independent reflections
 4554 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.092$
 $S = 0.98$
 5386 reflections
 318 parameters

 2 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.44$ e Å⁻³

Data collection: X -AREA (Stoe & Cie, 2008); cell refinement: X -AREA; data reduction: X -AREA; program(s) used to solve structure: $SHELXS97$ (Sheldrick, 2008); program(s) used to refine structure: $SHELXL97$ (Sheldrick, 2008); molecular graphics: XP in $SHELXTL$ (Sheldrick, 2008) and $DIAMOND$ (Brandenburg, 2011); software used to prepare material for publication: $XCIF$ in $SHELXTL$.

We gratefully acknowledge financial support by the DFG (project No. NA 720/3-1) and the State of Schleswig-Holstein. We thank Professor Dr Wolfgang Bensch for access to his experimental facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5887).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Boeckmann, J. & Näther, C. (2010). *Dalton Trans.* **39**, 11019–11026.
 Boeckmann, J. & Näther, C. (2011). *Chem. Commun.* **47**, 7104–7106.
 Brandenburg, K. (2011). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stoe & Cie (2008). X -AREA, X -RED32 and X -SHAPE. Stoe & Cie, Darmstadt, Germany.
 Wriedt, M. & Näther, C. (2011). *Dalton Trans.* **40**, 886–898.
 Wu, A.-Q., Zheng, F.-K., Guo, G.-C. & Huang, J.-S. (2004). *Acta Cryst.* **E60**, m373–m375.

supporting information

Acta Cryst. (2012). E68, m745 [doi:10.1107/S1600536812019691]

Bis(dicyanamido- κ N)tetrakis(pyridine- κ N)nickel(II)

Susanne Wöhlert, Mario Wriedt, Inke Jess and Christian Näther

S1. Comment

Recently we have reported on the thermal and magnetic properties of coordination polymers based on paramagnetic transition metal thio- and selenocyanato compounds in which the cations are linked by thio- or selenocyanato anions (Boeckmann & Näther; 2010; Boeckmann & Näther, 2011). The bridging compounds with *e.g.* pyridine can be prepared by thermal decomposition of discrete complexes with terminal bonded anionic ligands. In the course of systematic investigations we also started to investigate similar compounds based on dicyanamide (Wriedt & Näther, 2011). Within this project crystals of the title compound were obtained, which were characterized by single-crystal X-ray diffraction. In the crystal structure of the title compound each nickel(II) cation is six-coordinated by two dicyanamido anions and four pyridine ligands in a slightly distorted octahedral geometry (Fig. 1). The NiN₆ distances range from 2.057 (3) Å to 2.169 (3) Å and the angles are between 86.59 (11) ° and 178.96 (15)°. In the crystal structure, the discrete complexes are connected through intermolecular N—HC hydrogen bonds with an N···H distances of 2.559 Å (Fig. 2). The shortest nickel(II)—nickel(II) distance between the complexes is 9.157 Å.

It must be noted that a discrete nickel(II) dicyanamide complex is reported (Wu *et al.*, 2004) with 1,10-phenanthroline as co-ligands in which all ligands are *cis*-coordinated.

S2. Experimental

Nickel(II) chloride hexahydrate (NiCl₂·6H₂O), sodium dicyanamide (NaN(CN)₂) and pyridine were obtained from Alfa Aesar. All chemicals were used without further purification. 0.125 mmol (29.7 mg) NiCl₂·6H₂O, 0.25 mmol (22.3 mg) NaN(CN)₂ were reacted in 2.5 ml pyridine. Light-green single-crystal of the title compound were obtained after three days.

S3. Refinement

All H atoms were located in difference map but were positioned with idealized geometry and were refined isotropic with $U_{\text{eq}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ of the parent atom using a riding model with C—H = 0.95 Å. The structure is pseudo-centrosymmetric and can also be refined in the centrosymmetric space group *C2/c*. However, in *C2/c* the complexes are located on centres of inversion and the anionic ligands are strongly disordered which is not the case in space group *Cc*. Moreover, in *C2/c* the reliability factors are very poor and no reasonable structure model can be found. Therefore, *Cc* was selected, in which the structure can be very easily refined. In this case the absolute structure cannot be determined presumably, because of the pseudo-symmetry and therefore, a twin refinement for racemic twinning was performed leading to an BASF parameter of 0.53 (2).




Figure 1

The molecular structure of the title compound with labelling and displacement ellipsoids drawn at the 50% probability level.




Figure 2

Crystal structure of the title compound with a view of the discrete complexes (green = nickel(II), blue = nitrogen, grey = carbon and white = hydrogen).

Bis(dicyanamido- κ N)tetrakis(pyridine- κ N)nickel(II)

Crystal data

[Ni(C₂N₃)₂(C₅H₅N)₄]

$M_r = 507.21$

Monoclinic, *Cc*

Hall symbol: C -2yc

$a = 13.0439$ (6) Å

$b = 12.8557$ (8) Å

$c = 15.1294$ (7) Å

$\beta = 110.191$ (5)°

$V = 2381.1$ (2) Å³

$Z = 4$

$F(000) = 1048$

$D_x = 1.415$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 11154 reflections

$\theta = 3.0$ – 28.1 °

$\mu = 0.85$ mm⁻¹

$T = 170$ K

Block, light green

$0.05 \times 0.04 \times 0.04$ mm

Data collection

Stoe IPDS-1
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi scan

Absorption correction: numerical

(*X-SHAPE* and *X-RED32*; Stoe & Cie, 2008)

$T_{\min} = 0.911$, $T_{\max} = 0.973$

11154 measured reflections

5386 independent reflections

4554 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\max} = 28.1$ °, $\theta_{\min} = 3.0$ °

$h = -17 \rightarrow 17$

$k = -16 \rightarrow 16$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.092$

$S = 0.98$

5386 reflections

318 parameters

2 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0593P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.35 \text{ e } \text{Å}^{-3}$

$$\Delta\rho_{\min} = -0.44 \text{ e } \text{Å}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0106 (6)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.53427 (5)	0.75512 (3)	0.65840 (5)	0.01803 (10)
N1	0.5772 (2)	0.68096 (18)	0.7947 (2)	0.0204 (6)
C1	0.5414 (3)	0.7163 (3)	0.8622 (3)	0.0254 (7)
H1	0.4997	0.7786	0.8508	0.031*
C2	0.5620 (3)	0.6669 (3)	0.9474 (3)	0.0314 (8)
H2	0.5360	0.6956	0.9935	0.038*
C3	0.6207 (3)	0.5749 (3)	0.9651 (3)	0.0332 (8)
H3	0.6345	0.5385	1.0227	0.040*
C4	0.6587 (3)	0.5376 (3)	0.8962 (3)	0.0338 (8)
H4	0.6992	0.4747	0.9057	0.041*
C5	0.6369 (3)	0.5932 (2)	0.8131 (3)	0.0280 (7)
H5	0.6655	0.5682	0.7672	0.034*
N11	0.6453 (2)	0.88091 (19)	0.7206 (2)	0.0230 (6)
C11	0.7525 (3)	0.8647 (3)	0.7619 (3)	0.0364 (9)
H11	0.7792	0.7954	0.7664	0.044*
C12	0.8266 (4)	0.9446 (3)	0.7983 (4)	0.0512 (12)
H12	0.9024	0.9299	0.8262	0.061*
C13	0.7897 (4)	1.0450 (3)	0.7938 (4)	0.0480 (11)
H13	0.8391	1.1008	0.8188	0.058*
C14	0.6809 (4)	1.0628 (3)	0.7529 (3)	0.0416 (10)
H14	0.6527	1.1315	0.7484	0.050*
C15	0.6115 (3)	0.9793 (2)	0.7179 (3)	0.0331 (8)
H15	0.5355	0.9929	0.6903	0.040*
N21	0.4158 (2)	0.63412 (19)	0.5948 (2)	0.0223 (6)
C21	0.4425 (3)	0.5333 (2)	0.6051 (3)	0.0285 (8)
H21	0.5166	0.5153	0.6378	0.034*
C22	0.3677 (4)	0.4536 (3)	0.5706 (3)	0.0403 (10)
H22	0.3902	0.3829	0.5799	0.048*
C23	0.2604 (4)	0.4788 (3)	0.5226 (3)	0.0419 (10)
H23	0.2071	0.4259	0.4988	0.050*
C24	0.2317 (3)	0.5822 (3)	0.5098 (3)	0.0406 (9)

H24	0.1585	0.6020	0.4756	0.049*
C25	0.3110 (3)	0.6567 (3)	0.5475 (3)	0.0297 (8)
H25	0.2901	0.7279	0.5393	0.036*
N31	0.4925 (2)	0.81948 (19)	0.5191 (2)	0.0228 (6)
C31	0.5106 (3)	0.7632 (3)	0.4508 (3)	0.0269 (7)
H31	0.5396	0.6950	0.4657	0.032*
C32	0.4890 (3)	0.7998 (3)	0.3597 (3)	0.0359 (8)
H32	0.5032	0.7575	0.3138	0.043*
C33	0.4467 (3)	0.8983 (3)	0.3369 (3)	0.0368 (9)
H33	0.4307	0.9249	0.2750	0.044*
C34	0.4282 (4)	0.9574 (3)	0.4057 (3)	0.0384 (9)
H34	0.4000	1.0260	0.3924	0.046*
C35	0.4513 (3)	0.9153 (3)	0.4948 (3)	0.0326 (8)
H35	0.4371	0.9565	0.5414	0.039*
N41	0.6587 (2)	0.67389 (19)	0.6335 (2)	0.0241 (6)
C41	0.7342 (3)	0.6558 (2)	0.6128 (2)	0.0234 (7)
N42	0.8215 (3)	0.6277 (3)	0.5954 (3)	0.0398 (8)
C42	0.8367 (3)	0.6685 (3)	0.5206 (3)	0.0445 (10)
N43	0.8561 (4)	0.6974 (3)	0.4559 (3)	0.0723 (14)
N51	0.4106 (2)	0.8376 (2)	0.6812 (2)	0.0248 (6)
C51	0.3343 (3)	0.8681 (2)	0.6955 (2)	0.0198 (6)
N52	0.2530 (3)	0.9113 (2)	0.7117 (2)	0.0304 (6)
C52	0.1669 (3)	0.8567 (2)	0.7082 (2)	0.0266 (6)
N53	0.0854 (3)	0.8178 (3)	0.7044 (3)	0.0512 (9)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.01894 (14)	0.01636 (15)	0.02019 (15)	0.00311 (14)	0.00853 (10)	-0.00008 (15)
N1	0.0215 (15)	0.0189 (11)	0.0215 (17)	0.0028 (9)	0.0083 (13)	0.0001 (10)
C1	0.0266 (16)	0.0255 (15)	0.0252 (19)	0.0042 (13)	0.0103 (15)	0.0020 (14)
C2	0.0326 (18)	0.0417 (18)	0.0224 (18)	0.0036 (14)	0.0127 (15)	-0.0012 (14)
C3	0.037 (2)	0.0382 (18)	0.024 (2)	0.0060 (14)	0.0090 (17)	0.0072 (14)
C4	0.043 (2)	0.0283 (15)	0.030 (2)	0.0167 (14)	0.0136 (17)	0.0102 (13)
C5	0.0342 (18)	0.0260 (14)	0.0257 (18)	0.0106 (12)	0.0127 (15)	0.0035 (12)
N11	0.0242 (15)	0.0221 (12)	0.0234 (16)	0.0002 (10)	0.0093 (13)	-0.0014 (10)
C11	0.033 (2)	0.0281 (16)	0.044 (3)	-0.0006 (13)	0.0074 (19)	-0.0027 (14)
C12	0.033 (2)	0.044 (2)	0.061 (3)	-0.0111 (17)	-0.003 (2)	-0.008 (2)
C13	0.051 (3)	0.0321 (18)	0.060 (3)	-0.0195 (17)	0.019 (2)	-0.0114 (18)
C14	0.048 (2)	0.0229 (15)	0.055 (3)	-0.0078 (15)	0.018 (2)	-0.0056 (15)
C15	0.035 (2)	0.0189 (14)	0.047 (2)	-0.0006 (13)	0.0160 (18)	-0.0047 (14)
N21	0.0225 (15)	0.0185 (11)	0.0259 (16)	-0.0004 (9)	0.0084 (13)	-0.0028 (10)
C21	0.0280 (17)	0.0188 (13)	0.038 (2)	0.0029 (12)	0.0106 (16)	-0.0011 (12)
C22	0.048 (2)	0.0187 (15)	0.052 (3)	-0.0014 (14)	0.014 (2)	-0.0050 (14)
C23	0.035 (2)	0.0349 (18)	0.050 (3)	-0.0111 (15)	0.0068 (19)	-0.0073 (16)
C24	0.0274 (19)	0.0347 (18)	0.050 (3)	-0.0043 (14)	0.0009 (18)	-0.0037 (16)
C25	0.0222 (16)	0.0257 (14)	0.034 (2)	0.0013 (12)	0.0006 (15)	-0.0023 (13)
N31	0.0239 (15)	0.0220 (12)	0.0221 (17)	0.0011 (10)	0.0075 (13)	0.0024 (10)

C31	0.0275 (16)	0.0328 (17)	0.0208 (18)	0.0002 (13)	0.0089 (14)	-0.0039 (13)
C32	0.0334 (19)	0.053 (2)	0.023 (2)	-0.0027 (16)	0.0118 (16)	-0.0039 (15)
C33	0.033 (2)	0.051 (2)	0.026 (2)	-0.0024 (15)	0.0106 (17)	0.0107 (15)
C34	0.043 (2)	0.0395 (19)	0.030 (2)	0.0065 (15)	0.0077 (18)	0.0153 (15)
C35	0.040 (2)	0.0297 (15)	0.029 (2)	0.0076 (13)	0.0123 (17)	0.0056 (12)
N41	0.0250 (15)	0.0243 (12)	0.0258 (17)	0.0077 (10)	0.0122 (14)	0.0019 (10)
C41	0.0243 (16)	0.0230 (14)	0.0232 (18)	-0.0021 (11)	0.0085 (14)	-0.0037 (11)
N42	0.0305 (17)	0.0533 (19)	0.045 (2)	0.0087 (14)	0.0245 (15)	-0.0011 (14)
C42	0.048 (2)	0.0416 (18)	0.060 (3)	-0.0219 (16)	0.039 (2)	-0.0256 (17)
N43	0.103 (3)	0.067 (3)	0.082 (3)	-0.046 (2)	0.076 (3)	-0.033 (2)
N51	0.0223 (15)	0.0225 (12)	0.0307 (18)	0.0055 (10)	0.0106 (14)	-0.0014 (10)
C51	0.0244 (15)	0.0164 (12)	0.0168 (15)	0.0023 (10)	0.0048 (12)	-0.0002 (10)
N52	0.0295 (14)	0.0243 (13)	0.0441 (18)	0.0036 (11)	0.0213 (13)	-0.0051 (11)
C52	0.0266 (15)	0.0241 (13)	0.0324 (18)	0.0097 (11)	0.0144 (13)	0.0082 (11)
N53	0.0377 (17)	0.0438 (17)	0.079 (3)	0.0064 (14)	0.0294 (18)	0.0222 (17)

Geometric parameters (Å, °)

Ni1—N51	2.057 (3)	N21—C25	1.338 (5)
Ni1—N41	2.071 (3)	C21—C22	1.386 (5)
Ni1—N31	2.152 (3)	C21—H21	0.9500
Ni1—N11	2.158 (3)	C22—C23	1.375 (6)
Ni1—N1	2.162 (3)	C22—H22	0.9500
Ni1—N21	2.169 (3)	C23—C24	1.377 (6)
N1—C1	1.341 (4)	C23—H23	0.9500
N1—C5	1.344 (4)	C24—C25	1.381 (5)
C1—C2	1.377 (5)	C24—H24	0.9500
C1—H1	0.9500	C25—H25	0.9500
C2—C3	1.383 (5)	N31—C35	1.343 (4)
C2—H2	0.9500	N31—C31	1.348 (5)
C3—C4	1.385 (5)	C31—C32	1.391 (5)
C3—H3	0.9500	C31—H31	0.9500
C4—C5	1.388 (5)	C32—C33	1.376 (6)
C4—H4	0.9500	C32—H32	0.9500
C5—H5	0.9500	C33—C34	1.376 (6)
N11—C15	1.336 (4)	C33—H33	0.9500
N11—C11	1.337 (5)	C34—C35	1.385 (5)
C11—C12	1.388 (6)	C34—H34	0.9500
C11—H11	0.9500	C35—H35	0.9500
C12—C13	1.371 (6)	N41—C41	1.154 (4)
C12—H12	0.9500	C41—N42	1.306 (4)
C13—C14	1.358 (7)	N42—C42	1.322 (5)
C13—H13	0.9500	C42—N43	1.154 (5)
C14—C15	1.386 (5)	N51—C51	1.157 (4)
C14—H14	0.9500	C51—N52	1.294 (4)
C15—H15	0.9500	N52—C52	1.309 (4)
N21—C21	1.337 (4)	C52—N53	1.159 (4)

N51—Ni1—N41	178.96 (15)	N11—C15—C14	123.8 (4)
N51—Ni1—N31	91.03 (12)	N11—C15—H15	118.1
N41—Ni1—N31	88.00 (11)	C14—C15—H15	118.1
N51—Ni1—N11	89.29 (11)	C21—N21—C25	116.7 (3)
N41—Ni1—N11	90.40 (11)	C21—N21—Ni1	121.8 (3)
N31—Ni1—N11	92.51 (11)	C25—N21—Ni1	121.4 (2)
N51—Ni1—N1	91.56 (11)	N21—C21—C22	123.5 (4)
N41—Ni1—N1	89.43 (11)	N21—C21—H21	118.3
N31—Ni1—N1	176.39 (11)	C22—C21—H21	118.3
N11—Ni1—N1	90.04 (11)	C23—C22—C21	118.7 (3)
N51—Ni1—N21	87.97 (11)	C23—C22—H22	120.6
N41—Ni1—N21	92.33 (11)	C21—C22—H22	120.6
N31—Ni1—N21	86.59 (11)	C22—C23—C24	118.6 (4)
N11—Ni1—N21	177.09 (12)	C22—C23—H23	120.7
N1—Ni1—N21	90.98 (11)	C24—C23—H23	120.7
C1—N1—C5	117.1 (3)	C23—C24—C25	118.9 (4)
C1—N1—Ni1	122.3 (2)	C23—C24—H24	120.5
C5—N1—Ni1	120.5 (2)	C25—C24—H24	120.5
N1—C1—C2	123.3 (3)	N21—C25—C24	123.5 (3)
N1—C1—H1	118.3	N21—C25—H25	118.2
C2—C1—H1	118.3	C24—C25—H25	118.2
C1—C2—C3	119.5 (3)	C35—N31—C31	116.2 (3)
C1—C2—H2	120.3	C35—N31—Ni1	124.3 (2)
C3—C2—H2	120.3	C31—N31—Ni1	119.5 (2)
C2—C3—C4	117.9 (3)	N31—C31—C32	123.3 (4)
C2—C3—H3	121.1	N31—C31—H31	118.3
C4—C3—H3	121.1	C32—C31—H31	118.3
C3—C4—C5	119.3 (3)	C33—C32—C31	119.1 (4)
C3—C4—H4	120.3	C33—C32—H32	120.5
C5—C4—H4	120.3	C31—C32—H32	120.5
N1—C5—C4	122.8 (3)	C32—C33—C34	118.6 (4)
N1—C5—H5	118.6	C32—C33—H33	120.7
C4—C5—H5	118.6	C34—C33—H33	120.7
C15—N11—C11	116.3 (3)	C33—C34—C35	118.9 (4)
C15—N11—Ni1	122.1 (3)	C33—C34—H34	120.5
C11—N11—Ni1	121.6 (2)	C35—C34—H34	120.5
N11—C11—C12	122.9 (4)	N31—C35—C34	123.9 (4)
N11—C11—H11	118.5	N31—C35—H35	118.1
C12—C11—H11	118.5	C34—C35—H35	118.1
C13—C12—C11	119.5 (5)	C41—N41—Ni1	161.0 (3)
C13—C12—H12	120.2	N41—C41—N42	174.3 (4)
C11—C12—H12	120.2	C41—N42—C42	117.5 (4)
C14—C13—C12	118.4 (4)	N43—C42—N42	174.2 (5)
C14—C13—H13	120.8	C51—N51—Ni1	168.7 (3)
C12—C13—H13	120.8	N51—C51—N52	174.3 (3)
C13—C14—C15	119.1 (4)	C51—N52—C52	120.7 (3)
C13—C14—H14	120.5	N53—C52—N52	173.2 (3)
C15—C14—H14	120.5		
