

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Ethyl-6-(4-fluorophenyl)-7H-1,2,4-triazolo[3,4-b][1,3,4]thiadiazine

S. Jeyaseelan,^a H. C. Devarajgowda,^{a*} R. Sathishkumar,^b Agnes Sylvia D'souza^c and Alphonsus D'souza^c

^aDepartment of Physics, Yuvaraja's College (Constituent College), University of Mysore, Mysore 570 005, Karnataka, India, ^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 12, Karnataka, India, and ^cDepartment of Physics, St Philomena's College (Autonomous), Mysore 570 015, Karnataka, India
Correspondence e-mail: devarajgowda@yahoo.com


Received 12 May 2012; accepted 14 May 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.106; data-to-parameter ratio = 12.9.

In the title compound, $\text{C}_{12}\text{H}_{11}\text{FN}_4\text{S}$, the thiadiazine ring adopts a twist-boat conformation. The dihedral angle between the triazolothiadiazine system and the benzene ring is $10.54(9)^\circ$. The crystal structure is characterized by $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds. The crystal packing also exhibits $\pi-\pi$ interactions, with a centroid-centroid distance of $3.6348(15)$ Å.

Related literature

For biological properties of triazolothiadiazines, see: Feng *et al.* (1992); Mohan & Anjaneyalu (1987); Holla *et al.* (2001); Walser *et al.* (1991); Hirota *et al.* (1991); Bradbury & Rivett (1991); Heindel & Reid (1980); Heidelberger *et al.* (1957). For related structures, see: Andersson & MacGowan (2003); Novak *et al.* (2006).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{11}\text{FN}_4\text{S}$
 $M_r = 262.31$
Monoclinic, $P2_1/c$
 $a = 13.322(3)$ Å
 $b = 13.017(3)$ Å
 $c = 7.1912(16)$ Å
 $\beta = 105.308(4)^\circ$

$V = 1202.8(4)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.27$ mm⁻¹
 $T = 293$ K
 $0.24 \times 0.20 \times 0.12$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\min} = 0.770$, $T_{\max} = 1.000$
11097 measured reflections
2119 independent reflections
1828 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.106$
 $S = 1.06$
2119 reflections
164 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|--|-------|-------------|-------------|---------------|
| $\text{C1}-\text{H1}\cdots\text{N2}^i$ | 0.93 | 2.51 | 3.428 (3) | 172 |
| $\text{C8}-\text{H8A}\cdots\text{N1}^{ii}$ | 0.97 | 2.50 | 3.410 (3) | 156 |
| $\text{C8}-\text{H8B}\cdots\text{N2}^i$ | 0.97 | 2.30 | 3.228 (3) | 160 |

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97.

The authors thank Professor T. N. Guru Row, Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, for the data collection. SJ thanks the University Grants Commission (India), for the award of a Teacher Fellowship under the Faculty Development Programme (UGC-SWRO File No.: FIP/11 th Plan/KAMY006 TF, Dt.: 06/08/2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5922).

References

- Andersson, M. I. & MacGowan, A. P. (2003). *J. Antimicrob. Chemother.* **51**, 1–11.
Bradbury, R. H. & Rivett, J. E. (1991). *J. Med. Chem.* **34**, 151–157.
Bruker (2001). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Feng, X. M., Chen, R. & Yang, W. D. (1992). *Chem. J. Chin. Univ.* **13**, 187–194.
Heidelberger, C., Chaudhuri, N. K., Danneberg, P., Mooren, D., Greisbach, L., Duschinsky, R., Sennitzer, R. J., Pleaven, E. & Scheiner, J. (1957). *Nature (London)*, **179**, 663–666.
Heindel, N. D. & Reid, J. R. (1980). *J. Heterocycl. Chem.* **17**, 1087–1088.
Hirota, T., Sasaki, K., Yamamoto, H. & Nakayama, T. (1991). *J. Heterocycl. Chem.* **28**, 257–261.
Holla, B. S., Akberali, P. M. & Shivananda, M. K. (2001). *Il Farmaco*, **56**, 919–927.
Mohan, J. & Anjaneyalu, G. S. R. (1987). *Pol. J. Chem.* **61**, 547–551.
Novak, M. J., Baum, J. C. & Buhrow, J. A. (2006). *Olson. Surf. Sci.* **600**, L269–L273.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Walser, A., Flynn, T. & Mason, C. (1991). *J. Heterocycl. Chem.* **28**, 1121–1125.

supporting information

Acta Cryst. (2012). E68, o1860 [doi:10.1107/S160053681202185X]

3-Ethyl-6-(4-fluorophenyl)-7H-1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazine

S. Jeyaseelan, H. C. Devarajegowda, R. Sathishkumar, Agnes Sylvia D'souza and Alphonsus D'souza

S1. Comment

Triazoles fused with six membered ring systems are found to be associated with diverse pharmacological activity. A number of thiadiazines have been shown to exhibit antimicrobial (Feng *et al.*, 1992) and diuretic properties (Mohan *et al.*, 1987) and also serves as photographic couplers (Holla *et al.*, 2001). The analgesic, anti-asthmatic, diuretic, anti-hypertensive, anti-cholinergic, antibacterial, antifungal, anti-inflammatory, hypoglycemic, anti-tubercular and antiviral properties exhibited by various N-bridged heterocycles derived from a variety of 4-amino-5-mercapto-1,2,4-triazoles, have made them an important chemotherapeutic agents (Walser *et al.*, 1991; Hirota *et al.*, 1991; Bradbury *et al.*, 1991). The 1,2,4- triazoles nucleus has recently been incorporated into a wide variety of therapeutically interesting drugs including H1/H2 histamine receptor blockers, cholinesterase active agents, CNS stimulants, anti-anxiety agents and sedatives (Heindel *et al.*, 1980). Further fluorinated heterocycles have been shown to possess wide variety of biocidal activities. Compounds such as fluorouracil and fluoroquinolone have been used as anticancer agents and antibiotics (Heidelberger *et al.*, 1957; Andersson *et al.*, 2003; Novak *et al.*, 2006).

The asymmetric unit of 3-ethyl-6-(4-fluorophenyl)-7H-[1,2,4]triazolo[3,4-*b*] [1,3,4]thiadiazine is shown in Fig. 1. The triazolo-thiadiazine ring system is not planar. The dihedral angle between the triazolo-thiadiazine ring system (S1/N1–N4/C7–C10) and the benzene ring (C1–C6) is 10.54 (9)°.

In the crystal structure (Fig. 2), the molecules are connected *via* intermolecular C1—H1⋯N2, C8—H8A⋯N1 and C8—H8B⋯N2 hydrogen bonds (Table 1). Furthermore, the crystal structure features a π - π interaction, with a centroid-centroid $Cg1$ (C9/C10/N1–N3) distance of 3.5728 (16) Å.

S2. Experimental

A mixture of triazole (1) (0.01 mol) and *p*-fluorophenacyl bromide (0.01 mol) in ethanol (25 ml) was heated under reflux for 1–2 hrs. The reaction mixture was cooled to room temperature and neutralized with sodium acetate (5%). The precipitated triazolothiadiazines were collected by filtration, washed with water and recrystallized from ethanol. Yield 82%; m.p.455 K.

S3. Refinement

All H atoms were positioned geometrically, with C—H = 0.93 Å for aromatic H, C—H = 0.97 Å for methylene H and C—H = 0.96 Å for methyl H, and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for all other H.




Figure 1

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

The packing of molecules in the title structure.

3-Ethyl-6-(4-fluorophenyl)-7H-1,2,4- triazolo[3,4-b][1,3,4]thiadiazine

Crystal data

$C_{12}H_{11}FN_4S$

$M_r = 262.31$

Monoclinic, $P2_1/c$

Hall symbol: $-P 2_1/c$

$a = 13.322 (3) \text{ \AA}$

$b = 13.017 (3) \text{ \AA}$

$c = 7.1912 (16) \text{ \AA}$

$\beta = 105.308 (4)^\circ$

$V = 1202.8 (4) \text{ \AA}^3$

$Z = 4$

$F(000) = 544$

$D_x = 1.449 \text{ Mg m}^{-3}$

Melting point: 455 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2119 reflections
 $\theta = 2.2\text{--}25.0^\circ$
 $\mu = 0.27\text{ mm}^{-1}$

$T = 293\text{ K}$
 Plate, colourless
 $0.24 \times 0.20 \times 0.12\text{ mm}$

Data collection

Bruker SMART CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω and φ scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2001)
 $T_{\min} = 0.770$, $T_{\max} = 1.000$

11097 measured reflections
 2119 independent reflections
 1828 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -15 \rightarrow 15$
 $k = -15 \rightarrow 15$
 $l = -8 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.106$
 $S = 1.06$
 2119 reflections
 164 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0487P)^2 + 0.4632P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.22\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.22\text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|----|--------------|--------------|-------------|----------------------------------|
| S1 | 0.50854 (4) | 0.90687 (4) | 0.19250 (9) | 0.0542 (2) |
| F1 | 0.05432 (12) | 1.31988 (11) | -0.1252 (2) | 0.0845 (5) |
| N1 | 0.37247 (14) | 0.64191 (13) | 0.1417 (3) | 0.0511 (5) |
| N2 | 0.46299 (14) | 0.70215 (13) | 0.1828 (3) | 0.0550 (5) |
| N3 | 0.32718 (12) | 0.80332 (11) | 0.1211 (2) | 0.0397 (4) |
| N4 | 0.26399 (12) | 0.88834 (12) | 0.0605 (2) | 0.0403 (4) |
| C1 | 0.28158 (17) | 1.16575 (16) | 0.0713 (3) | 0.0534 (6) |
| H1 | 0.3526 | 1.1746 | 0.1256 | 0.064* |
| C2 | 0.21920 (19) | 1.25088 (18) | 0.0115 (4) | 0.0626 (6) |
| H2 | 0.2476 | 1.3166 | 0.0253 | 0.075* |
| C3 | 0.11625 (18) | 1.23664 (17) | -0.0673 (4) | 0.0566 (6) |
| C4 | 0.07182 (18) | 1.14193 (19) | -0.0911 (4) | 0.0625 (6) |
| H4 | 0.0008 | 1.1346 | -0.1464 | 0.075* |

| | | | | |
|------|--------------|--------------|-------------|------------|
| C5 | 0.13374 (16) | 1.05684 (17) | -0.0319 (3) | 0.0536 (6) |
| H5 | 0.1042 | 0.9916 | -0.0479 | 0.064* |
| C6 | 0.24016 (15) | 1.06767 (15) | 0.0515 (3) | 0.0407 (5) |
| C7 | 0.30545 (14) | 0.97569 (14) | 0.1164 (3) | 0.0378 (4) |
| C8 | 0.41148 (15) | 0.98739 (15) | 0.2543 (3) | 0.0446 (5) |
| H8A | 0.4073 | 0.9702 | 0.3833 | 0.054* |
| H8B | 0.4330 | 1.0586 | 0.2552 | 0.054* |
| C9 | 0.43346 (15) | 0.79735 (15) | 0.1685 (3) | 0.0436 (5) |
| C10 | 0.29301 (15) | 0.70301 (14) | 0.1038 (3) | 0.0423 (5) |
| C11 | 0.18178 (17) | 0.67302 (16) | 0.0591 (4) | 0.0547 (6) |
| H11A | 0.1431 | 0.7122 | -0.0514 | 0.066* |
| H11B | 0.1548 | 0.6907 | 0.1677 | 0.066* |
| C12 | 0.1639 (2) | 0.56042 (18) | 0.0163 (4) | 0.0671 (7) |
| H12A | 0.1819 | 0.5441 | -0.1011 | 0.101* |
| H12B | 0.0920 | 0.5443 | 0.0024 | 0.101* |
| H12C | 0.2066 | 0.5210 | 0.1204 | 0.101* |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|-------------|--------------|
| S1 | 0.0360 (3) | 0.0450 (3) | 0.0800 (4) | -0.0006 (2) | 0.0126 (3) | 0.0070 (3) |
| F1 | 0.0731 (10) | 0.0583 (9) | 0.1191 (13) | 0.0317 (8) | 0.0204 (9) | 0.0218 (8) |
| N1 | 0.0518 (11) | 0.0354 (9) | 0.0640 (12) | 0.0038 (8) | 0.0113 (9) | -0.0002 (8) |
| N2 | 0.0464 (10) | 0.0411 (10) | 0.0745 (13) | 0.0081 (8) | 0.0108 (9) | 0.0033 (9) |
| N3 | 0.0371 (8) | 0.0311 (8) | 0.0479 (9) | 0.0031 (7) | 0.0060 (7) | 0.0004 (7) |
| N4 | 0.0357 (9) | 0.0329 (9) | 0.0489 (10) | 0.0036 (7) | 0.0053 (7) | -0.0006 (7) |
| C1 | 0.0437 (12) | 0.0385 (12) | 0.0739 (15) | 0.0028 (9) | 0.0084 (10) | 0.0023 (10) |
| C2 | 0.0622 (15) | 0.0361 (12) | 0.0880 (18) | 0.0041 (11) | 0.0175 (13) | 0.0021 (11) |
| C3 | 0.0581 (14) | 0.0450 (13) | 0.0682 (15) | 0.0194 (11) | 0.0194 (11) | 0.0112 (11) |
| C4 | 0.0413 (12) | 0.0615 (15) | 0.0780 (16) | 0.0107 (11) | 0.0039 (11) | 0.0074 (13) |
| C5 | 0.0426 (12) | 0.0425 (12) | 0.0702 (15) | 0.0008 (9) | 0.0049 (10) | 0.0036 (10) |
| C6 | 0.0404 (11) | 0.0367 (10) | 0.0443 (11) | 0.0030 (8) | 0.0099 (9) | 0.0008 (8) |
| C7 | 0.0364 (10) | 0.0350 (10) | 0.0414 (10) | -0.0004 (8) | 0.0091 (8) | 0.0005 (8) |
| C8 | 0.0394 (11) | 0.0332 (10) | 0.0550 (12) | -0.0019 (8) | 0.0013 (9) | -0.0002 (9) |
| C9 | 0.0383 (10) | 0.0400 (11) | 0.0511 (12) | 0.0050 (9) | 0.0091 (9) | 0.0035 (9) |
| C10 | 0.0483 (11) | 0.0312 (10) | 0.0446 (11) | 0.0006 (9) | 0.0071 (9) | -0.0015 (8) |
| C11 | 0.0498 (13) | 0.0416 (12) | 0.0682 (14) | -0.0055 (10) | 0.0075 (11) | -0.0017 (10) |
| C12 | 0.0722 (17) | 0.0486 (13) | 0.0808 (17) | -0.0161 (12) | 0.0210 (14) | -0.0076 (12) |

Geometric parameters (Å, °)

| | | | |
|--------|-----------|--------|-----------|
| S1—C9 | 1.724 (2) | C4—C5 | 1.379 (3) |
| S1—C8 | 1.809 (2) | C4—H4 | 0.9300 |
| F1—C3 | 1.359 (2) | C5—C6 | 1.393 (3) |
| N1—C10 | 1.294 (2) | C5—H5 | 0.9300 |
| N1—N2 | 1.403 (2) | C6—C7 | 1.481 (3) |
| N2—C9 | 1.296 (2) | C7—C8 | 1.504 (3) |
| N3—C9 | 1.368 (3) | C8—H8A | 0.9700 |

| | | | |
|--------------|-------------|---------------|--------------|
| N3—C10 | 1.377 (2) | C8—H8B | 0.9700 |
| N3—N4 | 1.389 (2) | C10—C11 | 1.483 (3) |
| N4—C7 | 1.282 (2) | C11—C12 | 1.504 (3) |
| C1—C6 | 1.383 (3) | C11—H11A | 0.9700 |
| C1—C2 | 1.384 (3) | C11—H11B | 0.9700 |
| C1—H1 | 0.9300 | C12—H12A | 0.9600 |
| C2—C3 | 1.351 (3) | C12—H12B | 0.9600 |
| C2—H2 | 0.9300 | C12—H12C | 0.9600 |
| C3—C4 | 1.359 (3) | | |
| | | | |
| C9—S1—C8 | 94.03 (9) | N4—C7—C8 | 123.28 (17) |
| C10—N1—N2 | 108.10 (16) | C6—C7—C8 | 119.81 (16) |
| C9—N2—N1 | 106.96 (17) | C7—C8—S1 | 112.80 (14) |
| C9—N3—C10 | 105.32 (16) | C7—C8—H8A | 109.0 |
| C9—N3—N4 | 128.67 (16) | S1—C8—H8A | 109.0 |
| C10—N3—N4 | 124.63 (15) | C7—C8—H8B | 109.0 |
| C7—N4—N3 | 115.66 (15) | S1—C8—H8B | 109.0 |
| C6—C1—C2 | 121.1 (2) | H8A—C8—H8B | 107.8 |
| C6—C1—H1 | 119.4 | N2—C9—N3 | 110.27 (17) |
| C2—C1—H1 | 119.4 | N2—C9—S1 | 128.82 (16) |
| C3—C2—C1 | 118.7 (2) | N3—C9—S1 | 120.86 (14) |
| C3—C2—H2 | 120.6 | N1—C10—N3 | 109.33 (17) |
| C1—C2—H2 | 120.6 | N1—C10—C11 | 126.78 (18) |
| C2—C3—C4 | 122.5 (2) | N3—C10—C11 | 123.81 (17) |
| C2—C3—F1 | 119.1 (2) | C10—C11—C12 | 113.32 (19) |
| C4—C3—F1 | 118.4 (2) | C10—C11—H11A | 108.9 |
| C3—C4—C5 | 119.0 (2) | C12—C11—H11A | 108.9 |
| C3—C4—H4 | 120.5 | C10—C11—H11B | 108.9 |
| C5—C4—H4 | 120.5 | C12—C11—H11B | 108.9 |
| C4—C5—C6 | 120.6 (2) | H11A—C11—H11B | 107.7 |
| C4—C5—H5 | 119.7 | C11—C12—H12A | 109.5 |
| C6—C5—H5 | 119.7 | C11—C12—H12B | 109.5 |
| C1—C6—C5 | 118.08 (19) | H12A—C12—H12B | 109.5 |
| C1—C6—C7 | 121.91 (18) | C11—C12—H12C | 109.5 |
| C5—C6—C7 | 120.01 (18) | H12A—C12—H12C | 109.5 |
| N4—C7—C6 | 116.73 (16) | H12B—C12—H12C | 109.5 |
| | | | |
| C10—N1—N2—C9 | 0.3 (2) | N4—C7—C8—S1 | 46.9 (2) |
| C9—N3—N4—C7 | -26.4 (3) | C6—C7—C8—S1 | -138.27 (16) |
| C10—N3—N4—C7 | 168.98 (18) | C9—S1—C8—C7 | -49.59 (16) |
| C6—C1—C2—C3 | -0.1 (4) | N1—N2—C9—N3 | 0.5 (2) |
| C1—C2—C3—C4 | -0.4 (4) | N1—N2—C9—S1 | -177.06 (16) |
| C1—C2—C3—F1 | 179.6 (2) | C10—N3—C9—N2 | -1.0 (2) |
| C2—C3—C4—C5 | 0.3 (4) | N4—N3—C9—N2 | -167.93 (18) |
| F1—C3—C4—C5 | -179.7 (2) | C10—N3—C9—S1 | 176.72 (15) |
| C3—C4—C5—C6 | 0.2 (4) | N4—N3—C9—S1 | 9.8 (3) |
| C2—C1—C6—C5 | 0.6 (3) | C8—S1—C9—N2 | -156.4 (2) |
| C2—C1—C6—C7 | -179.4 (2) | C8—S1—C9—N3 | 26.27 (18) |

| | | | |
|-------------|--------------|----------------|-------------|
| C4—C5—C6—C1 | -0.6 (3) | N2—N1—C10—N3 | -1.0 (2) |
| C4—C5—C6—C7 | 179.3 (2) | N2—N1—C10—C11 | -177.8 (2) |
| N3—N4—C7—C6 | 179.22 (16) | C9—N3—C10—N1 | 1.2 (2) |
| N3—N4—C7—C8 | -5.8 (3) | N4—N3—C10—N1 | 168.80 (17) |
| C1—C6—C7—N4 | -167.62 (19) | C9—N3—C10—C11 | 178.14 (19) |
| C5—C6—C7—N4 | 12.5 (3) | N4—N3—C10—C11 | -14.3 (3) |
| C1—C6—C7—C8 | 17.2 (3) | N1—C10—C11—C12 | -13.0 (3) |
| C5—C6—C7—C8 | -162.73 (19) | N3—C10—C11—C12 | 170.6 (2) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H... <i>A</i> | <i>D</i> —H | H... <i>A</i> | <i>D</i> ... <i>A</i> | <i>D</i> —H... <i>A</i> |
|------------------------------------|-------------|---------------|-----------------------|-------------------------|
| C1—H1...N2 ⁱ | 0.93 | 2.51 | 3.428 (3) | 172 |
| C8—H8 <i>A</i> ...N1 ⁱⁱ | 0.97 | 2.50 | 3.410 (3) | 156 |
| C8—H8 <i>B</i> ...N2 ⁱ | 0.97 | 2.30 | 3.228 (3) | 160 |

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$; (ii) $x, -y+3/2, z+1/2$.