

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

DL-Alaninium iodide

Kevin Lamberts and Ulli Englert*

Institut für Anorganische Chemie, RWTH Aachen University, Landoltweg 1, 52074 Aachen, Germany

Correspondence e-mail: ullrich.englert@ac.rwth-aachen.de


Received 9 May 2012; accepted 15 May 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.026; wR factor = 0.062; data-to-parameter ratio = 27.4.

The crystal structure of DL-alanine hydroiodide (1-carboxy-ethanaminium iodide), $\text{C}_3\text{H}_8\text{NO}_2^+\cdot\text{I}^-$, is that of an organic salt consisting of *N*-protonated cations and iodide anions. The compound features homochiral helices of $\text{N}-\text{H}\cdots\text{O}$ hydrogen-bonded cations in the [010] direction; neighbouring chains are related by crystallographic inversion centers and hence show opposite chirality. The iodide counter-anions act as hydrogen-bond acceptors towards H atoms of the ammonium and carboxy groups, and cross-link the chains along [100]. Thus, an overall two-dimensional network is formed in the *ab* plane. No short contacts occur between iodide anions.

Related literature

For related structures of L-alanine hydrochloride, see: Di Blasio *et al.* (1977), D-alanine alaninium bromide, see: Fischer (2006), L-alanine hydrochloride monohydrate, see: Yamada *et al.* (2008) and DL-alanine hydrochloride, see: Trotter (1962).



Experimental

Crystal data

 $\text{C}_3\text{H}_8\text{NO}_2^+\cdot\text{I}^-$ $M_r = 217.00$

Monoclinic, $P2_1/n$
 $a = 7.6975$ (11) Å
 $b = 5.7776$ (8) Å
 $c = 16.034$ (2) Å
 $\beta = 98.999$ (2)°
 $V = 704.30$ (17) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 4.46$ mm⁻¹
 $T = 100$ K
 $0.30 \times 0.11 \times 0.05$ mm

Data collection

Bruker D8 goniometer with
 SMART APEX CCD detector
 Absorption correction: multi-scan
 (SADABS; Bruker, 2001)
 $T_{\min} = 0.348$, $T_{\max} = 0.808$

10196 measured reflections
 2109 independent reflections
 1887 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.062$
 $S = 1.05$
 2109 reflections
 77 parameters
 3 restraints

H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 1.17$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.87$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{I1}$	0.79 (4)	2.61 (4)	3.391 (2)	171 (3)
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{i}}$	0.87 (3)	2.05 (3)	2.861 (3)	155 (3)
$\text{N1}-\text{H1B}\cdots\text{I1}^{\text{ii}}$	0.88 (3)	2.71 (3)	3.557 (2)	163 (3)
$\text{N1}-\text{H1C}\cdots\text{I1}^{\text{iii}}$	0.87 (3)	2.80 (3)	3.580 (2)	150 (3)

Symmetry codes: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x - 1, y + 1, z$; (iii) $x - 1, y, z$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: publCIF (Westrip, 2010).

Dr Nadine Boymans is gratefully acknowledged for providing us with DL-alanine.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2163).

References

- Bruker (2001). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Di Blasio, B., Pavone, V. & Padone, C. (1977). *Cryst. Struct. Commun.* **6**, 745–748.
 Fischer, A. (2006). *Acta Cryst.* **E62**, o5786–o5788.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Trotter, J. (1962). *Can. J. Chem.* **40**, 1218–1220.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
 Yamada, K., Sato, A., Shimizu, T., Yamazaki, T. & Yokoyama, S. (2008). *Acta Cryst.* **E64**, o806.

supporting information

Acta Cryst. (2012). E68, o1846 [doi:10.1107/S1600536812022003]

DL-Alaninium iodide**Kevin Lamberts and Ulli Englert****S1. Comment**

Our attempt to synthesize a coordination compound from manganese(II)iodide and the racemic α -amino acid DL-alanine failed and unexpectedly led to the formation of the title compound.

The structure of this organic salt consists of one protonated alanine cation and one iodide anion in the asymmetric unit (Fig. 1); the compound crystallizes in the monoclinic space group $P2_1/n$.


All H atoms bonded to electronegative partners find an acceptor in suitable geometry (Table 1), thus forming the maximum number of classical hydrogen bonds. These interactions give rise to double layers (Fig. 2), with the iodide acting as acceptor for one donor from the carboxylic acid OH and two from the ammonium group; the halide adopts a trigonal-planar geometry with respect to these hydrogen bonds. A fourth hydrogen bond is formed between the remaining proton in the ammonium group and a neighbouring carboxylic acid O atom, forming a helical structure along the b -axis (Fig. 3). Each helix is homochiral, but the centrosymmetry of the space group implies the presence of left- and right-handed helices related by crystallographic inversion.

S2. Experimental


MnI₂·4H₂O (0.2 mmol, 74 mg) and DL-alanine (0.4 mmol, 36 mg) were dissolved in 5 ml H₂O/MeOH (1:1) and were left in an open flask at room temperature. After slow evaporation of the solvent a yellow oil remained which was placed in a desiccator. Colorless needles of DL-alanine hydroiodide formed after several weeks.

S3. Refinement


Hydrogen atoms bonded to carbon were included as riding in standard geometry with C—H = 1.00 Å for the methine and C—H = 0.98 Å for the methyl C atom. Coordinates of the hydrogen atoms in the ammonium and in the carboxylic acid groups were refined freely, with the N—H distances restrained to equal length. For all H atoms, $U_{\text{iso}}(\text{H})$ was constrained to 1.2 U_{eq} of the non-H reference atom.

**Figure 1**

PLATON (Spek, 2009) plot with displacement ellipsoids scaled to 80% probability, H atoms shown as spheres with arbitrary radii.

**Figure 2**

Representation of the *C*-face, showing a top view of the two-dimensional layer built by hydrogen bonds (Spek, 2009).

**Figure 3**

View on the *A*-face; homochiral helices extend along [010] (Spek, 2009).

1-carboxyethanaminium iodide

Crystal data

$\text{C}_3\text{H}_8\text{NO}_2^+\text{I}^-$

$M_r = 217.00$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 7.6975$ (11) Å

$b = 5.7776$ (8) Å

$c = 16.034$ (2) Å

$\beta = 98.999$ (2)°

$V = 704.30$ (17) Å³

$Z = 4$

$F(000) = 408$

$D_x = 2.047$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3131 reflections

$\theta = 2.6\text{--}30.3^\circ$

$\mu = 4.46$ mm⁻¹

$T = 100$ K

Needle, colourless

$0.30 \times 0.11 \times 0.05$ mm

Data collection

Bruker D8 goniometer with SMART APEX

CCD detector

diffractometer

Radiation source: INCOATEC microsource

Multilayer optics monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.348$, $T_{\max} = 0.808$

10196 measured reflections

2109 independent reflections

1887 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.055$

$\theta_{\max} = 30.7^\circ$, $\theta_{\min} = 2.6^\circ$

$h = -10 \rightarrow 11$

$k = -8 \rightarrow 8$

$l = -22 \rightarrow 22$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.026$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.062$	$w = 1/[\sigma^2(F_o^2) + (0.020P)^2 + 0.4P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
2109 reflections	$(\Delta/\sigma)_{\max} = 0.001$
77 parameters	$\Delta\rho_{\max} = 1.17 \text{ e } \text{\AA}^{-3}$
3 restraints	$\Delta\rho_{\min} = -1.87 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	0.78595 (2)	0.38517 (3)	0.868059 (10)	0.01622 (7)
O1	0.4337 (3)	0.6529 (4)	0.92827 (13)	0.0248 (5)
H1	0.508 (5)	0.577 (6)	0.913 (2)	0.030*
O2	0.2836 (3)	0.6020 (3)	0.79725 (13)	0.0193 (4)
C1	0.3013 (3)	0.6870 (5)	0.86701 (16)	0.0162 (5)
C2	0.1684 (3)	0.8526 (5)	0.89481 (17)	0.0158 (5)
H2	0.1169	0.7800	0.9420	0.019*
C3	0.2528 (4)	1.0818 (5)	0.9252 (2)	0.0249 (6)
H3A	0.1619	1.1888	0.9380	0.030*
H3B	0.3120	1.1483	0.8809	0.030*
H3C	0.3390	1.0558	0.9761	0.030*
N1	0.0254 (3)	0.8877 (4)	0.82138 (14)	0.0143 (4)
H1A	0.060 (4)	0.928 (5)	0.7744 (16)	0.017*
H1B	-0.052 (4)	0.988 (5)	0.834 (2)	0.017*
H1C	-0.036 (4)	0.761 (4)	0.811 (2)	0.017*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.01442 (11)	0.01740 (11)	0.01628 (10)	0.00331 (6)	0.00066 (7)	0.00025 (6)
O1	0.0196 (10)	0.0337 (12)	0.0190 (10)	0.0129 (9)	-0.0036 (8)	-0.0060 (9)
O2	0.0183 (10)	0.0227 (11)	0.0159 (9)	0.0043 (7)	0.0000 (8)	-0.0040 (7)
C1	0.0153 (12)	0.0167 (13)	0.0164 (12)	0.0011 (9)	0.0021 (9)	0.0011 (9)
C2	0.0146 (12)	0.0185 (13)	0.0136 (11)	0.0032 (9)	-0.0005 (9)	-0.0007 (9)

C3	0.0220 (14)	0.0223 (14)	0.0272 (15)	0.0040 (11)	-0.0059 (12)	-0.0102 (12)
N1	0.0146 (11)	0.0154 (11)	0.0119 (10)	0.0010 (8)	-0.0006 (8)	0.0005 (8)

Geometric parameters (Å, °)

O1—C1	1.315 (3)	C3—H3A	0.98
O1—H1	0.79 (4)	C3—H3B	0.98
O2—C1	1.210 (3)	C3—H3C	0.98
C1—C2	1.517 (4)	N1—H1A	0.87 (2)
C2—N1	1.495 (3)	N1—H1B	0.88 (2)
C2—C3	1.521 (4)	N1—H1C	0.87 (2)
C2—H2	1.00		
C1—O1—H1	111 (3)	C2—C3—H3B	109.5
O2—C1—O1	126.3 (3)	H3A—C3—H3B	109.5
O2—C1—C2	123.0 (2)	C2—C3—H3C	109.5
O1—C1—C2	110.8 (2)	H3A—C3—H3C	109.5
N1—C2—C1	107.5 (2)	H3B—C3—H3C	109.5
N1—C2—C3	111.1 (2)	C2—N1—H1A	115 (2)
C1—C2—C3	111.7 (2)	C2—N1—H1B	110 (2)
N1—C2—H2	108.8	H1A—N1—H1B	110 (3)
C1—C2—H2	108.8	C2—N1—H1C	110 (2)
C3—C2—H2	108.8	H1A—N1—H1C	107 (3)
C2—C3—H3A	109.5	H1B—N1—H1C	103 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...I1	0.79 (4)	2.61 (4)	3.391 (2)	171 (3)
N1—H1A...O2 ⁱ	0.87 (3)	2.05 (3)	2.861 (3)	155 (3)
N1—H1B...I1 ⁱⁱ	0.88 (3)	2.71 (3)	3.557 (2)	163 (3)
N1—H1C...I1 ⁱⁱⁱ	0.87 (3)	2.80 (3)	3.580 (2)	150 (3)

Symmetry codes: (i) $-x+1/2, y+1/2, -z+3/2$; (ii) $x-1, y+1, z$; (iii) $x-1, y, z$.