

2-{[3-Methyl-4-(2,2,2-trifluoroethoxy)pyridin-2-yl]methylsulfanyl}-1*H*-benzimidazole propan-2-ol monosolvate: a second monoclinic polymorph

Jin-Ju Ma,^{a,b} Ming-Hui Qi,^b Ming-Huang Hong,^b Jie Lu^c and Guo-Bin Ren^{a,b*}

^aSchool of Chemical Engineering and Energy, Zhengzhou University, Zhengzhou 450001, People's Republic of China, ^bPharmaceutical Crystal Engineering Research Group, Shanghai Institute of Pharmaceutical Industry, 1320 Beijing Road (West), Shanghai 200040, People's Republic of China, and ^cNational Engineering Laboratory for Cereal Fermentation Technology, School of Chemical & Material Engineering, Jiangnan University, Wuxi 214122, People's Republic of China

Correspondence e-mail: renguobin2557@yahoo.com.cn


Received 10 May 2012; accepted 16 May 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.040; wR factor = 0.116; data-to-parameter ratio = 13.4.

In the crystal structure of the title compound, $\text{C}_{16}\text{H}_{14}\text{F}_3\text{N}_3\text{OS}\cdots\text{C}_3\text{H}_8\text{O}$, the molecules are linked into chains along [010] via $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds. The triclinic form was reported by Ren *et al.* [(2011). *Acta Cryst. E67*, o270] and the first monoclinic form by Chen *et al.* [(2012). *Acta Cryst. E68*, o2015–o2016]. The fused five-and six-membered rings make a dihedral angle of $1.22(2)^\circ$, while the benzene and pyridine rings make a dihedral angle of $10.15(2)^\circ$.

Related literature

For the use of the title compound as an intermediate in the synthesis of the anti-ulcer drug lansoprazole [systematic name: (*RS*)-2-([3-methyl-4-(2,2,2-trifluoroethoxy)pyridin-2-yl)methylsulfanyl)-1*H*-benzo[*d*]imidazole], see: Del Rio *et al.* (2007); Reddy *et al.* (2008); Iwahashi *et al.* (1991). For related structures, see: Swamy & Ravikumar (2007); Hakim *et al.* (2010). For the triclinic polymorph of the title propan-2-ol solvo-polymorph, see: Ren *et al.* (2011) and for the monoclinic monohydrate, see: Chen *et al.* (2012).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{14}\text{F}_3\text{N}_3\text{OS}\cdots\text{C}_3\text{H}_8\text{O}$	$V = 1969.60(5)\text{ \AA}^3$
$M_r = 413.46$	$Z = 4$
Monoclinic, $P2_1/c$	$\text{Cu } K\alpha$ radiation
$a = 17.4583(2)\text{ \AA}$	$\mu = 1.89\text{ mm}^{-1}$
$b = 7.4162(1)\text{ \AA}$	$T = 296\text{ K}$
$c = 16.9622(2)\text{ \AA}$	$0.31 \times 0.23 \times 0.15\text{ mm}$
$\beta = 116.255(2)^\circ$	

Data collection

Bruker APEXII diffractometer	13351 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	3410 independent reflections
$(SADABS$; Bruker, 2009)	3266 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.592$, $T_{\max} = 0.765$	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	254 parameters
$wR(F^2) = 0.116$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\max} = 0.38\text{ e \AA}^{-3}$
3410 reflections	$\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2B \cdots N1 ⁱ	0.82	2.01	2.8142 (18)	167
N2—H2A \cdots O2 ⁱⁱ	0.86	1.98	2.8027 (18)	161

Symmetry codes: (i) $x, y, z - 1$; (ii) $x, y - 1, z + 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the New Drug Innovation (2009ZX09301-007) project of the Ministry of Science and Technology of China and the National Natural Science Foundation of China (Nos. 81102391 and 21176102).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2460).

References

- Bruker (2009). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, Y.-F., Chen, J.-Y., Hong, M.-H., Lu, J. & Ren, G.-B. (2012). *Acta Cryst. E68*, o2015–o2016.
- Del Rio, R. E., Wang, B., Achab, S. & Bohe, L. (2007). *Org. Lett.* **9**, 2265–2268.
- Hakim Al-arique, Q. N. M., Jasinski, J. P., Butcher, R. J., Yathirajan, H. S. & Narayana, B. (2010). *Acta Cryst. E66*, o1507–o1508.
- Iwahashi, T., Satoh, H., Nakao, M., Iwasaki, T., Yamazaki, T., Kubo, K., Tamura, T. & Imada, A. (1991). *Antimicrob. Agents Chemother.* **35**, 490–496.
- Reddy, G. M., Mukkanti, K., Kumar, T., Babu, J., Moses, M. & Reddy, P. P. (2008). *Synth. Commun.* **38**, 3477–3489.
- Ren, G.-B., Hong, M.-H., Zhong, J.-L., Yi, D.-X. & Xu, L.-H. (2011). *Acta Cryst. E67*, o270.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Swamy, G. Y. S. K. & Ravikumar, K. (2007). *J. Struct. Chem.* **48**, 715–718.

supporting information

Acta Cryst. (2012). E68, o2017 [https://doi.org/10.1107/S1600536812022143]

2-{[3-Methyl-4-(2,2,2-trifluoroethoxy)pyridin-2-yl]methylsulfanyl}-1*H*-benzimidazole propan-2-ol monosolvate: a second monoclinic polymorph

Jin-Ju Ma, Ming-Hui Qi, Ming-Huang Hong, Jie Lu and Guo-Bin Ren

S1. Comment

Lansoprazole (Del Rio *et al.*, 2007; Reddy *et al.*, 2008) and many of its analogues are characterized by an anti-ulcer effect (Iwahashi *et al.*, 1991). The title compound, (I), is the critical reaction intermediate of lansoprazole. Recently, the compound was successfully crystallized from 2-propanol, and the crystal structure is now firstly reported.

The crystal structure (Fig 1) contains one independent molecule and one 2-propanol solvato which are involved in the formation of hydrogen-bonded chains running along [010] via N—H···O and O—H···N hydrogen bonds (Table 1 and Fig. 2). The 2-propanol molecules acts as a hydrogen-bond bridge, providing further stability to the crystal lattice.

S2. Experimental

The raw material was kindly provided by Shanghai Enran Sci-Tech Investment Management Co., Ltd. The compound was dissolved in 2-propanol and suitable crystals of X-ray were obtained by slow evaporation at room temperature over a period of one week.

S3. Refinement

All C—H atoms were constrained to an ideal geometry with C—H distances of 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH; 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH_2 ; 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for CH_3 ; 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for OH atoms; and 0.86 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for NH atoms.





Figure 1

The content of asymmetric unit (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. Dashed lines denote hydrogen bonds. H atoms have been omitted for clarity.

**Figure 2**

A packing diagram, projected along the [010] chains.

2-{{[3-Methyl-4-(2,2,2-trifluoroethoxy)pyridin-2-yl]methylsulfanyl}- 1*H*-benzimidazole propan-2-ol monosolvate

Crystal data



$M_r = 413.46$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 17.4583 (2)$ Å

$b = 7.4162 (1)$ Å

$c = 16.9622 (2)$ Å

$\beta = 116.255 (2)^\circ$

$V = 1969.60 (5)$ Å³

$Z = 4$

$F(000) = 864$

$D_x = 1.394$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 3266 reflections

$\theta = 6.1\text{--}67.0^\circ$

$\mu = 1.89$ mm⁻¹

$T = 296$ K

Column, colorless

0.31 × 0.23 × 0.15 mm

Data collection

Bruker APEXII
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 0 pixels mm⁻¹

phi and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2009)

$T_{\min} = 0.592$, $T_{\max} = 0.765$

13351 measured reflections

3410 independent reflections

3266 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 67.0^\circ$, $\theta_{\min} = 6.1^\circ$

$h = -19 \rightarrow 20$

$k = -8 \rightarrow 8$

$l = -20 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.116$

$S = 1.06$

3410 reflections

254 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0697P)^2 + 0.5861P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick,

$$2008), Fc^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$$

Extinction coefficient: 0.0024 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.22271 (3)	0.11157 (5)	1.05242 (3)	0.04524 (17)
O1	0.09818 (8)	0.61043 (16)	0.71885 (7)	0.0497 (3)
O2	0.32047 (8)	0.67645 (16)	0.17242 (10)	0.0573 (4)
H2B	0.3016	0.5801	0.1800	0.086*
N1	0.28104 (9)	0.32274 (18)	1.20227 (9)	0.0417 (3)
N2	0.29844 (8)	0.02492 (18)	1.22301 (8)	0.0419 (3)
H2A	0.2958	-0.0880	1.2105	0.050*
N3	0.14436 (10)	0.1600 (2)	0.87641 (9)	0.0476 (4)
F1	0.13495 (9)	0.8359 (2)	0.61276 (11)	0.0938 (5)
F2	0.01560 (8)	0.89596 (15)	0.61011 (8)	0.0681 (3)
F3	0.02040 (11)	0.76215 (19)	0.50116 (8)	0.0893 (5)
C1	0.26983 (10)	0.1637 (2)	1.16472 (10)	0.0375 (3)
C1'	0.47122 (13)	0.6930 (4)	0.26857 (17)	0.0747 (6)
H1'A	0.4639	0.6167	0.3104	0.112*
H1'B	0.4658	0.8169	0.2816	0.112*
H1'C	0.5269	0.6734	0.2717	0.112*
C2	0.32070 (10)	0.2857 (2)	1.29233 (11)	0.0415 (4)
C2'	0.40380 (12)	0.6487 (3)	0.17709 (15)	0.0594 (5)
H2'A	0.4095	0.5217	0.1646	0.071*
C3	0.34706 (13)	0.4014 (3)	1.36387 (13)	0.0556 (5)
H3B	0.3384	0.5251	1.3561	0.067*
C3'	0.41146 (15)	0.7633 (4)	0.10808 (17)	0.0729 (6)
H3'A	0.3673	0.7311	0.0513	0.109*
H3'B	0.4663	0.7440	0.1092	0.109*
H3'C	0.4057	0.8880	0.1196	0.109*
C4	0.38645 (14)	0.3274 (3)	1.44666 (13)	0.0631 (5)
H4A	0.4048	0.4029	1.4954	0.076*
C5	0.39947 (13)	0.1429 (3)	1.45925 (13)	0.0595 (5)
H5A	0.4272	0.0979	1.5162	0.071*

C6	0.37235 (11)	0.0253 (3)	1.38953 (11)	0.0519 (4)
H6A	0.3803	-0.0985	1.3980	0.062*
C7	0.33245 (10)	0.0997 (2)	1.30578 (11)	0.0408 (4)
C8	0.20875 (11)	0.3377 (2)	1.00826 (11)	0.0443 (4)
H8A	0.1736	0.4076	1.0281	0.053*
H8B	0.2638	0.3969	1.0285	0.053*
C9	0.16630 (10)	0.3261 (2)	0.90919 (10)	0.0385 (3)
C10	0.15158 (10)	0.4812 (2)	0.85880 (10)	0.0393 (4)
C11	0.11201 (10)	0.4568 (2)	0.76766 (10)	0.0405 (4)
C12	0.08915 (12)	0.2862 (2)	0.73202 (11)	0.0493 (4)
H12A	0.0628	0.2686	0.6715	0.059*
C13	0.10670 (13)	0.1429 (2)	0.78931 (12)	0.0536 (5)
H13A	0.0913	0.0278	0.7657	0.064*
C14	0.05724 (12)	0.5917 (2)	0.62635 (11)	0.0454 (4)
H14A	-0.0010	0.5500	0.6070	0.054*
H14B	0.0875	0.5051	0.6076	0.054*
C15	0.05779 (11)	0.7721 (3)	0.58855 (11)	0.0495 (4)
C16	0.17477 (13)	0.6669 (3)	0.89758 (12)	0.0548 (5)
H16A	0.1592	0.7539	0.8511	0.082*
H16B	0.2352	0.6730	0.9344	0.082*
H16C	0.1448	0.6924	0.9320	0.082*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0616 (3)	0.0325 (3)	0.0362 (2)	-0.00018 (16)	0.0168 (2)	0.00201 (14)
O1	0.0649 (8)	0.0411 (7)	0.0345 (6)	-0.0002 (5)	0.0143 (5)	0.0050 (5)
O2	0.0481 (7)	0.0306 (6)	0.0920 (10)	-0.0005 (5)	0.0299 (7)	0.0013 (6)
N1	0.0517 (8)	0.0324 (7)	0.0404 (7)	0.0030 (6)	0.0199 (6)	0.0036 (5)
N2	0.0523 (8)	0.0304 (7)	0.0395 (7)	0.0015 (6)	0.0172 (6)	0.0040 (5)
N3	0.0613 (9)	0.0367 (7)	0.0395 (8)	-0.0012 (6)	0.0174 (7)	0.0000 (6)
F1	0.0650 (8)	0.1046 (11)	0.1088 (12)	-0.0198 (7)	0.0357 (8)	0.0334 (9)
F2	0.0844 (8)	0.0471 (7)	0.0706 (8)	0.0098 (5)	0.0322 (7)	0.0056 (5)
F3	0.1476 (14)	0.0680 (8)	0.0367 (6)	-0.0046 (8)	0.0267 (7)	0.0095 (6)
C1	0.0419 (8)	0.0318 (8)	0.0383 (8)	0.0007 (6)	0.0172 (7)	0.0037 (6)
C1'	0.0519 (11)	0.0715 (15)	0.0859 (16)	0.0019 (10)	0.0171 (11)	0.0122 (12)
C2	0.0444 (8)	0.0411 (9)	0.0408 (9)	0.0024 (7)	0.0203 (7)	0.0022 (7)
C2'	0.0493 (10)	0.0350 (9)	0.0896 (15)	0.0018 (8)	0.0268 (10)	-0.0054 (9)
C3	0.0682 (12)	0.0485 (11)	0.0517 (11)	-0.0001 (9)	0.0280 (9)	-0.0078 (8)
C3'	0.0667 (13)	0.0789 (16)	0.0785 (15)	-0.0071 (11)	0.0369 (12)	-0.0080 (12)
C4	0.0703 (12)	0.0766 (14)	0.0434 (10)	-0.0044 (11)	0.0260 (9)	-0.0130 (9)
C5	0.0584 (11)	0.0791 (14)	0.0382 (9)	0.0024 (10)	0.0188 (8)	0.0082 (9)
C6	0.0554 (10)	0.0533 (11)	0.0442 (10)	0.0035 (8)	0.0195 (8)	0.0122 (8)
C7	0.0419 (8)	0.0412 (9)	0.0401 (8)	0.0006 (6)	0.0190 (7)	0.0044 (6)
C8	0.0566 (10)	0.0334 (8)	0.0365 (9)	0.0009 (7)	0.0147 (7)	0.0028 (7)
C9	0.0404 (8)	0.0363 (8)	0.0365 (8)	0.0010 (6)	0.0149 (6)	0.0004 (6)
C10	0.0406 (8)	0.0374 (8)	0.0367 (8)	0.0014 (6)	0.0143 (6)	0.0014 (6)
C11	0.0425 (8)	0.0388 (8)	0.0375 (8)	0.0016 (7)	0.0152 (6)	0.0044 (7)

C12	0.0595 (10)	0.0467 (10)	0.0337 (8)	-0.0012 (8)	0.0134 (7)	-0.0017 (7)
C13	0.0726 (12)	0.0376 (9)	0.0421 (9)	-0.0051 (8)	0.0176 (9)	-0.0053 (7)
C14	0.0536 (9)	0.0453 (10)	0.0341 (8)	0.0029 (7)	0.0165 (7)	0.0017 (7)
C15	0.0548 (10)	0.0539 (11)	0.0365 (8)	-0.0026 (8)	0.0170 (7)	0.0045 (7)
C16	0.0733 (12)	0.0383 (9)	0.0422 (9)	-0.0022 (8)	0.0159 (9)	0.0010 (7)

Geometric parameters (\AA , $^\circ$)

S1—C1	1.7517 (16)	C3'—H3'A	0.9600
S1—C8	1.8088 (17)	C3'—H3'B	0.9600
O1—C11	1.366 (2)	C3'—H3'C	0.9600
O1—C14	1.414 (2)	C4—C5	1.388 (3)
O2—C2'	1.436 (2)	C4—H4A	0.9300
O2—H2B	0.8200	C5—C6	1.374 (3)
N1—C1	1.313 (2)	C5—H5A	0.9300
N1—C2	1.397 (2)	C6—C7	1.390 (2)
N2—C1	1.360 (2)	C6—H6A	0.9300
N2—C7	1.376 (2)	C8—C9	1.510 (2)
N2—H2A	0.8600	C8—H8A	0.9700
N3—C13	1.331 (2)	C8—H8B	0.9700
N3—C9	1.337 (2)	C9—C10	1.387 (2)
F1—C15	1.311 (2)	C10—C11	1.398 (2)
F2—C15	1.326 (2)	C10—C16	1.502 (2)
F3—C15	1.331 (2)	C11—C12	1.383 (3)
C1'—C2'	1.512 (3)	C12—C13	1.380 (3)
C1'—H1'A	0.9600	C12—H12A	0.9300
C1'—H1'B	0.9600	C13—H13A	0.9300
C1'—H1'C	0.9600	C14—C15	1.486 (3)
C2—C3	1.388 (3)	C14—H14A	0.9700
C2—C7	1.398 (2)	C14—H14B	0.9700
C2'—C3'	1.499 (3)	C16—H16A	0.9600
C2'—H2'A	0.9800	C16—H16B	0.9600
C3—C4	1.375 (3)	C16—H16C	0.9600
C3—H3B	0.9300		
C1—S1—C8	99.09 (8)	C7—C6—H6A	121.5
C11—O1—C14	117.24 (13)	N2—C7—C6	132.68 (16)
C2'—O2—H2B	109.5	N2—C7—C2	105.36 (14)
C1—N1—C2	104.42 (13)	C6—C7—C2	121.96 (17)
C1—N2—C7	106.91 (13)	C9—C8—S1	108.55 (11)
C1—N2—H2A	126.5	C9—C8—H8A	110.0
C7—N2—H2A	126.5	S1—C8—H8A	110.0
C13—N3—C9	117.35 (15)	C9—C8—H8B	110.0
N1—C1—N2	113.52 (14)	S1—C8—H8B	110.0
N1—C1—S1	128.56 (12)	H8A—C8—H8B	108.4
N2—C1—S1	117.92 (12)	N3—C9—C10	124.55 (15)
C2'—C1'—H1'A	109.5	N3—C9—C8	115.14 (14)
C2'—C1'—H1'B	109.5	C10—C9—C8	120.30 (14)

H1'A—C1'—H1'B	109.5	C9—C10—C11	116.09 (15)
C2'—C1'—H1'C	109.5	C9—C10—C16	123.33 (14)
H1'A—C1'—H1'C	109.5	C11—C10—C16	120.58 (14)
H1'B—C1'—H1'C	109.5	O1—C11—C12	123.98 (14)
C3—C2—N1	130.27 (16)	O1—C11—C10	115.48 (14)
C3—C2—C7	119.94 (16)	C12—C11—C10	120.53 (15)
N1—C2—C7	109.79 (15)	C13—C12—C11	117.74 (16)
O2—C2'—C3'	108.28 (17)	C13—C12—H12A	121.1
O2—C2'—C1'	109.59 (18)	C11—C12—H12A	121.1
C3'—C2'—C1'	112.37 (18)	N3—C13—C12	123.72 (17)
O2—C2'—H2'A	108.8	N3—C13—H13A	118.1
C3'—C2'—H2'A	108.8	C12—C13—H13A	118.1
C1'—C2'—H2'A	108.8	O1—C14—C15	107.09 (14)
C4—C3—C2	117.96 (19)	O1—C14—H14A	110.3
C4—C3—H3B	121.0	C15—C14—H14A	110.3
C2—C3—H3B	121.0	O1—C14—H14B	110.3
C2'—C3'—H3'A	109.5	C15—C14—H14B	110.3
C2'—C3'—H3'B	109.5	H14A—C14—H14B	108.6
H3'A—C3'—H3'B	109.5	F1—C15—F2	106.44 (17)
C2'—C3'—H3'C	109.5	F1—C15—F3	107.37 (16)
H3'A—C3'—H3'C	109.5	F2—C15—F3	106.68 (16)
H3'B—C3'—H3'C	109.5	F1—C15—C14	113.07 (16)
C3—C4—C5	121.61 (19)	F2—C15—C14	113.23 (14)
C3—C4—H4A	119.2	F3—C15—C14	109.68 (15)
C5—C4—H4A	119.2	C10—C16—H16A	109.5
C6—C5—C4	121.53 (18)	C10—C16—H16B	109.5
C6—C5—H5A	119.2	H16A—C16—H16B	109.5
C4—C5—H5A	119.2	C10—C16—H16C	109.5
C5—C6—C7	116.97 (19)	H16A—C16—H16C	109.5
C5—C6—H6A	121.5	H16B—C16—H16C	109.5
C2—N1—C1—N2	0.47 (18)	C13—N3—C9—C10	0.8 (3)
C2—N1—C1—S1	179.51 (12)	C13—N3—C9—C8	179.99 (15)
C7—N2—C1—N1	-0.99 (18)	S1—C8—C9—N3	3.65 (19)
C7—N2—C1—S1	179.85 (11)	S1—C8—C9—C10	-177.14 (12)
C8—S1—C1—N1	6.99 (17)	N3—C9—C10—C11	-0.9 (2)
C8—S1—C1—N2	-174.00 (12)	C8—C9—C10—C11	179.94 (14)
C1—N1—C2—C3	-179.33 (18)	N3—C9—C10—C16	178.21 (17)
C1—N1—C2—C7	0.23 (18)	C8—C9—C10—C16	-0.9 (2)
N1—C2—C3—C4	-178.76 (18)	C14—O1—C11—C12	0.6 (2)
C7—C2—C3—C4	1.7 (3)	C14—O1—C11—C10	-179.17 (14)
C2—C3—C4—C5	-0.3 (3)	C9—C10—C11—O1	-179.78 (13)
C3—C4—C5—C6	-1.1 (3)	C16—C10—C11—O1	1.1 (2)
C4—C5—C6—C7	1.0 (3)	C9—C10—C11—C12	0.5 (2)
C1—N2—C7—C6	-178.20 (18)	C16—C10—C11—C12	-178.69 (17)
C1—N2—C7—C2	1.05 (17)	O1—C11—C12—C13	-179.70 (17)
C5—C6—C7—N2	179.62 (17)	C10—C11—C12—C13	0.0 (3)
C5—C6—C7—C2	0.5 (3)	C9—N3—C13—C12	-0.2 (3)

C3—C2—C7—N2	178.81 (15)	C11—C12—C13—N3	−0.2 (3)
N1—C2—C7—N2	−0.81 (18)	C11—O1—C14—C15	−174.15 (14)
C3—C2—C7—C6	−1.8 (3)	O1—C14—C15—F1	59.7 (2)
N1—C2—C7—C6	178.54 (15)	O1—C14—C15—F2	−61.42 (19)
C1—S1—C8—C9	−178.94 (12)	O1—C14—C15—F3	179.55 (15)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O2—H2B···N1 ⁱ	0.82	2.01	2.8142 (18)	167
N2—H2A···O2 ⁱⁱ	0.86	1.98	2.8027 (18)	161

Symmetry codes: (i) $x, y, z-1$; (ii) $x, y-1, z+1$.