

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(4E)-N-[(2-Chlorophenyl)methoxy]-1,3-dimethyl-2,6-diphenylpiperidin-4-imine

Chennan Ramalingan,^{a,‡} Seik Weng Ng^{b,c} and Edward R. T. Tiekink^{b,*}

^aCentre for Nanotechnology, Department of Chemistry, Kalasalingam University, Krishnankoil 626 126, Tamilnadu, India, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department and Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia
Correspondence e-mail: edward.tiekink@gmail.com


Received 25 June 2012; accepted 25 June 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.045; wR factor = 0.118; data-to-parameter ratio = 18.8.

In the title compound, $\text{C}_{26}\text{H}_{27}\text{ClN}_2\text{O}$, the piperidine ring has a chair conformation and all of the ring substituents at C_{sp^3} atoms occupy equatorial positions. The dihedral angle formed between the phenyl rings is 48.11 (9°). The chlorobenzene ring occupies a position orthogonal to the methoxy(methylidene)amine residue [$\text{N}-\text{O}-\text{C}-\text{C}$ torsion angle = -87.90 (15°)]. The conformation about the imine $\text{C}=\text{N}$ bond [1.278 (2) Å] is *E*, and the chloro substituent is *anti* to the piperidine N atom. Helical supramolecular chains along [010] are sustained by $\text{C}-\text{H}\cdots\pi$ interactions in the crystal packing.

Related literature

For the biological activity of molecules having a 2,6-diaryl-piperidine core, see: Ramachandran *et al.* (2011); Ramalingan *et al.* (2004). For the structure of the bromo derivative, see: Ramalingan *et al.* (2012). For the synthesis, see: Ramalingan *et al.* (2006).



Experimental

Crystal data

 $\text{C}_{26}\text{H}_{27}\text{ClN}_2\text{O}$ $M_r = 418.95$

Monoclinic, $C2/c$
 $a = 20.3043$ (8) Å
 $b = 6.8811$ (3) Å
 $c = 32.2244$ (12) Å
 $\beta = 98.478$ (4°)
 $V = 4453.1$ (3) Å³

$Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.19$ mm⁻¹
 $T = 100$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2012)
 $T_{\min} = 0.846$, $T_{\max} = 1.000$

14305 measured reflections
5108 independent reflections
3847 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.118$
 $S = 1.02$
5108 reflections

271 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29$ e Å⁻³
 $\Delta\rho_{\min} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and Cg2 are the centroids of the C1–C6 and C15–C20 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C17}-\text{H17}\cdots\text{Cg1}^{\text{i}}$	0.95	2.69	3.556 (2)	151
$\text{C3}-\text{H3}\cdots\text{Cg2}^{\text{ii}}$	0.95	2.90	3.6852 (19)	141

Symmetry codes: (i) $x + \frac{3}{2}, y + \frac{1}{2}, z + 1$; (ii) $x + \frac{3}{2}, y + \frac{3}{2}, z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors are grateful for facilities provided by the Chairman/Management of Kalasalingam University, and thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5958).

References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Ramachandran, R., Rani, M., Senthana, S., Jeong, Y.-T. & Kabilan, S. (2011). *Eur. J. Med. Chem.* **46**, 1926–1934.
Ramalingan, C., Balasubramanian, S., Kabilan, S. & Vasudevan, M. (2004). *Eur. J. Med. Chem.* **39**, 527–533.
Ramalingan, C., Ng, S. E. & Tiekink, E. R. T. (2012). *Acta Cryst.* **E68**, o2267.
Ramalingan, C., Park, Y.-T. & Kabilan, S. (2006). *Eur. J. Med. Chem.* **41**, 683–696.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

[‡] Additional correspondence author, e-mail: ramalinganc@gmail.com.

supporting information

Acta Cryst. (2012). E68, o2268 [https://doi.org/10.1107/S1600536812028899]

(4*E*)-*N*-[(2-Chlorophenyl)methoxy]-1,3-dimethyl-2,6-diphenylpiperidin-4-imine**Chennan Ramalingan, Seik Weng Ng and Edward R. T. Tiekink****S1. Comment**

A diverse range of molecules possessing a 2,6-diaryl piperidine core exhibit potent biological activities (Ramachandran *et al.*, 2011; Ramalingan *et al.*, 2004). In a wide search program towards generating efficient biological agents, the title compound, (I), was synthesized (Ramalingan *et al.*, 2006). Herein, the crystal and molecular structure of (I) is described.

In (I), Fig. 1, the piperidine ring has a chair conformation and all of the ring-substituents occupy equatorial positions. In the bromo derivative (Ramalingan *et al.*, 2012) all but the *N*-bound substituent, which occupies a bisectonal position, also occupy equatorial positions, in accord with (I). The dihedral angle formed between the C15–C20 and C21–C26 phenyl rings is 48.11 (9)°, and each forms a dihedral angle of 64.68 (8) and 72.57 (9)°, respectively, with the chlorobenzene ring, which occupies a position orthogonal to the methoxy(methylidene)amine residue as seen in the N1—O1—C7—C6 torsion angle of -87.90 (15)°. The conformation about the imine C8=N1 bond [1.278 (2) Å] is *E*. The chloro substituent is *anti* to the piperidine-N atom.

In the crystal packing, helical supramolecular chains along [0 1 0] are sustained by C—H⋯ π interactions, Fig. 2 and Table 1. These assemble into a three-dimensional architecture without specific intermolecular interactions between the chains, Fig. 3.

S2. Experimental

For full details of the synthesis, refer to Ramalingan *et al.* (2006). Re-crystallization was performed by slow evaporation of an ethanolic solution of (I) which afforded colourless crystals. *M.pt.*: 361–362 K.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H = 0.95–0.99 Å, $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. Owing to poor agreement, two reflections, *i.e.* (0 1 0) and (1 2 10), were omitted from the final refinement.




Figure 1

The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Figure 2

A view of the helical supramolecular chain in (I) sustained by C—H... π interactions which are shown as purple dashed lines.




Figure 3

A view in projection down the *b* axis of the unit-cell contents for (I). The C—H... π interactions are shown as purple dashed lines. One chain has been highlighted in space-filling mode.

(4E)-N-[(2-Chlorophenyl)methoxy]-1,3-dimethyl-2,6-diphenylpiperidin-4-imine

Crystal data

$C_{26}H_{27}ClN_2O$

$M_r = 418.95$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 20.3043\ (8)\ \text{\AA}$

$b = 6.8811\ (3)\ \text{\AA}$

$c = 32.2244\ (12)\ \text{\AA}$

$\beta = 98.478\ (4)^\circ$

$V = 4453.1\ (3)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1776$

$D_x = 1.250\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 4543 reflections

$\theta = 2.2\text{--}27.5^\circ$

$\mu = 0.19\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Prism, colourless

$0.30 \times 0.25 \times 0.20\ \text{mm}$

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas detector
Radiation source: SuperNova (Mo) X-ray
Source
Mirror monochromator
Detector resolution: 10.4041 pixels mm⁻¹
 ω scan
Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.846$, $T_{\max} = 1.000$
14305 measured reflections
5108 independent reflections
3847 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 $\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -19 \rightarrow 26$
 $k = -8 \rightarrow 8$
 $l = -41 \rightarrow 40$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.118$
 $S = 1.02$
5108 reflections
271 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0478P)^2 + 2.1668P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.76604 (2)	0.38812 (7)	0.564062 (15)	0.03838 (15)
O1	0.63144 (6)	0.72592 (17)	0.47058 (3)	0.0276 (3)
N1	0.64825 (7)	0.6556 (2)	0.43184 (4)	0.0244 (3)
N2	0.56479 (6)	0.9658 (2)	0.32797 (4)	0.0210 (3)
C1	0.68498 (8)	0.3178 (2)	0.54424 (5)	0.0233 (4)
C2	0.66062 (9)	0.1447 (3)	0.55791 (5)	0.0287 (4)
H2	0.6880	0.0643	0.5773	0.034*
C3	0.59581 (9)	0.0907 (3)	0.54287 (5)	0.0280 (4)
H3	0.5783	-0.0271	0.5521	0.034*
C4	0.55655 (8)	0.2081 (3)	0.51448 (5)	0.0258 (4)
H4	0.5122	0.1705	0.5041	0.031*
C5	0.58196 (8)	0.3814 (2)	0.50108 (5)	0.0221 (4)
H5	0.5546	0.4610	0.4815	0.027*
C6	0.64648 (8)	0.4393 (2)	0.51585 (5)	0.0199 (3)
C7	0.67419 (9)	0.6302 (3)	0.50347 (5)	0.0256 (4)
H7A	0.6815	0.7164	0.5283	0.031*

H7B	0.7179	0.6071	0.4943	0.031*
C8	0.61070 (8)	0.7273 (2)	0.40023 (5)	0.0227 (4)
C9	0.55388 (8)	0.8646 (3)	0.40074 (5)	0.0260 (4)
H9A	0.5550	0.9163	0.4295	0.031*
H9B	0.5115	0.7929	0.3932	0.031*
C10	0.55575 (8)	1.0353 (2)	0.37012 (5)	0.0219 (4)
H10	0.5940	1.1217	0.3810	0.026*
C11	0.62664 (8)	0.8506 (2)	0.33000 (5)	0.0214 (4)
H11	0.6646	0.9332	0.3431	0.026*
C12	0.62416 (8)	0.6675 (2)	0.35738 (5)	0.0223 (4)
H12	0.5858	0.5863	0.3443	0.027*
C13	0.68726 (9)	0.5454 (3)	0.35879 (5)	0.0286 (4)
H13A	0.6845	0.4323	0.3769	0.043*
H13B	0.7261	0.6240	0.3700	0.043*
H13C	0.6917	0.5016	0.3304	0.043*
C14	0.56727 (9)	1.1334 (3)	0.30007 (5)	0.0288 (4)
H14A	0.5749	1.0881	0.2723	0.043*
H14B	0.6037	1.2199	0.3118	0.043*
H14C	0.5250	1.2040	0.2975	0.043*
C15	0.49163 (8)	1.1488 (2)	0.36936 (5)	0.0213 (3)
C16	0.49022 (8)	1.3168 (2)	0.39308 (5)	0.0237 (4)
H16	0.5306	1.3691	0.4074	0.028*
C17	0.43008 (9)	1.4094 (3)	0.39607 (5)	0.0271 (4)
H17	0.4296	1.5236	0.4126	0.033*
C18	0.37128 (9)	1.3360 (3)	0.37519 (5)	0.0277 (4)
H18	0.3302	1.3975	0.3778	0.033*
C19	0.37231 (9)	1.1719 (3)	0.35031 (5)	0.0295 (4)
H19	0.3320	1.1231	0.3352	0.035*
C20	0.43214 (8)	1.0787 (3)	0.34742 (5)	0.0269 (4)
H20	0.4325	0.9663	0.3304	0.032*
C21	0.63939 (8)	0.7940 (2)	0.28645 (5)	0.0220 (4)
C22	0.69737 (9)	0.8528 (3)	0.27206 (5)	0.0283 (4)
H22	0.7291	0.9288	0.2897	0.034*
C23	0.70963 (10)	0.8021 (3)	0.23233 (6)	0.0354 (5)
H23	0.7494	0.8442	0.2228	0.042*
C24	0.66437 (10)	0.6911 (3)	0.20661 (6)	0.0346 (5)
H24	0.6727	0.6571	0.1793	0.042*
C25	0.60686 (10)	0.6291 (3)	0.22063 (6)	0.0340 (4)
H25	0.5758	0.5511	0.2030	0.041*
C26	0.59421 (9)	0.6802 (3)	0.26037 (5)	0.0289 (4)
H26	0.5545	0.6373	0.2698	0.035*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0221 (2)	0.0391 (3)	0.0494 (3)	-0.00554 (19)	-0.00987 (19)	0.0090 (2)
O1	0.0368 (7)	0.0258 (7)	0.0208 (6)	0.0004 (5)	0.0067 (5)	0.0050 (5)
N1	0.0284 (8)	0.0227 (8)	0.0231 (7)	-0.0036 (6)	0.0070 (6)	0.0011 (6)

N2	0.0207 (7)	0.0217 (7)	0.0205 (7)	0.0014 (6)	0.0030 (5)	0.0023 (6)
C1	0.0201 (8)	0.0235 (9)	0.0254 (8)	-0.0021 (7)	0.0002 (6)	-0.0009 (7)
C2	0.0303 (10)	0.0243 (10)	0.0293 (9)	0.0004 (7)	-0.0023 (7)	0.0061 (8)
C3	0.0324 (10)	0.0208 (9)	0.0302 (9)	-0.0057 (7)	0.0029 (7)	0.0031 (8)
C4	0.0230 (9)	0.0264 (9)	0.0271 (9)	-0.0069 (7)	0.0005 (7)	-0.0020 (8)
C5	0.0224 (9)	0.0249 (9)	0.0182 (8)	-0.0005 (7)	0.0002 (6)	0.0003 (7)
C6	0.0230 (8)	0.0215 (9)	0.0154 (7)	-0.0013 (6)	0.0042 (6)	-0.0005 (7)
C7	0.0275 (9)	0.0259 (10)	0.0222 (8)	-0.0054 (7)	0.0000 (7)	0.0034 (7)
C8	0.0211 (9)	0.0205 (9)	0.0264 (9)	-0.0034 (7)	0.0036 (7)	0.0053 (7)
C9	0.0264 (9)	0.0273 (10)	0.0256 (9)	0.0020 (7)	0.0088 (7)	0.0045 (7)
C10	0.0217 (9)	0.0226 (9)	0.0211 (8)	-0.0029 (7)	0.0017 (6)	-0.0015 (7)
C11	0.0178 (8)	0.0231 (9)	0.0230 (8)	-0.0016 (6)	0.0020 (6)	0.0004 (7)
C12	0.0191 (8)	0.0241 (9)	0.0238 (8)	-0.0010 (7)	0.0033 (6)	0.0025 (7)
C13	0.0287 (10)	0.0310 (10)	0.0266 (9)	0.0065 (8)	0.0058 (7)	0.0053 (8)
C14	0.0345 (10)	0.0281 (10)	0.0244 (9)	0.0041 (8)	0.0057 (7)	0.0051 (8)
C15	0.0237 (9)	0.0211 (9)	0.0194 (8)	-0.0012 (7)	0.0041 (6)	0.0016 (7)
C16	0.0292 (9)	0.0235 (9)	0.0187 (8)	-0.0060 (7)	0.0047 (7)	-0.0001 (7)
C17	0.0384 (11)	0.0199 (9)	0.0255 (9)	0.0007 (7)	0.0129 (7)	-0.0006 (7)
C18	0.0287 (10)	0.0285 (10)	0.0275 (9)	0.0063 (7)	0.0092 (7)	0.0026 (8)
C19	0.0229 (9)	0.0339 (10)	0.0309 (9)	-0.0002 (7)	0.0008 (7)	-0.0027 (8)
C20	0.0251 (9)	0.0260 (9)	0.0295 (9)	-0.0011 (7)	0.0031 (7)	-0.0078 (8)
C21	0.0223 (9)	0.0219 (9)	0.0213 (8)	0.0037 (7)	0.0018 (6)	0.0038 (7)
C22	0.0309 (10)	0.0261 (10)	0.0286 (9)	-0.0006 (7)	0.0069 (7)	0.0043 (8)
C23	0.0391 (11)	0.0359 (11)	0.0347 (10)	0.0077 (9)	0.0173 (8)	0.0091 (9)
C24	0.0441 (12)	0.0369 (11)	0.0234 (9)	0.0187 (9)	0.0069 (8)	0.0024 (8)
C25	0.0356 (11)	0.0360 (11)	0.0273 (9)	0.0094 (8)	-0.0058 (8)	-0.0036 (8)
C26	0.0233 (9)	0.0362 (11)	0.0261 (9)	0.0011 (8)	-0.0004 (7)	-0.0018 (8)

Geometric parameters (Å, °)

C11—C1	1.7439 (17)	C12—C13	1.527 (2)
O1—N1	1.4266 (17)	C12—H12	1.0000
O1—C7	1.4280 (19)	C13—H13A	0.9800
N1—C8	1.278 (2)	C13—H13B	0.9800
N2—C14	1.467 (2)	C13—H13C	0.9800
N2—C11	1.479 (2)	C14—H14A	0.9800
N2—C10	1.477 (2)	C14—H14B	0.9800
C1—C2	1.386 (2)	C14—H14C	0.9800
C1—C6	1.391 (2)	C15—C16	1.389 (2)
C2—C3	1.384 (2)	C15—C20	1.393 (2)
C2—H2	0.9500	C16—C17	1.393 (2)
C3—C4	1.382 (2)	C16—H16	0.9500
C3—H3	0.9500	C17—C18	1.377 (3)
C4—C5	1.393 (2)	C17—H17	0.9500
C4—H4	0.9500	C18—C19	1.387 (3)
C5—C6	1.385 (2)	C18—H18	0.9500
C5—H5	0.9500	C19—C20	1.389 (2)
C6—C7	1.506 (2)	C19—H19	0.9500

C7—H7A	0.9900	C20—H20	0.9500
C7—H7B	0.9900	C21—C26	1.391 (2)
C8—C9	1.493 (2)	C21—C22	1.388 (2)
C8—C12	1.504 (2)	C22—C23	1.385 (2)
C9—C10	1.538 (2)	C22—H22	0.9500
C9—H9A	0.9900	C23—C24	1.375 (3)
C9—H9B	0.9900	C23—H23	0.9500
C10—C15	1.515 (2)	C24—C25	1.380 (3)
C10—H10	1.0000	C24—H24	0.9500
C11—C21	1.515 (2)	C25—C26	1.388 (2)
C11—C12	1.543 (2)	C25—H25	0.9500
C11—H11	1.0000	C26—H26	0.9500
N1—O1—C7	107.18 (12)	C13—C12—C11	111.60 (13)
C8—N1—O1	112.01 (13)	C8—C12—H12	107.7
C14—N2—C11	110.05 (12)	C13—C12—H12	107.7
C14—N2—C10	109.23 (13)	C11—C12—H12	107.7
C11—N2—C10	110.66 (12)	C12—C13—H13A	109.5
C2—C1—C6	122.22 (15)	C12—C13—H13B	109.5
C2—C1—C11	118.92 (13)	H13A—C13—H13B	109.5
C6—C1—C11	118.84 (13)	C12—C13—H13C	109.5
C3—C2—C1	118.97 (16)	H13A—C13—H13C	109.5
C3—C2—H2	120.5	H13B—C13—H13C	109.5
C1—C2—H2	120.5	N2—C14—H14A	109.5
C4—C3—C2	120.05 (16)	N2—C14—H14B	109.5
C4—C3—H3	120.0	H14A—C14—H14B	109.5
C2—C3—H3	120.0	N2—C14—H14C	109.5
C3—C4—C5	120.12 (16)	H14A—C14—H14C	109.5
C3—C4—H4	119.9	H14B—C14—H14C	109.5
C5—C4—H4	119.9	C16—C15—C20	118.76 (16)
C6—C5—C4	120.97 (15)	C16—C15—C10	120.51 (15)
C6—C5—H5	119.5	C20—C15—C10	120.56 (15)
C4—C5—H5	119.5	C15—C16—C17	120.54 (16)
C5—C6—C1	117.67 (15)	C15—C16—H16	119.7
C5—C6—C7	122.10 (15)	C17—C16—H16	119.7
C1—C6—C7	120.18 (14)	C18—C17—C16	120.28 (16)
O1—C7—C6	112.89 (13)	C18—C17—H17	119.9
O1—C7—H7A	109.0	C16—C17—H17	119.9
C6—C7—H7A	109.0	C17—C18—C19	119.67 (16)
O1—C7—H7B	109.0	C17—C18—H18	120.2
C6—C7—H7B	109.0	C19—C18—H18	120.2
H7A—C7—H7B	107.8	C18—C19—C20	120.18 (16)
N1—C8—C9	127.36 (15)	C18—C19—H19	119.9
N1—C8—C12	117.26 (15)	C20—C19—H19	119.9
C9—C8—C12	115.38 (14)	C19—C20—C15	120.51 (16)
C8—C9—C10	112.57 (13)	C19—C20—H20	119.7
C8—C9—H9A	109.1	C15—C20—H20	119.7
C10—C9—H9A	109.1	C26—C21—C22	118.62 (15)

C8—C9—H9B	109.1	C26—C21—C11	121.13 (15)
C10—C9—H9B	109.1	C22—C21—C11	120.25 (15)
H9A—C9—H9B	107.8	C23—C22—C21	120.83 (18)
N2—C10—C15	112.02 (13)	C23—C22—H22	119.6
N2—C10—C9	111.17 (13)	C21—C22—H22	119.6
C15—C10—C9	107.56 (13)	C24—C23—C22	120.14 (18)
N2—C10—H10	108.7	C24—C23—H23	119.9
C15—C10—H10	108.7	C22—C23—H23	119.9
C9—C10—H10	108.7	C23—C24—C25	119.79 (17)
N2—C11—C21	110.91 (13)	C23—C24—H24	120.1
N2—C11—C12	111.35 (12)	C25—C24—H24	120.1
C21—C11—C12	110.13 (13)	C24—C25—C26	120.29 (18)
N2—C11—H11	108.1	C24—C25—H25	119.9
C21—C11—H11	108.1	C26—C25—H25	119.9
C12—C11—H11	108.1	C21—C26—C25	120.32 (17)
C8—C12—C13	112.86 (14)	C21—C26—H26	119.8
C8—C12—C11	109.08 (14)	C25—C26—H26	119.8
C7—O1—N1—C8	177.41 (14)	N1—C8—C12—C11	-131.39 (16)
C6—C1—C2—C3	0.0 (3)	C9—C8—C12—C11	49.02 (18)
C11—C1—C2—C3	178.66 (13)	N2—C11—C12—C8	-55.90 (17)
C1—C2—C3—C4	0.4 (3)	C21—C11—C12—C8	-179.37 (13)
C2—C3—C4—C5	-0.3 (3)	N2—C11—C12—C13	178.71 (13)
C3—C4—C5—C6	-0.1 (2)	C21—C11—C12—C13	55.24 (18)
C4—C5—C6—C1	0.4 (2)	N2—C10—C15—C16	-138.80 (15)
C4—C5—C6—C7	-177.07 (15)	C9—C10—C15—C16	98.76 (17)
C2—C1—C6—C5	-0.3 (2)	N2—C10—C15—C20	46.1 (2)
C11—C1—C6—C5	-179.05 (12)	C9—C10—C15—C20	-76.34 (18)
C2—C1—C6—C7	177.20 (16)	C20—C15—C16—C17	2.3 (2)
C11—C1—C6—C7	-1.5 (2)	C10—C15—C16—C17	-172.88 (14)
N1—O1—C7—C6	-87.90 (15)	C15—C16—C17—C18	-0.6 (2)
C5—C6—C7—O1	-9.4 (2)	C16—C17—C18—C19	-1.5 (3)
C1—C6—C7—O1	173.12 (14)	C17—C18—C19—C20	1.9 (3)
O1—N1—C8—C9	-1.6 (2)	C18—C19—C20—C15	-0.1 (3)
O1—N1—C8—C12	178.88 (13)	C16—C15—C20—C19	-2.0 (2)
N1—C8—C9—C10	133.48 (17)	C10—C15—C20—C19	173.19 (15)
C12—C8—C9—C10	-47.0 (2)	N2—C11—C21—C26	-60.8 (2)
C14—N2—C10—C15	60.47 (17)	C12—C11—C21—C26	62.9 (2)
C11—N2—C10—C15	-178.23 (13)	N2—C11—C21—C22	120.20 (16)
C14—N2—C10—C9	-179.16 (13)	C12—C11—C21—C22	-116.07 (17)
C11—N2—C10—C9	-57.86 (17)	C26—C21—C22—C23	1.0 (3)
C8—C9—C10—N2	49.92 (19)	C11—C21—C22—C23	-179.98 (16)
C8—C9—C10—C15	172.88 (14)	C21—C22—C23—C24	-0.4 (3)
C14—N2—C11—C21	-54.15 (17)	C22—C23—C24—C25	-0.4 (3)
C10—N2—C11—C21	-174.97 (13)	C23—C24—C25—C26	0.7 (3)
C14—N2—C11—C12	-177.17 (13)	C22—C21—C26—C25	-0.7 (3)
C10—N2—C11—C12	62.01 (17)	C11—C21—C26—C25	-179.72 (16)
N1—C8—C12—C13	-6.7 (2)	C24—C25—C26—C21	-0.1 (3)

C9—C8—C12—C13 173.67 (14)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 and Cg2 are the centroids of the C1–C6 and C15–C20 rings, respectively.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C17—H17 \cdots Cg1 ⁱ	0.95	2.69	3.556 (2)	151
C3—H3 \cdots Cg2 ⁱⁱ	0.95	2.90	3.6852 (19)	141

Symmetry codes: (i) $x+3/2, y+5/2, z+1$; (ii) $x+3/2, y+3/2, z+1$.