

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Chloroanilinium 2-carboxylate

Min-Min Zhao

College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China

Correspondence e-mail: chenxinyuanseu@yahoo.com.cn


Received 23 May 2012; accepted 5 June 2012

 Key indicators: single-crystal X-ray study; $T = 153$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.088; data-to-parameter ratio = 16.5.

In the title molecular salt, $\text{C}_6\text{H}_7\text{ClN}^+\cdot\text{C}_3\text{H}_3\text{O}_4^-$, the components are linked by $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, leading to a two-dimensional network parallel to the bc plane. Weak $\text{C}-\text{H}\cdots\text{O}$ interactions are also observed.

Related literature

For the structures and properties of related compounds, see: Chen *et al.* (2001); Wang *et al.* (2002); Xue *et al.* (2002); Huang *et al.* (1999); Zhang *et al.* (2001); Ye *et al.* (2008).



Experimental

Crystal data

$\text{C}_6\text{H}_7\text{ClN}^+\cdot\text{C}_3\text{H}_3\text{O}_4^-$
 $M_r = 231.63$
 Monoclinic, $P2_1/c$
 $a = 12.8272$ (19) Å
 $b = 9.2273$ (10) Å
 $c = 8.4114$ (13) Å
 $\beta = 93.809$ (2)°

$V = 993.4$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.38$ mm⁻¹
 $T = 153$ K
 $0.10 \times 0.05 \times 0.05$ mm

Data collection

Rigaku Mercury CCD diffractometer
 Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.910$, $T_{\max} = 1.000$

6935 measured reflections
 2259 independent reflections
 1985 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.088$
 $S = 1.06$
 2259 reflections
 137 parameters

4 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.32$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O4}-\text{H4}\cdots\text{O1}^{\text{i}}$	0.82	1.71	2.5314 (14)	176
$\text{N1}-\text{H1B}\cdots\text{O2}^{\text{ii}}$	0.89	1.87	2.7546 (16)	176
$\text{N1}-\text{H1C}\cdots\text{O4}^{\text{i}}$	0.89	2.25	2.8702 (15)	127
$\text{N1}-\text{H1C}\cdots\text{O2}^{\text{iii}}$	0.89	2.25	2.9313 (15)	133
$\text{N1}-\text{H1A}\cdots\text{O3}$	0.89	1.91	2.7848 (15)	166
$\text{C3}-\text{H3A}\cdots\text{O3}$	0.93	2.55	3.2599 (18)	134

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by a start-up grant from Southeast University, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2414).

References

- Chen, Z.-F., Li, B.-Q., Xie, Y.-R., Xiong, R.-G., You, X.-Z. & Feng, X.-L. (2001). *Inorg. Chem. Commun.* **4**, 346–349.
 Huang, S.-P.-D., Xiong, R.-G., Han, J.-D. & Weiner, B. R. (1999). *Inorg. Chim. Acta* **294**, 95–98.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, L.-Z., Wang, X.-S., Li, Y.-H., Bai, Z.-P., Xiong, R.-G., Xiong, M. & Li, G.-W. (2002). *Chin. J. Inorg. Chem.* **18**, 1191–1194.
 Xue, X., Abrahams, B. F., Xiong, R.-G. & You, X.-Z. (2002). *Aust. J. Chem.* **55**, 495–497.
 Ye, Q., Fu, D.-W., Hang, T., Xiong, R.-G., Chan, P. W. H. & Huang, S.-P.-D. (2008). *Inorg. Chem.* **47**, 772–774.
 Zhang, J., Xiong, R.-G., Chen, X.-T., Che, C.-M., Xue, Z.-L. & You, X.-Z. (2001). *Organometallics* **20**, 4118–4121.

supporting information

Acta Cryst. (2012). E68, o2079 [https://doi.org/10.1107/S1600536812025469]

4-Chloroanilinium 2-carboxyacetate

Min-Min Zhao

S1. Comment

Simple organic salts containing strong intramolecular H-bonds have attracted an attention as materials which display ferroelectric–paraelectric phase transitions (Chen *et al.*, 2001; Huang *et al.*, 1999; Zhang *et al.*, 2001). With the purpose of obtaining phase transition crystals of organic salts, various organic molecules have been studied and a series of new crystal materials have been elaborated (Wang *et al.*, 2002; Xue *et al.*, 2002; Ye *et al.*, 2008). Herewith, we present the synthesis and crystal structure of the title compound, 4-chloroanilinium 2-carboxyacetate.

In the title compound (Fig. 1), the bond lengths and angles have normal values. The asymmetric unit contains one 4-chloroanilinium cation and one 2-carboxyacetate anion. The protonated N atom is involved in strong intramolecular N—H \cdots O hydrogen bonds with the N \cdots O distances of N1—H1A \cdots O3 = 2.7848 (15)Å; N1—H1B \cdots O2 = 2.7546 (16)Å, N1—H1C \cdots O2 = 2.9313 (15)Å and N1—H1C \cdots O4 = 2.8702 (15)Å. The N—H \cdots O and O—H \cdots O H-bonding interactions connected the components into a 2D network parallel to the *bc*-plane. A weak non-classical intermolecular C3—H3A \cdots O3 interaction is observed, Table 1.

S2. Experimental

The malonic acid (10 mmol), 4-chloroaniline (10 mmol) and ethanol (50 mL) were added into a 100 mL flask. The mixture was stirred at 333 K for 2 h, and then the precipitate was filtrated out. Colourless crystals suitable for X-ray diffraction were obtained by slow evaporation of the solution.

S3. Refinement

All the H atoms attached to C atoms were placed into the idealized positions and treated as riding with C—H = 0.93Å (aromatic) and C—H = 0.97Å (methylene) with $U_{iso}(H) = 1.2U_{eq}(C)$. The H atoms based on N and O were placed into the calculated positions with the H—N = 0.89Å and H—O = 0.82Å and refined with $U_{iso}(H) = 1.5U_{eq}(N \text{ and } O)$.




Figure 1

A view of the asymmetric unit with the atomic numbering scheme. The displacement ellipsoids were drawn at the 30% probability level.

4-Chloroanilinium 2-carboxylate

Crystal data

$C_6H_7ClN^+ \cdot C_3H_3O_4^-$

$M_r = 231.63$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 12.8272\ (19)\ \text{\AA}$

$b = 9.2273\ (10)\ \text{\AA}$

$c = 8.4114\ (13)\ \text{\AA}$

$\beta = 93.809\ (2)^\circ$

$V = 993.4\ (2)\ \text{\AA}^3$

$Z = 4$

$F(000) = 480$

$D_x = 1.549\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2259 reflections

$\theta = 2.7\text{--}27.5^\circ$

$\mu = 0.38\ \text{mm}^{-1}$

$T = 153\ \text{K}$

Block, colourless

$0.10 \times 0.05 \times 0.05\ \text{mm}$

Data collection

Rigaku Mercury CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $13.6612\ \text{pixels mm}^{-1}$

CCD profile fitting scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.910$, $T_{\max} = 1.000$

6935 measured reflections

2259 independent reflections

1985 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.7^\circ$

$h = -16 \rightarrow 16$

$k = -9 \rightarrow 11$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.088$

$S = 1.06$

2259 reflections

137 parameters

4 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0471P)^2 + 0.2506P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.58798 (3)	0.24099 (5)	0.04636 (6)	0.03142 (14)
O1	0.05355 (8)	0.92647 (10)	0.25990 (11)	0.0154 (2)
O2	0.12165 (8)	1.07190 (10)	0.45291 (12)	0.0164 (2)
O3	0.16242 (8)	0.58923 (10)	0.41276 (12)	0.0176 (2)
O4	-0.00722 (7)	0.64921 (10)	0.40573 (11)	0.0150 (2)
H4	-0.0212	0.5744	0.3562	0.022*
N1	0.16273 (9)	0.31425 (13)	0.27635 (14)	0.0144 (3)
H1A	0.1611	0.3942	0.3353	0.022*
H1B	0.1469	0.2379	0.3348	0.022*
H1C	0.1165	0.3220	0.1930	0.022*
C9	0.09274 (11)	0.67518 (15)	0.43794 (15)	0.0125 (3)
C7	0.09565 (10)	0.95030 (15)	0.39848 (16)	0.0119 (3)
C4	0.26723 (10)	0.29563 (15)	0.22045 (15)	0.0137 (3)
C1	0.46232 (12)	0.26233 (17)	0.11081 (19)	0.0203 (3)
C8	0.11608 (11)	0.82132 (15)	0.51033 (16)	0.0139 (3)
H8A	0.1889	0.8233	0.5495	0.017*
H8B	0.0743	0.8332	0.6014	0.017*
C6	0.39676 (12)	0.14367 (17)	0.11557 (19)	0.0217 (3)
H6A	0.4186	0.0533	0.0817	0.026*
C5	0.29804 (12)	0.16053 (16)	0.17131 (17)	0.0186 (3)
H5A	0.2532	0.0816	0.1755	0.022*
C2	0.43096 (12)	0.39847 (17)	0.15827 (18)	0.0206 (3)
H2A	0.4755	0.4776	0.1531	0.025*
C3	0.33224 (11)	0.41483 (16)	0.21357 (17)	0.0174 (3)
H3A	0.3098	0.5054	0.2459	0.021*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0211 (2)	0.0287 (3)	0.0460 (3)	0.00583 (15)	0.01430 (18)	0.00397 (18)
O1	0.0212 (5)	0.0124 (5)	0.0121 (5)	0.0001 (4)	-0.0013 (4)	-0.0001 (4)
O2	0.0213 (5)	0.0111 (5)	0.0167 (5)	-0.0022 (4)	0.0002 (4)	-0.0015 (4)

O3	0.0179 (5)	0.0131 (5)	0.0219 (5)	0.0021 (4)	0.0030 (4)	-0.0023 (4)
O4	0.0165 (5)	0.0108 (5)	0.0171 (5)	0.0002 (4)	-0.0019 (4)	-0.0018 (4)
N1	0.0143 (5)	0.0126 (6)	0.0161 (6)	0.0000 (4)	-0.0005 (5)	-0.0010 (4)
C9	0.0177 (6)	0.0113 (7)	0.0086 (6)	0.0002 (5)	0.0012 (5)	0.0032 (5)
C7	0.0105 (6)	0.0126 (7)	0.0129 (6)	0.0013 (5)	0.0028 (5)	-0.0005 (5)
C4	0.0141 (6)	0.0146 (7)	0.0122 (6)	0.0012 (5)	-0.0005 (5)	0.0002 (5)
C1	0.0154 (7)	0.0235 (8)	0.0223 (7)	0.0046 (6)	0.0043 (6)	0.0032 (6)
C8	0.0164 (6)	0.0122 (7)	0.0126 (6)	0.0007 (5)	-0.0011 (5)	-0.0008 (5)
C6	0.0251 (8)	0.0154 (8)	0.0249 (8)	0.0049 (6)	0.0032 (6)	-0.0008 (6)
C5	0.0205 (7)	0.0142 (8)	0.0210 (7)	-0.0005 (5)	0.0000 (6)	-0.0006 (5)
C2	0.0187 (7)	0.0187 (8)	0.0245 (8)	-0.0026 (6)	0.0028 (6)	0.0023 (6)
C3	0.0187 (7)	0.0131 (7)	0.0203 (7)	0.0011 (5)	0.0003 (6)	-0.0016 (5)

Geometric parameters (Å, °)

C11—C1	1.7455 (16)	C4—C5	1.379 (2)
O1—C7	1.2709 (16)	C4—C3	1.384 (2)
O2—C7	1.2488 (17)	C1—C6	1.383 (2)
O3—C9	1.2238 (17)	C1—C2	1.386 (2)
O4—C9	1.3149 (16)	C8—H8A	0.9700
O4—H4	0.8200	C8—H8B	0.9700
N1—C4	1.4598 (17)	C6—C5	1.388 (2)
N1—H1A	0.8900	C6—H6A	0.9300
N1—H1B	0.8900	C5—H5A	0.9300
N1—H1C	0.8900	C2—C3	1.386 (2)
C9—C8	1.5015 (19)	C2—H2A	0.9300
C7—C8	1.5289 (18)	C3—H3A	0.9300
C9—O4—H4	115.8	C9—C8—C7	115.35 (11)
C4—N1—H1A	109.5	C9—C8—H8A	108.4
C4—N1—H1B	109.5	C7—C8—H8A	108.4
H1A—N1—H1B	109.5	C9—C8—H8B	108.4
C4—N1—H1C	109.5	C7—C8—H8B	108.4
H1A—N1—H1C	109.5	H8A—C8—H8B	107.5
H1B—N1—H1C	109.5	C1—C6—C5	119.42 (14)
O3—C9—O4	124.03 (13)	C1—C6—H6A	120.3
O3—C9—C8	121.60 (12)	C5—C6—H6A	120.3
O4—C9—C8	114.37 (11)	C4—C5—C6	119.25 (14)
O2—C7—O1	125.35 (12)	C4—C5—H5A	120.4
O2—C7—C8	116.31 (11)	C6—C5—H5A	120.4
O1—C7—C8	118.34 (12)	C1—C2—C3	118.99 (14)
C5—C4—C3	121.36 (13)	C1—C2—H2A	120.5
C5—C4—N1	119.37 (12)	C3—C2—H2A	120.5
C3—C4—N1	119.25 (12)	C4—C3—C2	119.60 (13)
C6—C1—C2	121.38 (14)	C4—C3—H3A	120.2
C6—C1—C11	119.71 (12)	C2—C3—H3A	120.2
C2—C1—C11	118.91 (12)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O4—H4···O1 ⁱ	0.82	1.71	2.5314 (14)	176
N1—H1B···O2 ⁱⁱ	0.89	1.87	2.7546 (16)	176
N1—H1C···O4 ⁱ	0.89	2.25	2.8702 (15)	127
N1—H1C···O2 ⁱⁱⁱ	0.89	2.25	2.9313 (15)	133
N1—H1A···O3	0.89	1.91	2.7848 (15)	166
C3—H3A···O3	0.93	2.55	3.2599 (18)	134

Symmetry codes: (i) $-x, y-1/2, -z+1/2$; (ii) $x, y-1, z$; (iii) $x, -y+3/2, z-1/2$.