

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[Bis(pyridin-2-yl) selenide- κ^2N,N']tetra-chloridotin(IV)

Gunay Z. Mammadova,^{a*} Zhanna V. Matsulevich,^b
Vladimir K. Osmanov,^b Alexander V. Borisov^b and
Victor N. Khrustalev^c

^aBaku State University, Z. Khalilov St 23, Baku, AZ-1148, Azerbaijan, ^bR.E. Alekseev Nizhny Novgorod State Technical University, 24 Minin St, Nizhny Novgorod, 603950, Russian Federation, and ^cX-Ray Structural Centre, A.N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 28 Vavilov St, B-334, Moscow 119991, Russian Federation
Correspondence e-mail: gunka479@mail.ru


Received 4 June 2012; accepted 7 June 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.025; wR factor = 0.058; data-to-parameter ratio = 25.1.

The title compound, $[SnCl_4(C_{10}H_8N_2Se)]$, was obtained by the reaction of 2,2'-dipyridyl diselenide with tin tetrachloride. The Sn^{IV} ion is coordinated by two N atoms [$Sn-N = 2.266$ (2) and 2.274 (2) Å] from the bis(2-pyridyl)selenide ligand and four chloride anions [$Sn-Cl = 2.3717$ (6)– 2.3939 (6) Å] in a distorted octahedral geometry. The central six-membered chelate ring has a boat conformation with the Se and Sn atoms deviating by 0.692 (3) and 0.855 (3) Å, respectively, from the mean plane through the remaining four ring atoms. The pyridine rings are inclined to each other by a dihedral angle of 49.62 (8)°. The crystal packing exhibits short intermolecular $Se \cdots Cl$ contacts [3.5417 (7) and 3.5648 (7) Å], weak $C-H \cdots Cl$ hydrogen bonds and $\pi-\pi$ stacking interactions between the pyridine rings with a centroid-centroid distance of 3.683 (3) Å.

Related literature

For the crystal structure of the 2,2'-dipyridyl-selenide ligand, see: Dunne *et al.* (1995). For the crystal structures of related compounds, see: Tresoldi *et al.* (1992); Kondo *et al.* (1995); Blake *et al.* (2002); Teles *et al.* (2006); Zhao *et al.* (2007); Wriedt *et al.* (2008a,b,c).



Experimental

Crystal data

$[SnCl_4(C_{10}H_8N_2Se)]$
 $M_r = 495.63$
Monoclinic, $P2_1/c$
 $a = 8.0835$ (4) Å
 $b = 12.2153$ (5) Å
 $c = 14.4710$ (6) Å
 $\beta = 101.208$ (1)°
 $V = 1401.65$ (11) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 5.16$ mm⁻¹
 $T = 100$ K
 $0.30 \times 0.24 \times 0.15$ mm

Data collection

Bruker SMART 1K CCD diffractometer
Absorption correction: multi-scan [*SADABS*; Sheldrick, 1998]
 $T_{min} = 0.306$, $T_{max} = 0.511$
16245 measured reflections
4096 independent reflections
3723 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.058$
 $S = 1.00$
4096 reflections
163 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 1.64$ e Å⁻³
 $\Delta\rho_{min} = -1.10$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C3-H3 \cdots Cl3^i$	0.95	2.79	3.3965 (18)	122
$C8-H8 \cdots Cl2^ii$	0.95	2.83	3.3126 (18)	113

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - 1, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

We thank Professor Abel M. Maharramov for fruitful discussions and help in this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5309).

References

- Blake, A. J., George, M. W., Hubberstey, P., Garcia, R. L. & Wilson, C. (2002). *Acta Cryst.* **E58**, m96–m98.
Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Dunne, S. J., von Nagy-Felsobuki, E. I. & Mackay, M. F. (1995). *Acta Cryst.* **C51**, 1454–1457.
Kondo, M., Kawata, S., Kitagawa, S., Kiso, H. & Munakata, M. (1995). *Acta Cryst.* **C51**, 567–569.
Sheldrick, G. M. (1998). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Teles, W. M., Marinho, M. V., Yoshida, M. I., Speziali, N. L., Krambrock, K., Pinheiro, C. B., Pinhal, N. M., Leitão, A. A. & Machado, F. C. (2006). *Inorg. Chim. Acta*, **359**, 4613–4618.
Tresoldi, G., Rotondo, E., Piraino, P., Lanfranchi, M. & Tiripicchio, A. (1992). *Inorg. Chim. Acta*, **194**, 233–241.
Wriedt, M., Jess, I. & Näther, C. (2008a). *Acta Cryst.* **E64**, m10.
Wriedt, M., Jess, I. & Näther, C. (2008b). *Acta Cryst.* **E64**, m11.
Wriedt, M., Jess, I. & Näther, C. (2008c). *Acta Cryst.* **E64**, m315.
Zhao, Q.-H., Mu, X.-M. & Fang, R.-B. (2007). *Pol. J. Chem.* **81**, 1369–1373.

supporting information

Acta Cryst. (2012). E68, m983 [https://doi.org/10.1107/S160053681202586X]

[Bis(pyridin-2-yl) selenide- κ^2N,N']tetrachloridotin(IV)

Gunay Z. Mammadova, Zhanna V. Matsulevich, Vladimir K. Osmanov, Alexander V. Borisov and Victor N. Khrustalev

S1. Comment

2,2'-Dipyridyl sulfide plays prominent role as useful ligand in coordination chemistry (Tresoldi *et al.*, 1992; Kondo *et al.*, 1995; Blake *et al.*, 2002; Teles *et al.*, 2006; Zhao *et al.*, 2007; Wriedt *et al.*, 2008a, 2008b, 2008c). The important structural feature of these complexes is the practically unchangeable bond angle at sulfur atom. On the other hand, the most labile geometrical parameters in them are the dihedral angle between two pyridine rings as well as the deviation of metal atom from the mean plane of the central six-membered chelate ring passed through the two nitrogen and two carbon atoms due to the different coordination environment. It is interesting to note that 2,2'-dipyridyl selenide is also known (Dunne *et al.*, 1995), however, no structurally characterized metal complexes with this ligand were reported till now.

This article is dedicated to the first structural characterization of metal complex with 2,2'-dipyridyl selenide ligand - [bis(2-pyridyl)selenide- κ^2N,N']tetrachlorido-tin(IV), C₁₀H₈Cl₄N₂SeSn (**I**), which was obtained by the reaction of 2,2'-dipyridyl diselenide with tin tetrachloride (Figure 1).

The molecule of **I** possesses overall intrinsic C_s (*m*) symmetry (Figure 2). The tin ion is coordinated by two N atoms [Sn—N 2.266 (2), 2.274 (2) Å] from bis(2-pyridyl)selenide ligand and four chloride anions [Sn—Cl 2.3717 (6)–2.3939 (6) Å] in a distorted octahedral geometry. The central six-membered chelate ring has a *boat* conformation with the Se and Sn atoms deviating from the mean plane passed through the rest four atoms of the ring at 0.692 (3) and 0.855 (3) Å, respectively. Two pyridine rings are inclined to each other with a dihedral angle of 49.62 (8)°. Remarkably, the value of the bond angle at selenium atom in **I** (101.51 (10)°) is almost equal to that in the free 2,2'-dipyridyl selenide ligand (101.9 (2)°) (Dunne *et al.*, 1995).

In the crystal, the molecules of **I** form the chains along the *a* axis by the attractive intermolecular Se1⋯Cl2ⁱ [3.5417 (7) Å] and Se1⋯Cl4ⁱ [3.5648 (7) Å] interactions. The chains are further linked into a three-dimensional framework by weak C—H⋯Cl hydrogen bonds (Table 1) and π ⋯ π stacking interactions between the pyridine rings with a centroid-centroid distance of 3.683 (3) Å. Symmetry code: (i) *x* - 1, *y*, *z*.

S2. Experimental

A solution of SnCl₄ (0.13 g, 0.5 mmol) in CH₂Cl₂ (25 ml) was added to a solution of 2,2'-dipyridyl diselenide (0.16 g, 0.5 mmol) in CH₂Cl₂ (25 ml) with stirring at room temperature. After 10 min, solvent was evaporated *in vacuo*. An attempt to re-crystallization of the solid residue from CH₃CN led to formation of the powder Se which was separated by filtration of hot solution. The filtrate was concentrated *in vacuo*. The solid was re-crystallized from CH₃CN to give **I** as yellow crystals. Yield is 82%. *M.p.* = 541–543 K. ¹H NMR (DMSO-*d*₆, 300 MHz, 302 K): δ = 8.48 (d, 2H, J = 4.8), 7.70 (t, 2H, J = 7.8), 7.55 (d, 2H, J = 7.8), 7.28 (dd, 2H, J = 7.8, J = 4.8). Anal. Calcd for C₁₀H₈Cl₄N₂SeSn: C, 24.23; H, 1.63; N, 5.65. Found: C, 24.14; H, 1.59; N, 5.57.

S3. Refinement

The hydrogen atoms were placed in calculated positions with $C-H = 0.95 \text{ \AA}$ and refined in the riding model with fixed isotropic displacement parameters [$U_{iso}(H) = 1.2U_{eq}(C)$].




Figure 1

Reaction of 2,2'-dipyridyl diselenide with tin tetrachloride.




Figure 2

The molecular structure of **I** showing the atomic numbering and 50% probability displacement ellipsoids.

[Bis(pyridin-2-yl)selenide- κ^2N,N']tetrachloridotin(IV)

Crystal data

$[\text{SnCl}_4(\text{C}_{10}\text{H}_8\text{N}_2\text{Se})]$

$M_r = 495.63$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 8.0835 (4) \text{ \AA}$

$b = 12.2153 (5) \text{ \AA}$

$c = 14.4710 (6) \text{ \AA}$

$\beta = 101.208 (1)^\circ$

$V = 1401.65 (11) \text{ \AA}^3$

$Z = 4$

$F(000) = 936$

$D_x = 2.349 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9615 reflections

$\theta = 2.2-30.0^\circ$

$\mu = 5.16 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Prism, yellow

$0.30 \times 0.24 \times 0.15 \text{ mm}$

Data collection

Bruker SMART 1K CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
[SADABS; Sheldrick, 1998]

$T_{\min} = 0.306$, $T_{\max} = 0.511$

16245 measured reflections

4096 independent reflections

3723 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 30.0^\circ$, $\theta_{\min} = 2.2^\circ$

$h = -11 \rightarrow 11$

$k = -17 \rightarrow 17$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.058$

$S = 1.00$

4096 reflections

163 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.018P)^2 + 5.2P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 1.64 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -1.10 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.381830 (19)	0.755376 (12)	0.120929 (11)	0.01034 (5)
Se1	-0.08948 (3)	0.76213 (2)	0.108418 (18)	0.01534 (6)
Cl1	0.46143 (7)	0.84715 (5)	-0.00939 (4)	0.01648 (11)
Cl2	0.59551 (7)	0.84559 (5)	0.23278 (4)	0.01729 (11)
Cl3	0.27580 (8)	0.67096 (5)	0.24510 (4)	0.01741 (11)
Cl4	0.54619 (7)	0.59645 (5)	0.10454 (4)	0.01745 (11)
N1	0.2061 (3)	0.89921 (17)	0.12452 (14)	0.0129 (4)
N2	0.1674 (3)	0.68416 (17)	0.01238 (14)	0.0131 (4)
C1	0.0371 (3)	0.8955 (2)	0.11795 (17)	0.0145 (4)
C2	-0.0569 (3)	0.9903 (2)	0.12213 (18)	0.0177 (5)
H2	-0.1758	0.9862	0.1167	0.021*
C3	0.0241 (3)	1.0905 (2)	0.13423 (18)	0.0182 (5)
H3	-0.0382	1.1556	0.1387	0.022*
C4	0.1969 (3)	1.0950 (2)	0.13974 (18)	0.0184 (5)
H4	0.2551	1.1629	0.1477	0.022*
C5	0.2831 (3)	0.9981 (2)	0.13332 (17)	0.0156 (4)

H5	0.4011	1.0012	0.1352	0.019*
C6	0.0019 (3)	0.68436 (19)	0.01536 (17)	0.0134 (4)
C7	-0.1159 (3)	0.6267 (2)	-0.04941 (18)	0.0167 (5)
H7	-0.2316	0.6271	-0.0450	0.020*
C8	-0.0625 (3)	0.5688 (2)	-0.12034 (18)	0.0187 (5)
H8	-0.1401	0.5266	-0.1639	0.022*
C9	0.1063 (3)	0.5734 (2)	-0.12678 (17)	0.0176 (5)
H9	0.1451	0.5369	-0.1765	0.021*
C10	0.2175 (3)	0.6317 (2)	-0.06004 (17)	0.0153 (4)
H10	0.3328	0.6351	-0.0651	0.018*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.00923 (7)	0.01100 (8)	0.01039 (8)	0.00012 (5)	0.00091 (5)	-0.00026 (5)
Se1	0.01033 (11)	0.01823 (12)	0.01802 (12)	-0.00126 (8)	0.00412 (9)	-0.00212 (9)
Cl1	0.0171 (3)	0.0176 (3)	0.0157 (3)	-0.0012 (2)	0.0055 (2)	0.0024 (2)
Cl2	0.0152 (2)	0.0166 (3)	0.0178 (3)	-0.0017 (2)	-0.0023 (2)	-0.0027 (2)
Cl3	0.0206 (3)	0.0184 (3)	0.0138 (2)	-0.0021 (2)	0.0049 (2)	0.0021 (2)
Cl4	0.0161 (3)	0.0151 (3)	0.0203 (3)	0.0045 (2)	0.0014 (2)	-0.0015 (2)
N1	0.0133 (9)	0.0124 (9)	0.0131 (9)	0.0010 (7)	0.0025 (7)	0.0002 (7)
N2	0.0120 (9)	0.0135 (9)	0.0132 (9)	0.0004 (7)	0.0012 (7)	-0.0006 (7)
C1	0.0154 (10)	0.0161 (11)	0.0120 (10)	0.0006 (8)	0.0028 (8)	0.0000 (8)
C2	0.0177 (11)	0.0197 (12)	0.0168 (11)	0.0035 (9)	0.0058 (9)	0.0009 (9)
C3	0.0261 (13)	0.0138 (11)	0.0165 (11)	0.0055 (9)	0.0087 (10)	0.0033 (9)
C4	0.0250 (13)	0.0122 (10)	0.0184 (11)	0.0015 (9)	0.0053 (10)	-0.0007 (9)
C5	0.0160 (11)	0.0142 (10)	0.0169 (11)	0.0004 (8)	0.0038 (9)	0.0016 (9)
C6	0.0130 (10)	0.0112 (10)	0.0157 (11)	0.0003 (8)	0.0018 (8)	0.0012 (8)
C7	0.0138 (10)	0.0155 (11)	0.0187 (11)	-0.0013 (8)	-0.0018 (9)	0.0017 (9)
C8	0.0206 (12)	0.0142 (11)	0.0174 (11)	0.0022 (9)	-0.0061 (9)	0.0004 (9)
C9	0.0226 (12)	0.0159 (11)	0.0127 (11)	0.0031 (9)	-0.0007 (9)	-0.0016 (8)
C10	0.0154 (10)	0.0166 (11)	0.0138 (10)	0.0030 (8)	0.0024 (8)	-0.0021 (8)

Geometric parameters (Å, °)

Sn1—N1	2.266 (2)	C2—H2	0.9500
Sn1—N2	2.274 (2)	C3—C4	1.385 (4)
Sn1—Cl3	2.3717 (6)	C3—H3	0.9500
Sn1—Cl1	2.3873 (6)	C4—C5	1.385 (3)
Sn1—Cl4	2.3901 (6)	C4—H4	0.9500
Sn1—Cl2	2.3939 (6)	C5—H5	0.9500
Se1—C6	1.910 (2)	C6—C7	1.391 (3)
Se1—C1	1.914 (2)	C7—C8	1.383 (4)
N1—C1	1.351 (3)	C7—H7	0.9500
N1—C5	1.353 (3)	C8—C9	1.387 (4)
N2—C6	1.347 (3)	C8—H8	0.9500
N2—C10	1.356 (3)	C9—C10	1.382 (3)
C1—C2	1.393 (3)	C9—H9	0.9500

C2—C3	1.383 (4)	C10—H10	0.9500
N1—Sn1—N2	85.14 (7)	C3—C2—H2	120.2
N1—Sn1—Cl3	89.95 (5)	C1—C2—H2	120.2
N2—Sn1—Cl3	91.02 (5)	C2—C3—C4	119.3 (2)
N1—Sn1—Cl1	85.44 (5)	C2—C3—H3	120.4
N2—Sn1—Cl1	85.43 (5)	C4—C3—H3	120.4
Cl3—Sn1—Cl1	174.40 (2)	C3—C4—C5	118.5 (2)
N1—Sn1—Cl4	174.23 (5)	C3—C4—H4	120.7
N2—Sn1—Cl4	89.11 (5)	C5—C4—H4	120.7
Cl3—Sn1—Cl4	90.68 (2)	N1—C5—C4	122.7 (2)
Cl1—Sn1—Cl4	93.59 (2)	N1—C5—H5	118.6
N1—Sn1—Cl2	90.05 (5)	C4—C5—H5	118.6
N2—Sn1—Cl2	174.98 (5)	N2—C6—C7	122.1 (2)
Cl3—Sn1—Cl2	90.42 (2)	N2—C6—Se1	123.11 (18)
Cl1—Sn1—Cl2	92.76 (2)	C7—C6—Se1	114.76 (18)
Cl4—Sn1—Cl2	95.68 (2)	C8—C7—C6	119.2 (2)
C6—Se1—C1	101.51 (10)	C8—C7—H7	120.4
C1—N1—C5	118.5 (2)	C6—C7—H7	120.4
C1—N1—Sn1	127.01 (16)	C7—C8—C9	118.8 (2)
C5—N1—Sn1	114.52 (16)	C7—C8—H8	120.6
C6—N2—C10	118.3 (2)	C9—C8—H8	120.6
C6—N2—Sn1	127.28 (16)	C10—C9—C8	119.2 (2)
C10—N2—Sn1	114.38 (16)	C10—C9—H9	120.4
N1—C1—C2	121.4 (2)	C8—C9—H9	120.4
N1—C1—Se1	123.52 (18)	N2—C10—C9	122.2 (2)
C2—C1—Se1	114.99 (18)	N2—C10—H10	118.9
C3—C2—C1	119.5 (2)	C9—C10—H10	118.9
N2—Sn1—N1—C1	37.1 (2)	C6—Se1—C1—C2	136.94 (19)
Cl3—Sn1—N1—C1	-53.95 (19)	N1—C1—C2—C3	-0.7 (4)
Cl1—Sn1—N1—C1	122.9 (2)	Se1—C1—C2—C3	176.52 (19)
Cl2—Sn1—N1—C1	-144.38 (19)	C1—C2—C3—C4	1.5 (4)
N2—Sn1—N1—C5	-142.22 (17)	C2—C3—C4—C5	-0.3 (4)
Cl3—Sn1—N1—C5	126.76 (17)	C1—N1—C5—C4	2.7 (4)
Cl1—Sn1—N1—C5	-56.43 (16)	Sn1—N1—C5—C4	-177.9 (2)
Cl2—Sn1—N1—C5	36.33 (17)	C3—C4—C5—N1	-1.9 (4)
N1—Sn1—N2—C6	-42.8 (2)	C10—N2—C6—C7	4.4 (3)
Cl3—Sn1—N2—C6	47.1 (2)	Sn1—N2—C6—C7	-172.03 (18)
Cl1—Sn1—N2—C6	-128.6 (2)	C10—N2—C6—Se1	-176.05 (18)
Cl4—Sn1—N2—C6	137.7 (2)	Sn1—N2—C6—Se1	7.5 (3)
N1—Sn1—N2—C10	140.65 (18)	C1—Se1—C6—N2	40.4 (2)
Cl3—Sn1—N2—C10	-129.48 (17)	C1—Se1—C6—C7	-140.06 (19)
Cl1—Sn1—N2—C10	54.85 (17)	N2—C6—C7—C8	-1.2 (4)
Cl4—Sn1—N2—C10	-38.82 (17)	Se1—C6—C7—C8	179.23 (19)
C5—N1—C1—C2	-1.4 (4)	C6—C7—C8—C9	-2.4 (4)
Sn1—N1—C1—C2	179.31 (17)	C7—C8—C9—C10	2.7 (4)
C5—N1—C1—Se1	-178.36 (18)	C6—N2—C10—C9	-4.1 (4)

Sn1—N1—C1—Se1	2.4 (3)	Sn1—N2—C10—C9	172.81 (19)
C6—Se1—C1—N1	-45.9 (2)	C8—C9—C10—N2	0.5 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C3—H3···Cl3 ⁱ	0.95	2.79	3.3965 (18)	122
C8—H8···Cl2 ⁱⁱ	0.95	2.83	3.3126 (18)	113
C8—H8···Cl3 ⁱⁱⁱ	0.95	2.81	3.6870 (19)	154

Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $x-1, -y+3/2, z-1/2$; (iii) $-x, -y+1, -z$.