

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-[2-[(*E*)-(2-Benzoylhydrazin-1-yl-*idene*)methyl]phenoxy]acetic acidHoong-Kun Fun,^{a*} ‡ Tze Shyang Chia,^a Ahmed M. Alafeefy^b and Hatem A. Abdel-Aziz^c

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Pharmaceutical Chemistry, College of Pharmacy, Salman Bin Abdulaziz University, PO Box 173, Alkharj 11942, Saudi Arabia, and ^cDepartment of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, PO Box 2457, Riyadh 11451, Saudi Arabia
Correspondence e-mail: hkfun@usm.my


Received 21 June 2012; accepted 25 June 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.040; wR factor = 0.124; data-to-parameter ratio = 20.3.

In the title compound, $\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_4$, the dihedral angle between the aromatic rings is $12.45(6)^\circ$. The central $\text{C}(=\text{O})-\text{N}-\text{N}=\text{C}$ bridge is roughly planar (r.m.s. deviation = 0.0346 Å) and makes dihedral angles of $13.01(7)$ and $0.56(7)^\circ$ with the attached phenyl and benzene rings, respectively. The acetic acid unit (r.m.s. deviation = 0.0066 Å) is twisted from its attached benzene ring [dihedral angle = $19.48(6)^\circ$]. In the crystal, molecules are linked by $\text{O}-\text{H}\cdots(\text{O},\text{N})$, $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into sheets lying parallel to the bc plane. A weak aromatic $\pi-\pi$ stacking interaction is also observed [centroid-centroid distance = $3.7330(7)$ Å].

Related literature

For background to the biological activity of hydrazones, see: Abdel-Aziz & Mekawey (2009). For a related structure, see: Rassem *et al.* (2012). For further synthetic details, see: Desai *et al.* (2000). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_4$
 $M_r = 298.29$
Monoclinic, $P2_1/c$
 $a = 12.2173(7)$ Å
 $b = 7.8523(5)$ Å
 $c = 15.6025(9)$ Å
 $\beta = 108.577(1)^\circ$

$V = 1418.82(15)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 100$ K
 $0.32 \times 0.21 \times 0.14$ mm

Data collection

Bruker APEX DUO CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.969$, $T_{\max} = 0.986$

14301 measured reflections
4122 independent reflections
3463 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.124$
 $S = 1.07$
4122 reflections
203 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.42$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3}-\text{H1O3}\cdots\text{O1}^i$	0.88	1.99	2.7194 (14)	139
$\text{O3}-\text{H1O3}\cdots\text{N2}^j$	0.88	2.30	3.0464 (12)	142
$\text{N1}-\text{H1N1}\cdots\text{O4}^{ii}$	0.925 (17)	2.009 (16)	2.9131 (13)	165.4 (16)
$\text{C5}-\text{H5A}\cdots\text{O4}^{ii}$	0.93	2.55	3.4058 (16)	153
$\text{C11}-\text{H11A}\cdots\text{O1}^{iii}$	0.93	2.51	3.3791 (13)	155
$\text{C15}-\text{H15A}\cdots\text{O1}^{iv}$	0.97	2.59	3.5434 (15)	167

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 1, y - \frac{1}{2}, -z - \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + 1, -y + 1, -z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *S SAINT* (Bruker, 2009); data reduction: *S SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and TSC thank Universiti Sains Malaysia (USM) for the Research University Grant (1001/PFIZIK/811160). TSC also thanks the Malaysian Government and USM for the award of a research fellowship. The authors thank the Dean-ship of Scientific Research and the Research Center, College of Pharmacy, King Saud University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6868).

References

- Abdel-Aziz, H. A. & Mekawey, A. A. I. (2009). *Eur. J. Med. Chem.* **44**, 4985–4997.
Bruker (2009). *SADABS*, *APEX2* and *S SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
Desai, N. C., Dave, D., Shah, M. D. & Vyas, G. D. (2000). *Indian J. Chem. Sect. B*, **39**, 277–282.

Rassem, H. H., Salhin, A., Bin Salleh, B., Rosli, M. M. & Fun, H.-K. (2012).
Acta Cryst. **E68**, o1832.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o2260–o2261 [https://doi.org/10.1107/S1600536812028735]

2-{2-[(*E*)-(2-Benzoylhydrazin-1-ylidene)methyl]phenoxy}acetic acid**Hoong-Kun Fun, Tze Shyang Chia, Ahmed M. Alafeefy and Hatem A. Abdel-Aziz****S1. Comment**

In continuation to our interest in the chemistry and biological activities of hydrazones (Abdel-Aziz *et al.*, 2009), we now report herein the crystal structure of the title compound.

The asymmetric unit of the title compound is shown in Fig. 1. The C1–C6 and C9–C14 benzene rings are slightly twisted from each other as indicated by the dihedral angle of 12.45 (6)°. The central C7(=O1)—N1—N2=C8 bridge is nearly planar [r.m.s. deviation = 0.0346 Å], extended in a zigzag conformation and makes dihedral angles of 13.01 (7) and 0.56 (7)° with the C1–C6 and C9–C14 benzene rings, respectively. The acetic acid unit [C15/C16/O3/O4; r.m.s. deviation = 0.0066 Å] is twisted from its attached C9–C14 benzene ring with dihedral angle of 19.48 (6)°. Bond lengths and angles are comparable to those in a related structure (Rassem *et al.*, 2012).

In the crystal (Fig. 2), molecules are linked by O3—H1O3···O1, O3—H1O3···N2, N1—H1N1···O4, C5—H5A···O4, C11—H11A···O1 and C15—H15A···O1 hydrogen bonds (Table 1) into two-dimensional networks parallel to *bc* plane. π - π interaction is also observed with *Cg*1···*Cg*2 distance of 3.7330 (7) Å [symmetry operator = 1 - *x*, -*y*, -*z*], where *Cg*1 and *Cg*2 are the centroids of C1–C6 and C9–C14 rings, respectively.

S2. Experimental

The title compound was prepared by heating 2-(2-formylphenoxy)acetic acid with benzohydrazide in absolute ethanol for 4 h (Desai *et al.*, 2000). Colourless blocks were obtained by slow evaporation from EtOH/DMF.

S3. Refinement

The atoms H1O3 and H1N1 were located in a difference fourier map. Atom H1O3 was then fixed at its found location [O3—H1O3 = 0.8843 Å] and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$, whereas the atom H1N1 was refined freely [N1—H1N1 = 0.924 (17) Å]. The remaining H atoms were positioned geometrically [C—H = 0.93 and 0.97 Å] and refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title compound with 50% probability displacement ellipsoids.




Figure 2

The crystal packing of the title compound. The dashed lines represent the hydrogen bonds. For clarity sake, hydrogen atoms not involved in hydrogen bonding have been omitted.

2-{2-[(*E*)-(2-Benzoylhydrazin-1-ylidene)methyl]phenoxy}acetic acid

Crystal data

C₁₆H₁₄N₂O₄ $M_r = 298.29$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 12.2173$ (7) Å $b = 7.8523$ (5) Å $c = 15.6025$ (9) Å $\beta = 108.577$ (1)° $V = 1418.82$ (15) Å³ $Z = 4$ $F(000) = 624$ $D_x = 1.396$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5671 reflections

 $\theta = 2.8$ – 30.1 ° $\mu = 0.10$ mm⁻¹ $T = 100$ K

Block, colourless

 $0.32 \times 0.21 \times 0.14$ mm

Data collection

Bruker APEX DUO CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scansAbsorption correction: multi-scan
(*SADABS*; Bruker, 2009) $T_{\min} = 0.969$, $T_{\max} = 0.986$

14301 measured reflections

4122 independent reflections

3463 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.024$ $\theta_{\text{max}} = 30.1$ °, $\theta_{\text{min}} = 2.8$ ° $h = -17 \rightarrow 16$ $k = -11 \rightarrow 10$ $l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.124$ $S = 1.07$

4122 reflections

203 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0673P)^2 + 0.394P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.42$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.**Refinement.** Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.70132 (7)	0.06520 (12)	0.21655 (5)	0.0260 (2)
O2	0.35596 (7)	0.38508 (10)	-0.16701 (5)	0.01834 (17)

O3	0.48909 (8)	0.40025 (12)	-0.26386 (6)	0.0273 (2)
H1O3	0.5341	0.4088	-0.2982	0.033*
O4	0.35806 (9)	0.53387 (12)	-0.37939 (6)	0.0305 (2)
N1	0.64695 (7)	0.09189 (11)	0.06501 (6)	0.01427 (18)
N2	0.55352 (7)	0.18816 (11)	0.06821 (6)	0.01471 (18)
C1	0.88465 (10)	-0.14596 (14)	0.22331 (7)	0.0202 (2)
H1A	0.8587	-0.1404	0.2732	0.024*
C2	0.98417 (10)	-0.23668 (15)	0.22871 (9)	0.0262 (2)
H2A	1.0239	-0.2935	0.2818	0.031*
C3	1.02450 (10)	-0.24288 (17)	0.15531 (10)	0.0304 (3)
H3A	1.0920	-0.3018	0.1594	0.036*
C4	0.96375 (11)	-0.16065 (17)	0.07552 (10)	0.0299 (3)
H4A	0.9907	-0.1654	0.0261	0.036*
C5	0.86271 (10)	-0.07092 (15)	0.06869 (8)	0.0221 (2)
H5A	0.8220	-0.0170	0.0149	0.027*
C6	0.82339 (9)	-0.06286 (13)	0.14308 (7)	0.0162 (2)
C7	0.71940 (9)	0.03547 (13)	0.14467 (7)	0.01526 (19)
C8	0.49302 (8)	0.25514 (13)	-0.00687 (6)	0.01392 (19)
H8A	0.5128	0.2371	-0.0590	0.017*
C9	0.39275 (8)	0.36004 (13)	-0.01025 (6)	0.01350 (19)
C10	0.36525 (9)	0.40247 (14)	0.06745 (7)	0.0177 (2)
H10A	0.4114	0.3619	0.1233	0.021*
C11	0.27065 (10)	0.50378 (15)	0.06311 (7)	0.0220 (2)
H11A	0.2542	0.5322	0.1156	0.026*
C12	0.20069 (10)	0.56235 (15)	-0.02047 (7)	0.0206 (2)
H12A	0.1360	0.6278	-0.0239	0.025*
C13	0.22630 (9)	0.52432 (14)	-0.09895 (7)	0.0169 (2)
H13A	0.1793	0.5643	-0.1546	0.020*
C14	0.32297 (8)	0.42582 (13)	-0.09354 (6)	0.01383 (19)
C15	0.31286 (9)	0.48856 (14)	-0.24511 (7)	0.0180 (2)
H15A	0.3096	0.6062	-0.2272	0.022*
H15B	0.2352	0.4528	-0.2792	0.022*
C16	0.38970 (10)	0.47431 (14)	-0.30367 (7)	0.0196 (2)
H1N1	0.6582 (14)	0.066 (2)	0.0106 (11)	0.030 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0283 (4)	0.0379 (5)	0.0125 (3)	0.0126 (4)	0.0072 (3)	0.0028 (3)
O2	0.0218 (4)	0.0241 (4)	0.0112 (3)	0.0062 (3)	0.0082 (3)	0.0036 (3)
O3	0.0275 (4)	0.0309 (5)	0.0311 (4)	0.0042 (4)	0.0200 (4)	0.0053 (4)
O4	0.0533 (6)	0.0261 (4)	0.0172 (4)	0.0037 (4)	0.0184 (4)	0.0046 (3)
N1	0.0150 (4)	0.0165 (4)	0.0117 (4)	0.0012 (3)	0.0048 (3)	-0.0007 (3)
N2	0.0143 (4)	0.0158 (4)	0.0141 (4)	0.0005 (3)	0.0046 (3)	-0.0013 (3)
C1	0.0198 (5)	0.0167 (5)	0.0203 (5)	0.0007 (4)	0.0010 (4)	-0.0014 (4)
C2	0.0198 (5)	0.0185 (5)	0.0328 (6)	0.0023 (4)	-0.0023 (4)	-0.0009 (4)
C3	0.0182 (5)	0.0232 (6)	0.0491 (8)	0.0042 (4)	0.0097 (5)	-0.0008 (5)
C4	0.0269 (6)	0.0282 (6)	0.0414 (7)	0.0040 (5)	0.0202 (5)	-0.0003 (5)

C5	0.0216 (5)	0.0227 (5)	0.0247 (5)	0.0018 (4)	0.0111 (4)	0.0008 (4)
C6	0.0143 (4)	0.0145 (4)	0.0187 (5)	-0.0008 (4)	0.0037 (4)	-0.0016 (4)
C7	0.0160 (4)	0.0160 (4)	0.0138 (4)	-0.0006 (4)	0.0047 (3)	-0.0003 (3)
C8	0.0153 (4)	0.0142 (4)	0.0128 (4)	-0.0021 (3)	0.0051 (3)	-0.0007 (3)
C9	0.0147 (4)	0.0133 (4)	0.0127 (4)	-0.0022 (3)	0.0047 (3)	-0.0007 (3)
C10	0.0218 (5)	0.0194 (5)	0.0122 (4)	0.0003 (4)	0.0055 (4)	0.0000 (4)
C11	0.0273 (5)	0.0247 (5)	0.0170 (5)	0.0037 (4)	0.0114 (4)	-0.0021 (4)
C12	0.0206 (5)	0.0225 (5)	0.0214 (5)	0.0044 (4)	0.0103 (4)	-0.0003 (4)
C13	0.0164 (4)	0.0188 (5)	0.0157 (4)	0.0015 (4)	0.0051 (4)	0.0018 (4)
C14	0.0159 (4)	0.0152 (4)	0.0115 (4)	-0.0017 (3)	0.0059 (3)	-0.0007 (3)
C15	0.0187 (5)	0.0234 (5)	0.0120 (4)	0.0017 (4)	0.0050 (4)	0.0040 (4)
C16	0.0289 (6)	0.0170 (5)	0.0160 (4)	-0.0020 (4)	0.0117 (4)	-0.0007 (4)

Geometric parameters (Å, °)

O1—C7	1.2325 (12)	C5—C6	1.3918 (15)
O2—C14	1.3686 (11)	C5—H5A	0.9300
O2—C15	1.4191 (12)	C6—C7	1.4936 (14)
O3—C16	1.3107 (15)	C8—C9	1.4631 (14)
O3—H1O3	0.8843	C8—H8A	0.9300
O4—C16	1.2136 (13)	C9—C10	1.3975 (13)
N1—C7	1.3504 (13)	C9—C14	1.4059 (13)
N1—N2	1.3831 (11)	C10—C11	1.3869 (15)
N1—H1N1	0.924 (17)	C10—H10A	0.9300
N2—C8	1.2821 (13)	C11—C12	1.3901 (16)
C1—C2	1.3884 (16)	C11—H11A	0.9300
C1—C6	1.3985 (15)	C12—C13	1.3895 (14)
C1—H1A	0.9300	C12—H12A	0.9300
C2—C3	1.3843 (19)	C13—C14	1.3916 (14)
C2—H2A	0.9300	C13—H13A	0.9300
C3—C4	1.389 (2)	C15—C16	1.5084 (14)
C3—H3A	0.9300	C15—H15A	0.9700
C4—C5	1.3956 (16)	C15—H15B	0.9700
C4—H4A	0.9300		
C14—O2—C15	117.23 (8)	C9—C8—H8A	120.1
C16—O3—H1O3	109.9	C10—C9—C14	118.14 (9)
C7—N1—N2	116.90 (8)	C10—C9—C8	122.20 (9)
C7—N1—H1N1	121.8 (10)	C14—C9—C8	119.62 (8)
N2—N1—H1N1	121.3 (10)	C11—C10—C9	121.42 (9)
C8—N2—N1	115.68 (8)	C11—C10—H10A	119.3
C2—C1—C6	120.16 (11)	C9—C10—H10A	119.3
C2—C1—H1A	119.9	C10—C11—C12	119.29 (9)
C6—C1—H1A	119.9	C10—C11—H11A	120.4
C3—C2—C1	120.23 (11)	C12—C11—H11A	120.4
C3—C2—H2A	119.9	C13—C12—C11	120.79 (10)
C1—C2—H2A	119.9	C13—C12—H12A	119.6
C2—C3—C4	119.72 (11)	C11—C12—H12A	119.6

C2—C3—H3A	120.1	C12—C13—C14	119.41 (9)
C4—C3—H3A	120.1	C12—C13—H13A	120.3
C3—C4—C5	120.67 (12)	C14—C13—H13A	120.3
C3—C4—H4A	119.7	O2—C14—C13	123.35 (9)
C5—C4—H4A	119.7	O2—C14—C9	115.76 (9)
C6—C5—C4	119.43 (11)	C13—C14—C9	120.88 (9)
C6—C5—H5A	120.3	O2—C15—C16	110.21 (9)
C4—C5—H5A	120.3	O2—C15—H15A	109.6
C5—C6—C1	119.78 (10)	C16—C15—H15A	109.6
C5—C6—C7	123.97 (9)	O2—C15—H15B	109.6
C1—C6—C7	116.22 (9)	C16—C15—H15B	109.6
O1—C7—N1	121.36 (9)	H15A—C15—H15B	108.1
O1—C7—C6	120.86 (9)	O4—C16—O3	126.30 (10)
N1—C7—C6	117.77 (9)	O4—C16—C15	119.63 (11)
N2—C8—C9	119.75 (8)	O3—C16—C15	114.03 (9)
N2—C8—H8A	120.1		
C7—N1—N2—C8	172.93 (9)	N2—C8—C9—C14	-176.62 (9)
C6—C1—C2—C3	1.17 (17)	C14—C9—C10—C11	1.25 (16)
C1—C2—C3—C4	-1.25 (19)	C8—C9—C10—C11	179.21 (10)
C2—C3—C4—C5	0.4 (2)	C9—C10—C11—C12	0.95 (17)
C3—C4—C5—C6	0.59 (19)	C10—C11—C12—C13	-1.72 (18)
C4—C5—C6—C1	-0.67 (17)	C11—C12—C13—C14	0.25 (17)
C4—C5—C6—C7	177.28 (11)	C15—O2—C14—C13	20.08 (14)
C2—C1—C6—C5	-0.20 (16)	C15—O2—C14—C9	-160.56 (9)
C2—C1—C6—C7	-178.31 (10)	C12—C13—C14—O2	-178.65 (10)
N2—N1—C7—O1	1.68 (15)	C12—C13—C14—C9	2.02 (16)
N2—N1—C7—C6	-177.63 (8)	C10—C9—C14—O2	177.88 (9)
C5—C6—C7—O1	-165.87 (11)	C8—C9—C14—O2	-0.13 (14)
C1—C6—C7—O1	12.14 (15)	C10—C9—C14—C13	-2.74 (15)
C5—C6—C7—N1	13.44 (16)	C8—C9—C14—C13	179.25 (9)
C1—C6—C7—N1	-168.54 (9)	C14—O2—C15—C16	157.84 (9)
N1—N2—C8—C9	-179.66 (8)	O2—C15—C16—O4	169.40 (10)
N2—C8—C9—C10	5.46 (15)	O2—C15—C16—O3	-12.74 (13)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H1O3 \cdots O1 ⁱ	0.88	1.99	2.7194 (14)	139
O3—H1O3 \cdots N2 ⁱ	0.88	2.30	3.0464 (12)	142
N1—H1N1 \cdots O4 ⁱⁱ	0.925 (17)	2.009 (16)	2.9131 (13)	165.4 (16)
C5—H5A \cdots O4 ⁱⁱ	0.93	2.55	3.4058 (16)	153
C11—H11A \cdots O1 ⁱⁱⁱ	0.93	2.51	3.3791 (13)	155
C15—H15A \cdots O1 ^{iv}	0.97	2.59	3.5434 (15)	167

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x+1, y-1/2, -z-1/2$; (iii) $-x+1, y+1/2, -z+1/2$; (iv) $-x+1, -y+1, -z$.