

Bis(dimethylformamide- κ O){4,4',6,6'-tetrachloro-2,2-[butane-1,4-diyl(nitrilomethanylylidene)]diphenolato- κ^4 O,N,N',O'}nickel(II)

Hadi Kargar,^{a*} Reza Kia,^{b‡} Amir Adabi Ardakani^a and Muhammad Nawaz Tahir^{c*}

^aDepartment of Chemistry, Payame Noor University, PO Box 19395-3697 Tehran, I. R. of IRAN, ^bDepartment of Chemistry, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^cDepartment of Physics, University of Sargodha, Punjab, Pakistan

Correspondence e-mail: h.kargar@pnu.ac.ir, dmntahir_uos@yahoo.com


Received 14 June 2012; accepted 25 June 2012

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.044; wR factor = 0.123; data-to-parameter ratio = 19.6.

In the title Schiff base complex, $[\text{Ni}(\text{C}_{18}\text{H}_{14}\text{Cl}_4\text{N}_2\text{O}_2)(\text{C}_3\text{H}_7\text{NO})_2]$, the geometry around the Ni^{II} atom is distorted octahedral. It is coordinated by the N_2O_2 donor atoms of the tetradentate Schiff base ligand and the O atoms of two dimethylformamide molecules, which are *cis* to one another. The benzene rings are almost normal to each other [dihedral angle = $88.60(14)^\circ$]. The various intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds make $S(5)$ and $S(6)$ ring motifs. In the crystal, molecules are linked by pairs of weak $\text{C}-\text{H}\cdots\text{Cl}$ interactions, forming inversion dimers.

Related literature

For standard bond lengths, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For background to Schiff base ligands and their complexes, see: Kargar, Kia, Abbasian *et al.* (2012); Kargar *et al.* (2011); Kia *et al.* (2010). For the crystal structure of the ligand, see: Kargar, Kia, Ardakani *et al.* (2012).



Experimental

Crystal data

$[\text{Ni}(\text{C}_{18}\text{H}_{14}\text{Cl}_4\text{N}_2\text{O}_2)(\text{C}_3\text{H}_7\text{NO})_2]$

$M_r = 637.01$

Monoclinic, $P2_1/n$

$a = 9.7392(11)$ Å

$b = 19.165(2)$ Å

$c = 15.0197(14)$ Å

$\beta = 93.236(3)^\circ$

$V = 2799.0(5)$ Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 1.11$ mm⁻¹

$T = 291$ K

$0.36 \times 0.28 \times 0.26$ mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(SADABS; Bruker, 2005)

$T_{\text{min}} = 0.690$, $T_{\text{max}} = 0.761$

23789 measured reflections

6633 independent reflections

4349 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.123$

$S = 1.02$

6633 reflections

338 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.39$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.58$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C9}-\text{H9B}\cdots\text{O4}$	0.97	2.58	3.327 (4)	134
$\text{C11}-\text{H11B}\cdots\text{O4}$	0.97	2.40	3.057 (4)	125
$\text{C19}-\text{H19}\cdots\text{O1}$	0.93	2.25	2.865 (4)	123
$\text{C8}-\text{H8A}\cdots\text{Cl3}^{\dagger}$	0.97	2.86	3.753 (3)	153

Symmetry code: (i) $-x, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HK and AAA thank PNU for financial support. MNT thanks GC University of Sargodha, Pakistan, for the research facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2463).

[‡] Present address: Structural Dynamics of (Bio)Chemical Systems, Max Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Göttingen, Germany.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Kargar, H., Kia, R., Abbasian, S. & Tahir, M. N. (2012). *Acta Cryst. E* **68**, m182.
- Kargar, H., Kia, R., Ardakani, A. A. S. & Tahir, M. N. (2012). *Acta Cryst. E* **68**, o2244–o2245.
- Kargar, H., Kia, R., Pahlavani, E. & Tahir, M. N. (2011). *Acta Cryst. E* **67**, o614.
- Kia, R., Kargar, H., Tahir, M. N. & Kianoosh, F. (2010). *Acta Cryst. E* **66**, o2296.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2012). E68, m997–m998 [https://doi.org/10.1107/S1600536812028681]

Bis(dimethylformamide- κO){4,4',6,6'-tetrachloro-2,2-[butane-1,4-diyl(nitrilomethanylylidene)]diphenolato- $\kappa^4 O,N,N',O'$ }nickel(II)

Hadi Kargar, Reza Kia, Amir Adabi Ardakani and Muhammad Nawaz Tahir

S1. Comment

In continuation of our work on the synthesis and crystal structure analysis of Schiff base ligands and their complexes (Kargar, Kia, Abbasian *et al.*, 2012; Kargar, Kia, Ardakani *et al.*, 2012; Kargar *et al.*, 2011; Kia *et al.*, 2010), we report herein on the synthesise and crystal structure of the title compound.

The asymmetric unit of the title compound, Fig. 1, comprises a Ni^{II} Schiff base complex. The geometry around Ni^{II} is distorted octahedral being coordinated by N₂O₂ donor atoms of the tetradentate ligand, 6,6'-((butane-1,4-diylbis(azanylylidene))bis(methanylylidene)) bis(2,4-dichlorophenol) [Kargar, Kia, Ardakani *et al.*, 2012] and by two oxygen atoms of dimethylformamide molecules that are *cis* to one another. The bond lengths (Allen *et al.*, 1987) and angles are within the normal range. The intramolecular C—H \cdots O hydrogen bonds makes *S*(5) and *S*(6) ring motif (Table 1; Bernstein *et al.*, 1995). The substituted benzene rings [C1–C6 and C13–C18] are almost normal [88.60 (14)°] to each other.


In the crystal structure molecules are linked by pairs of weak C—H \cdots Cl interactions into individual inversion dimers (Table 1 and Fig. 2).

S2. Experimental


The title compound was synthesized by adding 3,5-dichlorosalicylaldehyde-1,4-butylenediimine (1 mmol) to a solution of NiCl₂·6H₂O (1.1 mmol) in ethanol (30 ml). The mixture was refluxed with stirring for 30 min. The resultant solution was filtered. Green prismatic single crystals of the title compound, suitable for *X*-ray structure determination, were obtained by recrystallization from ethanol on slow evaporation of the solvents at room temperature over several days.

S3. Refinement

The H-atoms were included in calculated positions and treated as riding atoms: C–H = 0.93, 0.96 and 0.97 Å for CH, CH₃ and CH₂ H-atoms, respectively, with U_{iso}(H) = k × U_{eq}(parent C-atom), where k = 1.5 for CH₃ H-atoms and = 1.2 for other H-atoms.

**Figure 1**

The molecular structure of the title compound, showing 40% probability displacement ellipsoids and the atomic numbering. Dashed lines show the intramolecular C-H...O interactions (see Table 1 for details).

**Figure 2**

The crystal packing of the title compound viewed along the *a* axis, showing linking of molecules through weak C—H...Cl interactions (dashed lines; see Table 1 for details) into individual inversion dimers. Only the H atoms involved in these interactions are shown.

Bis(dimethylformamide- κ O){4,4',6,6'-tetrachloro-2,2-[butane-1,4-diyl(nitrilomethanylylidene)]diphenolato- κ^4 O,N,N',O'}nickel(II)

Crystal data

[Ni(C₁₈H₁₄Cl₄N₂O₂)(C₃H₇NO)₂]
 $M_r = 637.01$
 Monoclinic, $P2_1/n$
 Hall symbol: -P 2yn
 $a = 9.7392$ (11) Å
 $b = 19.165$ (2) Å
 $c = 15.0197$ (14) Å
 $\beta = 93.236$ (3)°
 $V = 2799.0$ (5) Å³
 $Z = 4$

$F(000) = 1312$
 $D_x = 1.512$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 3422 reflections
 $\theta = 2.8$ – 27.5 °
 $\mu = 1.11$ mm⁻¹
 $T = 291$ K
 Prism, green
 $0.36 \times 0.28 \times 0.26$ mm

Data collection

Bruker SMART APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.690$, $T_{\max} = 0.761$

23789 measured reflections
 6633 independent reflections
 4349 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$
 $\theta_{\max} = 27.9$ °, $\theta_{\min} = 1.7$ °
 $h = -12 \rightarrow 12$
 $k = -25 \rightarrow 25$
 $l = -18 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.123$
 $S = 1.02$
 6633 reflections
 338 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0542P)^2 + 0.7005P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.39$ e Å⁻³
 $\Delta\rho_{\min} = -0.58$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.03675 (3)	0.167782 (17)	0.39862 (2)	0.03835 (12)
Cl1	0.42524 (9)	0.25307 (4)	0.57798 (6)	0.0652 (2)
Cl2	0.34269 (10)	0.05880 (5)	0.83006 (6)	0.0714 (3)

Cl3	0.40497 (8)	0.00038 (5)	0.36167 (6)	0.0626 (2)
Cl4	0.18116 (14)	-0.08010 (6)	0.04365 (7)	0.0929 (4)
O1	0.1739 (2)	0.19835 (9)	0.49423 (13)	0.0440 (5)
O2	0.1643 (2)	0.09054 (10)	0.36689 (14)	0.0477 (5)
O3	0.1363 (2)	0.23648 (11)	0.31056 (15)	0.0545 (5)
O4	-0.0881 (2)	0.25534 (10)	0.43674 (16)	0.0536 (5)
N1	-0.0578 (2)	0.10534 (11)	0.48708 (16)	0.0393 (5)
N2	-0.0946 (2)	0.14157 (11)	0.29273 (16)	0.0413 (5)
N3	0.3381 (3)	0.28212 (13)	0.27093 (18)	0.0524 (6)
N4	-0.1082 (3)	0.34747 (13)	0.5284 (2)	0.0599 (7)
C1	0.2073 (3)	0.16609 (13)	0.56753 (19)	0.0375 (6)
C2	0.3275 (3)	0.18564 (14)	0.6196 (2)	0.0438 (7)
C3	0.3696 (3)	0.15504 (15)	0.6988 (2)	0.0507 (7)
H3	0.4488	0.1700	0.7307	0.061*
C4	0.2906 (3)	0.10097 (15)	0.7302 (2)	0.0477 (7)
C5	0.1725 (3)	0.08039 (14)	0.68435 (19)	0.0440 (7)
H5	0.1200	0.0449	0.7074	0.053*
C6	0.1289 (3)	0.11139 (13)	0.60378 (19)	0.0395 (6)
C7	0.0006 (3)	0.08645 (13)	0.5616 (2)	0.0425 (7)
H7	-0.0451	0.0524	0.5927	0.051*
C8	-0.1887 (3)	0.07074 (15)	0.4602 (2)	0.0499 (7)
H8A	-0.2166	0.0421	0.5093	0.060*
H8B	-0.1741	0.0400	0.4103	0.060*
C9	-0.3051 (3)	0.12120 (16)	0.4337 (2)	0.0539 (8)
H9A	-0.3881	0.1052	0.4601	0.065*
H9B	-0.2824	0.1666	0.4589	0.065*
C10	-0.3356 (3)	0.12976 (16)	0.3339 (2)	0.0535 (8)
H10A	-0.3411	0.0837	0.3071	0.064*
H10B	-0.4254	0.1513	0.3245	0.064*
C11	-0.2327 (3)	0.17269 (15)	0.2847 (2)	0.0485 (7)
H11A	-0.2630	0.1758	0.2222	0.058*
H11B	-0.2290	0.2196	0.3090	0.058*
C12	-0.0633 (3)	0.10065 (14)	0.22963 (19)	0.0416 (6)
H12	-0.1270	0.0977	0.1813	0.050*
C13	0.0590 (3)	0.05857 (14)	0.22437 (19)	0.0424 (7)
C14	0.0659 (3)	0.01658 (16)	0.1478 (2)	0.0510 (7)
H14	-0.0042	0.0189	0.1032	0.061*
C15	0.1742 (4)	-0.02746 (16)	0.1383 (2)	0.0556 (8)
C16	0.2789 (3)	-0.03246 (15)	0.2039 (2)	0.0540 (8)
H16	0.3522	-0.0628	0.1972	0.065*
C17	0.2736 (3)	0.00781 (14)	0.2790 (2)	0.0447 (7)
C18	0.1638 (3)	0.05554 (14)	0.2943 (2)	0.0411 (6)
C19	0.2586 (3)	0.24959 (15)	0.3267 (2)	0.0493 (7)
H19	0.2982	0.2358	0.3817	0.059*
C20	0.2846 (4)	0.30430 (19)	0.1833 (2)	0.0675 (10)
H20A	0.1949	0.2845	0.1712	0.101*
H20B	0.3450	0.2887	0.1391	0.101*
H20C	0.2783	0.3543	0.1818	0.101*

C21	0.4789 (4)	0.3010 (2)	0.2978 (3)	0.0774 (11)
H21A	0.5021	0.2828	0.3563	0.116*
H21B	0.4875	0.3509	0.2987	0.116*
H21C	0.5400	0.2818	0.2562	0.116*
C22	-0.0370 (3)	0.30254 (15)	0.4834 (2)	0.0518 (8)
H22	0.0583	0.3067	0.4872	0.062*
C23	-0.2567 (4)	0.3464 (2)	0.5240 (3)	0.0882 (14)
H23A	-0.2892	0.3089	0.4859	0.132*
H23B	-0.2912	0.3899	0.5003	0.132*
H23C	-0.2885	0.3396	0.5827	0.132*
C24	-0.0397 (5)	0.4028 (2)	0.5811 (4)	0.1100 (18)
H24A	-0.0736	0.4474	0.5606	0.165*
H24B	0.0577	0.4005	0.5746	0.165*
H24C	-0.0583	0.3970	0.6428	0.165*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.03646 (19)	0.03847 (19)	0.0406 (2)	-0.00509 (14)	0.00601 (15)	0.00000 (15)
Cl1	0.0587 (5)	0.0703 (5)	0.0656 (6)	-0.0261 (4)	-0.0058 (4)	0.0115 (4)
Cl2	0.0795 (6)	0.0831 (6)	0.0507 (5)	0.0078 (5)	-0.0038 (4)	0.0170 (4)
Cl3	0.0420 (4)	0.0766 (6)	0.0693 (6)	0.0060 (4)	0.0048 (4)	0.0036 (4)
Cl4	0.1229 (9)	0.0960 (8)	0.0603 (6)	0.0309 (7)	0.0097 (6)	-0.0273 (5)
O1	0.0462 (11)	0.0416 (10)	0.0439 (12)	-0.0094 (8)	0.0011 (9)	0.0038 (9)
O2	0.0433 (11)	0.0566 (12)	0.0432 (12)	0.0064 (9)	0.0019 (9)	-0.0071 (10)
O3	0.0435 (12)	0.0663 (13)	0.0537 (14)	-0.0126 (10)	0.0046 (10)	0.0147 (11)
O4	0.0486 (12)	0.0433 (11)	0.0691 (15)	-0.0007 (9)	0.0056 (11)	-0.0119 (10)
N1	0.0391 (12)	0.0368 (11)	0.0426 (15)	-0.0064 (9)	0.0079 (10)	-0.0031 (10)
N2	0.0363 (12)	0.0409 (12)	0.0466 (15)	-0.0010 (9)	0.0024 (10)	0.0035 (11)
N3	0.0482 (15)	0.0560 (15)	0.0540 (17)	-0.0036 (12)	0.0129 (12)	0.0131 (12)
N4	0.0649 (18)	0.0504 (15)	0.0648 (19)	0.0059 (13)	0.0066 (15)	-0.0137 (13)
C1	0.0394 (14)	0.0321 (12)	0.0416 (17)	0.0031 (11)	0.0080 (12)	-0.0032 (12)
C2	0.0434 (16)	0.0412 (14)	0.0471 (19)	-0.0034 (12)	0.0054 (13)	-0.0020 (13)
C3	0.0451 (16)	0.0573 (18)	0.049 (2)	0.0048 (14)	-0.0011 (14)	-0.0009 (15)
C4	0.0537 (18)	0.0508 (16)	0.0386 (18)	0.0111 (14)	0.0041 (14)	0.0046 (13)
C5	0.0542 (18)	0.0398 (14)	0.0393 (17)	0.0007 (12)	0.0138 (14)	-0.0016 (12)
C6	0.0453 (15)	0.0350 (13)	0.0394 (17)	0.0007 (11)	0.0123 (12)	-0.0021 (11)
C7	0.0463 (16)	0.0362 (14)	0.0468 (19)	-0.0054 (12)	0.0190 (14)	0.0010 (12)
C8	0.0482 (17)	0.0455 (16)	0.056 (2)	-0.0161 (13)	0.0061 (14)	0.0006 (14)
C9	0.0392 (16)	0.0607 (19)	0.063 (2)	-0.0134 (14)	0.0131 (14)	-0.0137 (15)
C10	0.0354 (15)	0.0563 (18)	0.069 (2)	-0.0025 (13)	0.0039 (14)	-0.0054 (16)
C11	0.0392 (15)	0.0496 (16)	0.056 (2)	0.0017 (12)	-0.0025 (14)	-0.0013 (14)
C12	0.0405 (15)	0.0442 (15)	0.0398 (17)	-0.0041 (12)	-0.0010 (12)	0.0014 (12)
C13	0.0447 (16)	0.0435 (15)	0.0396 (17)	-0.0009 (12)	0.0069 (13)	0.0062 (12)
C14	0.0596 (19)	0.0535 (18)	0.0401 (18)	-0.0003 (14)	0.0052 (14)	0.0011 (14)
C15	0.070 (2)	0.0538 (18)	0.0440 (19)	0.0072 (16)	0.0124 (17)	-0.0045 (15)
C16	0.0560 (19)	0.0482 (16)	0.060 (2)	0.0041 (14)	0.0239 (17)	0.0032 (15)
C17	0.0400 (15)	0.0465 (15)	0.0486 (18)	-0.0013 (12)	0.0109 (13)	0.0057 (13)

C18	0.0384 (14)	0.0417 (14)	0.0443 (18)	-0.0061 (11)	0.0125 (12)	0.0060 (13)
C19	0.0445 (17)	0.0552 (17)	0.0483 (19)	-0.0039 (13)	0.0056 (14)	0.0167 (14)
C20	0.075 (2)	0.075 (2)	0.054 (2)	-0.0091 (19)	0.0167 (19)	0.0089 (18)
C21	0.049 (2)	0.090 (3)	0.095 (3)	-0.0093 (19)	0.020 (2)	0.023 (2)
C22	0.0494 (18)	0.0416 (15)	0.065 (2)	-0.0016 (13)	0.0079 (15)	-0.0036 (15)
C23	0.070 (3)	0.073 (3)	0.126 (4)	0.011 (2)	0.042 (3)	-0.005 (2)
C24	0.119 (4)	0.080 (3)	0.129 (5)	0.011 (3)	-0.007 (3)	-0.056 (3)

Geometric parameters (Å, °)

Ni1—O1	1.993 (2)	C8—H8A	0.9700
Ni1—O2	2.0069 (19)	C8—H8B	0.9700
Ni1—N1	2.046 (2)	C9—C10	1.520 (5)
Ni1—N2	2.046 (2)	C9—H9A	0.9700
Ni1—O3	2.1370 (19)	C9—H9B	0.9700
Ni1—O4	2.1684 (19)	C10—C11	1.520 (4)
C11—C2	1.742 (3)	C10—H10A	0.9700
C12—C4	1.753 (3)	C10—H10B	0.9700
C13—C17	1.738 (3)	C11—H11A	0.9700
C14—C15	1.748 (3)	C11—H11B	0.9700
O1—C1	1.288 (3)	C12—C13	1.444 (4)
O2—C18	1.279 (3)	C12—H12	0.9300
O3—C19	1.229 (3)	C13—C14	1.408 (4)
O4—C22	1.232 (4)	C13—C18	1.425 (4)
N1—C7	1.279 (4)	C14—C15	1.364 (4)
N1—C8	1.473 (3)	C14—H14	0.9300
N2—C12	1.280 (4)	C15—C16	1.381 (5)
N2—C11	1.470 (3)	C16—C17	1.370 (4)
N3—C19	1.328 (4)	C16—H16	0.9300
N3—C20	1.451 (4)	C17—C18	1.436 (4)
N3—C21	1.452 (4)	C19—H19	0.9300
N4—C22	1.315 (4)	C20—H20A	0.9600
N4—C23	1.445 (5)	C20—H20B	0.9600
N4—C24	1.462 (5)	C20—H20C	0.9600
C1—C2	1.421 (4)	C21—H21A	0.9600
C1—C6	1.423 (4)	C21—H21B	0.9600
C2—C3	1.368 (4)	C21—H21C	0.9600
C3—C4	1.389 (4)	C22—H22	0.9300
C3—H3	0.9300	C23—H23A	0.9600
C4—C5	1.365 (4)	C23—H23B	0.9600
C5—C6	1.393 (4)	C23—H23C	0.9600
C5—H5	0.9300	C24—H24A	0.9600
C6—C7	1.450 (4)	C24—H24B	0.9600
C7—H7	0.9300	C24—H24C	0.9600
C8—C9	1.526 (4)		
O1—Ni1—O2	89.39 (8)	C11—C10—C9	116.1 (3)
O1—Ni1—N1	90.68 (9)	C11—C10—H10A	108.3

O2—Ni1—N1	91.73 (9)	C9—C10—H10A	108.3
O1—Ni1—N2	174.92 (8)	C11—C10—H10B	108.3
O2—Ni1—N2	90.14 (9)	C9—C10—H10B	108.3
N1—Ni1—N2	94.39 (9)	H10A—C10—H10B	107.4
O1—Ni1—O3	87.48 (8)	N2—C11—C10	111.5 (2)
O2—Ni1—O3	89.95 (8)	N2—C11—H11A	109.3
N1—Ni1—O3	177.50 (9)	C10—C11—H11A	109.3
N2—Ni1—O3	87.46 (9)	N2—C11—H11B	109.3
O1—Ni1—O4	86.86 (8)	C10—C11—H11B	109.3
O2—Ni1—O4	175.86 (8)	H11A—C11—H11B	108.0
N1—Ni1—O4	90.08 (8)	N2—C12—C13	127.9 (3)
N2—Ni1—O4	93.44 (9)	N2—C12—H12	116.1
O3—Ni1—O4	88.13 (8)	C13—C12—H12	116.1
C1—O1—Ni1	126.97 (17)	C14—C13—C18	120.9 (3)
C18—O2—Ni1	128.06 (19)	C14—C13—C12	116.2 (3)
C19—O3—Ni1	118.2 (2)	C18—C13—C12	122.8 (3)
C22—O4—Ni1	120.2 (2)	C15—C14—C13	120.8 (3)
C7—N1—C8	116.6 (2)	C15—C14—H14	119.6
C7—N1—Ni1	122.67 (18)	C13—C14—H14	119.6
C8—N1—Ni1	120.01 (19)	C14—C15—C16	120.8 (3)
C12—N2—C11	116.3 (3)	C14—C15—C14	120.5 (3)
C12—N2—Ni1	124.1 (2)	C16—C15—C14	118.6 (2)
C11—N2—Ni1	119.52 (19)	C17—C16—C15	119.2 (3)
C19—N3—C20	121.1 (3)	C17—C16—H16	120.4
C19—N3—C21	121.1 (3)	C15—C16—H16	120.4
C20—N3—C21	117.7 (3)	C16—C17—C18	123.7 (3)
C22—N4—C23	121.5 (3)	C16—C17—C13	118.8 (2)
C22—N4—C24	121.1 (3)	C18—C17—C13	117.5 (2)
C23—N4—C24	117.4 (3)	O2—C18—C13	125.1 (3)
O1—C1—C2	120.3 (2)	O2—C18—C17	120.3 (3)
O1—C1—C6	124.4 (3)	C13—C18—C17	114.6 (3)
C2—C1—C6	115.3 (3)	O3—C19—N3	124.4 (3)
C3—C2—C1	124.2 (3)	O3—C19—H19	117.8
C3—C2—C11	119.2 (2)	N3—C19—H19	117.8
C1—C2—C11	116.6 (2)	N3—C20—H20A	109.5
C2—C3—C4	118.1 (3)	N3—C20—H20B	109.5
C2—C3—H3	121.0	H20A—C20—H20B	109.5
C4—C3—H3	121.0	N3—C20—H20C	109.5
C5—C4—C3	120.7 (3)	H20A—C20—H20C	109.5
C5—C4—C12	119.6 (2)	H20B—C20—H20C	109.5
C3—C4—C12	119.7 (3)	N3—C21—H21A	109.5
C4—C5—C6	121.5 (3)	N3—C21—H21B	109.5
C4—C5—H5	119.3	H21A—C21—H21B	109.5
C6—C5—H5	119.3	N3—C21—H21C	109.5
C5—C6—C1	120.2 (3)	H21A—C21—H21C	109.5
C5—C6—C7	116.9 (2)	H21B—C21—H21C	109.5
C1—C6—C7	122.9 (3)	O4—C22—N4	124.4 (3)
N1—C7—C6	128.2 (2)	O4—C22—H22	117.8

N1—C7—H7	115.9	N4—C22—H22	117.8
C6—C7—H7	115.9	N4—C23—H23A	109.5
N1—C8—C9	113.9 (2)	N4—C23—H23B	109.5
N1—C8—H8A	108.8	H23A—C23—H23B	109.5
C9—C8—H8A	108.8	N4—C23—H23C	109.5
N1—C8—H8B	108.8	H23A—C23—H23C	109.5
C9—C8—H8B	108.8	H23B—C23—H23C	109.5
H8A—C8—H8B	107.7	N4—C24—H24A	109.5
C10—C9—C8	115.3 (3)	N4—C24—H24B	109.5
C10—C9—H9A	108.4	H24A—C24—H24B	109.5
C8—C9—H9A	108.4	N4—C24—H24C	109.5
C10—C9—H9B	108.4	H24A—C24—H24C	109.5
C8—C9—H9B	108.4	H24B—C24—H24C	109.5
H9A—C9—H9B	107.5		
O2—Ni1—O1—C1	-69.9 (2)	C4—C5—C6—C7	-178.8 (2)
N1—Ni1—O1—C1	21.8 (2)	O1—C1—C6—C5	-179.0 (2)
O3—Ni1—O1—C1	-159.9 (2)	C2—C1—C6—C5	-0.8 (4)
O4—Ni1—O1—C1	111.9 (2)	O1—C1—C6—C7	-0.6 (4)
O1—Ni1—O2—C18	-161.0 (2)	C2—C1—C6—C7	177.6 (2)
N1—Ni1—O2—C18	108.4 (2)	C8—N1—C7—C6	179.8 (3)
N2—Ni1—O2—C18	14.0 (2)	Ni1—N1—C7—C6	9.7 (4)
O3—Ni1—O2—C18	-73.5 (2)	C5—C6—C7—N1	-177.9 (3)
O1—Ni1—O3—C19	27.0 (2)	C1—C6—C7—N1	3.7 (4)
O2—Ni1—O3—C19	-62.4 (2)	C7—N1—C8—C9	129.0 (3)
N2—Ni1—O3—C19	-152.6 (2)	Ni1—N1—C8—C9	-60.6 (3)
O4—Ni1—O3—C19	113.9 (2)	N1—C8—C9—C10	101.6 (3)
O1—Ni1—O4—C22	15.5 (2)	C8—C9—C10—C11	-74.4 (3)
N1—Ni1—O4—C22	106.2 (2)	C12—N2—C11—C10	93.9 (3)
N2—Ni1—O4—C22	-159.4 (2)	Ni1—N2—C11—C10	-88.3 (3)
O3—Ni1—O4—C22	-72.0 (2)	C9—C10—C11—N2	60.4 (4)
O1—Ni1—N1—C7	-18.0 (2)	C11—N2—C12—C13	-173.9 (3)
O2—Ni1—N1—C7	71.5 (2)	Ni1—N2—C12—C13	8.4 (4)
N2—Ni1—N1—C7	161.7 (2)	N2—C12—C13—C14	178.4 (3)
O4—Ni1—N1—C7	-104.8 (2)	N2—C12—C13—C18	2.3 (4)
O1—Ni1—N1—C8	172.2 (2)	C18—C13—C14—C15	-0.8 (4)
O2—Ni1—N1—C8	-98.3 (2)	C12—C13—C14—C15	-177.0 (3)
N2—Ni1—N1—C8	-8.1 (2)	C13—C14—C15—C16	0.6 (5)
O4—Ni1—N1—C8	85.4 (2)	C13—C14—C15—C14	179.3 (2)
N1—Ni1—N2—C12	-105.1 (2)	C14—C15—C16—C17	-0.4 (5)
O3—Ni1—N2—C12	76.6 (2)	C14—C15—C16—C17	-179.1 (2)
O4—Ni1—N2—C12	164.6 (2)	C15—C16—C17—C18	0.3 (4)
O2—Ni1—N2—C11	169.0 (2)	C15—C16—C17—C13	179.3 (2)
N1—Ni1—N2—C11	77.3 (2)	Ni1—O2—C18—C13	-8.8 (4)
O3—Ni1—N2—C11	-101.0 (2)	Ni1—O2—C18—C17	172.30 (18)
O4—Ni1—N2—C11	-13.1 (2)	C14—C13—C18—O2	-178.4 (3)
Ni1—O1—C1—C2	165.65 (18)	C12—C13—C18—O2	-2.4 (4)
Ni1—O1—C1—C6	-16.2 (4)	C14—C13—C18—C17	0.6 (4)

O1—C1—C2—C3	179.0 (3)	C12—C13—C18—C17	176.6 (2)
C6—C1—C2—C3	0.6 (4)	C16—C17—C18—O2	178.6 (3)
O1—C1—C2—C11	-1.0 (3)	C13—C17—C18—O2	-0.4 (3)
C6—C1—C2—C11	-179.33 (19)	C16—C17—C18—C13	-0.4 (4)
C1—C2—C3—C4	0.6 (4)	C13—C17—C18—C13	-179.41 (19)
C11—C2—C3—C4	-179.4 (2)	Ni1—O3—C19—N3	169.0 (2)
C2—C3—C4—C5	-1.7 (4)	C20—N3—C19—O3	-1.0 (5)
C2—C3—C4—C12	178.5 (2)	C21—N3—C19—O3	174.7 (3)
C3—C4—C5—C6	1.6 (4)	Ni1—O4—C22—N4	-161.8 (3)
C12—C4—C5—C6	-178.6 (2)	C23—N4—C22—O4	-2.2 (5)
C4—C5—C6—C1	-0.3 (4)	C24—N4—C22—O4	-178.9 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C9—H9 <i>B</i> ...O4	0.97	2.58	3.327 (4)	134
C11—H11 <i>B</i> ...O4	0.97	2.40	3.057 (4)	125
C19—H19...O1	0.93	2.25	2.865 (4)	123
C8—H8 <i>A</i> ...C13 ⁱ	0.97	2.86	3.753 (3)	153

Symmetry code: (i) $-x, -y, -z+1$.