

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Redetermination of [Gd(NO₃)₃(H₂O)₄]·2H₂O

Ziyad A. Taha,^{a*} Abdulaziz Ajlouni,^a Ahmed K. Hijazi,^b
Fritz E. Kühn^c and Eberhardt Herdtweck^{d*}

^aDepartment of Applied Chemical Sciences, Jordan University of Science & Technology, Irbid 22110, Jordan, ^bDepartment of Chemistry, Faculty of Science, Jerash University, Jerash 26150, Jordan, ^cDepartment Chemie, Molekulare Katalyse, Technische Universität München, Lichtenbergstrasse 4, D-85747 Garching bei München, Germany, and ^dDepartment Chemie, Lehrstuhl für Anorganische Chemie, Technische Universität München, Lichtenbergstrasse 4, D-85747 Garching bei München, Germany

Correspondence e-mail: akh75_99@yahoo.com, eberhardt.herdtweck@ch.tum.de

Received 18 May 2012; accepted 20 June 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{O}-\text{N}) = 0.004$ Å; R factor = 0.016; wR factor = 0.041; data-to-parameter ratio = 10.2.

The crystal structure of the title compound, tetraaquatris-(nitrato- $\kappa^2\text{O},\text{O}'$)gadolinium(III) dihydrate, was redetermined from single-crystal X-ray data. In comparison with the first determination [Ma *et al.* (1991). *Wuji Huaxue Xuebao*, **7**, 351–353], all H atoms could be located, accompanied with higher accuracy and precision. The Gd^{III} atom shows a ten-coordination with three nitrate ligands behaving in a bidentate manner and the other positions being occupied by four water molecules, forming a distorted bicapped square antiprism. Two nitrate ions coordinate to the metal atom with similar bond lengths while the third shows a more asymmetric bonding behaviour. An intricate network of O—H...O hydrogen bonds, including the lattice water molecules, stabilizes the crystal packing.

Related literature

For a previous determination of the title compound, see: Ma *et al.* (1991). Isotypic [RE(NO₃)₃(H₂O)₄]·2H₂O structures were described for RE = Nd by Rogers *et al.* (1983), for Tb by Moret *et al.* (1990), for Sm by Kawashima *et al.* (2000), for Eu by Stumpf & Bolte (2001), for Dy by Gao *et al.* (1990), and for La by Eriksson *et al.* (1980). [RE(NO₃)₃(H₂O)₄]·H₂O structures with one less water molecule were described for RE = Eu by Ribár *et al.* (1986), for Gd by Stockhause & Meyer (1997), and for Yb by Junk *et al.* (1999).

Experimental

Crystal data

[Gd(NO ₃) ₃ (H ₂ O) ₄]·2H ₂ O	$a = 6.6996$ (2) Å
$M_r = 451.38$	$b = 9.1145$ (3) Å
Triclinic, $P\bar{1}$	$c = 11.6207$ (3) Å

$\alpha = 69.8257$ (10)°
 $\beta = 88.9290$ (11)°
 $\gamma = 69.2170$ (11)°
 $V = 618.36$ (3) Å³
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 5.45$ mm⁻¹
 $T = 173$ K
 $0.48 \times 0.46 \times 0.23$ mm

Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(SADABS; Bruker, 2008)
 $T_{\min} = 0.349$, $T_{\max} = 0.745$

19731 measured reflections
2256 independent reflections
2255 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.016$
 $wR(F^2) = 0.041$
 $S = 1.20$
2256 reflections

221 parameters
All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 1.04$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.10$ e Å⁻³

Table 1

Selected bond lengths (Å).

Gd1—O1	2.528 (2)	Gd1—O8	2.754 (2)
Gd1—O2	2.494 (3)	Gd1—O10	2.398 (2)
Gd1—O4	2.578 (2)	Gd1—O11	2.389 (2)
Gd1—O5	2.518 (3)	Gd1—O12	2.392 (3)
Gd1—O7	2.552 (2)	Gd1—O13	2.364 (3)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O10—H1...O15 ⁱ	0.76 (5)	1.97 (5)	2.720 (4)	171 (5)
O10—H2...O4 ⁱⁱ	0.75 (4)	2.21 (4)	2.956 (3)	178 (6)
O11—H3...O14 ⁱⁱⁱ	0.82 (4)	1.85 (4)	2.671 (3)	176 (5)
O11—H4...O7 ^{iv}	0.78 (5)	2.19 (5)	2.967 (4)	173 (5)
O12—H5...O4 ^v	0.77 (6)	2.50 (6)	3.185 (3)	149 (5)
O12—H5...O7 ^v	0.77 (6)	2.56 (6)	3.156 (4)	135 (5)
O12—H6...O8 ^{vi}	0.83 (5)	2.27 (5)	3.074 (3)	162 (4)
O12—H6...O9 ^{vi}	0.83 (5)	2.39 (5)	3.062 (4)	139 (4)
O13—H7...O15	0.90 (4)	1.84 (4)	2.721 (3)	169 (5)
O13—H8...O14	0.71 (4)	2.04 (4)	2.738 (4)	168 (5)
O14—H9...O9 ^{iv}	0.81 (5)	2.03 (5)	2.826 (4)	167 (4)
O14—H10...O3 ^v	0.76 (5)	2.45 (5)	3.008 (4)	132 (5)
O14—H10...O5 ^{vii}	0.76 (5)	2.31 (5)	2.888 (4)	134 (5)
O15—H11...O6 ⁱⁱ	0.80 (6)	2.02 (6)	2.819 (4)	176 (6)
O15—H12...O3 ^{viii}	0.76 (6)	2.30 (6)	2.903 (4)	139 (5)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y + 1, -z + 1$; (iii) $-x + 1, -y + 1, -z$; (iv) $-x + 2, -y + 1, -z$; (v) $x - 1, y, z$; (vi) $-x + 1, -y + 2, -z$; (vii) $x, y - 1, z$; (viii) $-x + 2, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SIR92 (Altomare *et al.*, 1994); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: PLATON.

The authors are grateful to the Deanship of Scientific Research at Jordan University of Science and Technology for the financial support of this work (grant No. 48/2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2634).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Bruker (2008). *APEX*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Eriksson, B., Larsson, L. O., Niinisto, L. & Valkonen, J. (1980). *Inorg. Chem.* **19**, 1207–1210.
- Gao, S., Ma, H. & Yang, Z. (1990). *J. Northwest Univ. (China)*, **20**, 53–58.
- Junk, P. C., Kepert, D. L., Skelton, B. W. & White, A. H. (1999). *Aust. J. Chem.* **52**, 497–505.
- Kawashima, R., Sasaki, M., Satoh, S., Isoda, H., Kino, Y. & Shiozaki, Y. (2000). *J. Phys. Soc. Jpn.* **69**, 3297–3303.
- Ma, H., Gao, S. & Zupci, Y. (1991). *Wuji Huaxue Xuebao*, **7**, 351–353.
- Moret, E., Bünzli, J.-C. G. & Schenk, K. J. (1990). *Inorg. Chim. Acta*, **178**, 83–88.
- Ribár, B., Kapor, A., Argay, Gy. & Kálmán, A. (1986). *Acta Cryst.* **C42**, 1450–1452.
- Rogers, D. J., Taylor, N. J. & Toogood, G. E. (1983). *Acta Cryst.* **C39**, 939–941.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stockhause, S. & Meyer, G. (1997). *Z. Kristallogr. New Cryst. Struct.* **212**, 315.
- Stumpf, T. & Bolte, M. (2001). *Acta Cryst.* **E57**, i10–i11.

supporting information

Acta Cryst. (2012). E68, i56–i57 [https://doi.org/10.1107/S1600536812028000]

Redetermination of $[\text{Gd}(\text{NO}_3)_3(\text{H}_2\text{O})_4]\cdot 2\text{H}_2\text{O}$

Ziyad A. Taha, Abdulaziz Ajlouni, Ahmed K. Hijazi, Fritz E. Kühn and Eberhardt Herdtweck

S1. Comment

Single crystals of gadolinium nitrate hexahydrate, $[\text{Gd}(\text{NO}_3)_3(\text{H}_2\text{O})_4]2\text{H}_2\text{O}$, were obtained and its structure was re-determined using single-crystal X-ray diffraction analysis.

As shown in Figure 1, the Gd^{III} atom is ten-coordinated, being bound to six nitrate-oxygen atoms, O1, O2, O4, O5, O7, and O8, and four water-oxygen atoms O10, O11, O12, and O13. The nitrate ions are coordinated to the central gadolinium atom in a bidentate mode and show clearly the changes in bond lengths and angles noted previously for the isotopic Eu(III) structure (Stumpf & Bolte, 2001). The distances between the Gd^{III} ion and the nitrate-O atoms ($\text{Gd}-\text{O}$) are in the range 2.494 (3)–2.754 (2) Å, with a mean value of 2.571 Å. The differences between the Gd^{III} atom and the two oxygen atoms of the same nitrate group are 0.034 Å and 0.060 Å for nitrate N1 and nitrate N2, respectively. Nitrate N1 and nitrate N2 groups appear to be more symmetrically bonded to the Gd^{III} atom because the third nitrate group N3 exhibits one $\text{Gd}-\text{O}$ distance that is 0.202 Å longer than the other. This asymmetric bonding seem to be associated with a steric effect of the coordinating water molecules. There is simply not space enough for all the oxygen atoms around the Gd^{III} atom at the same distance. The $\text{Gd}-\text{O}_{\text{water}}$ distances are in the range of 2.364 (3)–2.398 (2) Å with an average distance of 2.386 Å. Hence the water molecules are closer to the Gd^{III} atom than the nitrate groups by *ca.* 0.185 Å.

These results are comparable to that reported for other lanthanide complexes with bidentately coordinating nitrate groups. The coordination polyhedron around the Gd^{III} ion can be best described as a distorted bi-capped square antiprism (Fig. 1), as previously reported for other hydrated lanthanide(III) nitrate complexes (Ribár *et al.*, 1986; Moret *et al.*, 1990; Ma *et al.*, 1991; Stockhause & Meyer, 1997; Kawashima *et al.*, 2000; Stumpf & Bolte, 2001; Junk *et al.*, 1999; Gao *et al.*, 1990; Eriksson *et al.*, 1980; Rogers *et al.*, 1983). The structure contains additional two lattice water molecules, which are associated with the complex by a network of hydrogen bonds. All hydrogen atoms and except for O1 and O2 all oxygen atoms are involved in a complicated network of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, stabilizing the crystal packing (Table 1, Fig. 2). H5, H6 and H10 act as bifurcated bridging atoms. In contrast to the original work of Ma *et al.* (1991), we report much more precise results. All hydrogen atoms could be located in the difference Fourier maps and were allowed to refine freely. The final *R*1 value changed from 0.076 to 0.0158 and the overall e.s.d.'s for the positional parameters dropped from *e.g.* $\text{Gd1 } x = 0.8016$ (1) to 0.80352 (2). The same is valid for distances *e.g.* $\text{Gd1}-\text{O1}$ dropped from 2.558 (56) Å to 2.528 (2) Å and for the angels *e.g.* $\text{O1}-\text{Gd1}-\text{O2}$ dropped from 51.1 (22) to 50.56 (7).

S2. Experimental

Single crystals of $[\text{Gd}(\text{NO}_3)_3(\text{H}_2\text{O})_4]2\text{H}_2\text{O}$ have been obtained accidentally during the synthesis of a $\text{Gd}-N,N$ -bis-(salicylaldehyde)-*o*-phenylenediamine Schiff base complex. 1.0 mmol of the Schiff base were dissolved in 10 ml chloroform. To this solution was added in a drop-wise manner a 10 ml ethyl acetate solution of 2.0 mmol $\text{Gd}(\text{NO}_3)_3\cdot 6(\text{H}_2\text{O})$. The reaction mixture was stirred for 2 h at room temperature. The yellow precipitate was filtered, washed several times with ethyl acetate and chloroform. Single crystals suitable for X-ray were obtained after a few days

by the slow evaporation of the solvent in an open atmosphere.

S3. Refinement

Hydrogen atoms could be located in difference Fourier maps and were allowed to refine freely. The residual electron density of $\Delta\rho_{\max} = 1.04 \text{ e \AA}^{-3}$ is located 0.89 \AA next to Gd1, whereas $\Delta\rho_{\min} = 1.10 \text{ e \AA}^{-3}$ is located 0.96 \AA next to Gd1.




Figure 1

View of the title compound with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level.




Figure 2

Stereo plot of the expanded unit cell. Hydrogen bonds (dotted lines) are limited to a distance $\text{O}\cdots\text{H}$ of 2.35 \AA .

Tetraaquatris(nitrato- κ^2O,O')gadolinium(III) dihydrate

Crystal data

[Gd(NO₃)₃(H₂O)₄] \cdot 2H₂O $M_r = 451.38$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 6.6996$ (2) Å $b = 9.1145$ (3) Å $c = 11.6207$ (3) Å $\alpha = 69.8257$ (10) $^\circ$ $\beta = 88.9290$ (11) $^\circ$ $\gamma = 69.2170$ (11) $^\circ$ $V = 618.36$ (3) Å³ $Z = 2$ $F(000) = 434$ $D_x = 2.424$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9802 reflections

 $\theta = 1.9$ – 25.4 $^\circ$ $\mu = 5.45$ mm⁻¹ $T = 173$ K

Fragment, orange

 $0.48 \times 0.46 \times 0.23$ mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: rotating anode FR591

Graphite monochromator

Detector resolution: 16 pixels mm⁻¹ ϕ - and ω -rotation scans

Absorption correction: multi-scan

(SADABS; Bruker, 2008)

 $T_{\min} = 0.349$, $T_{\max} = 0.745$

19731 measured reflections

2256 independent reflections

2255 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.036$ $\theta_{\max} = 25.4$ $^\circ$, $\theta_{\min} = 1.9$ $^\circ$ $h = -8$ → 8 $k = -10$ → 10 $l = -13$ → 13

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.016$ $wR(F^2) = 0.041$ $S = 1.20$

2256 reflections

221 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: difference Fourier map

All H-atom parameters refined

Weighting scheme based on measured s.u.'s $w =$

$$1/[\sigma^2(F_o^2) + (0.0186P)^2 + 0.725P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

 $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 1.04$ e Å⁻³ $\Delta\rho_{\min} = -1.10$ e Å⁻³

Extinction correction: SHELXL97 (Sheldrick,

2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0129 (7)

Special details

Experimental. Diffractometer operator E. Herdtweck scanspeed 10 s per frame dx 45 4932 frames measured in 9 data sets ϕ -scan with $\Delta\phi = 0.50$ degree-scans with $\Delta\omega = 0.50$ degree

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All s.u.'s are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except for 0 with very negative F^2 or flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $R_{\text{factor_obs}}$ *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Gd1	0.80352 (2)	0.59425 (1)	0.22568 (1)	0.0147 (1)
O1	1.0817 (4)	0.3431 (3)	0.3878 (2)	0.0294 (7)
O2	1.1029 (4)	0.3531 (3)	0.20104 (19)	0.0286 (6)
O3	1.3338 (4)	0.1425 (3)	0.3490 (2)	0.0304 (6)
O4	1.0578 (3)	0.6801 (3)	0.3296 (2)	0.0268 (6)
O5	0.7353 (4)	0.8617 (3)	0.2653 (2)	0.0305 (7)
O6	0.9570 (4)	0.9177 (3)	0.3616 (2)	0.0319 (7)
O7	1.0495 (3)	0.6937 (3)	0.07347 (19)	0.0245 (6)
O8	0.7274 (4)	0.8794 (3)	0.0190 (2)	0.0336 (7)
O9	0.9806 (4)	0.9195 (3)	−0.0904 (2)	0.0322 (7)
O10	0.6976 (4)	0.5590 (3)	0.4287 (2)	0.0224 (7)
O11	0.7203 (4)	0.5521 (3)	0.04330 (19)	0.0223 (6)
O12	0.4279 (4)	0.7621 (3)	0.1754 (2)	0.0259 (7)
O13	0.6339 (4)	0.3963 (3)	0.2792 (2)	0.0262 (7)
N1	1.1767 (4)	0.2751 (3)	0.3141 (2)	0.0213 (7)
N2	0.9171 (4)	0.8234 (3)	0.3202 (2)	0.0233 (8)
N3	0.9183 (4)	0.8342 (3)	−0.0015 (2)	0.0212 (7)
O14	0.5912 (4)	0.2215 (3)	0.1372 (2)	0.0239 (7)
O15	0.6255 (4)	0.2025 (3)	0.5159 (2)	0.0258 (7)
H1	0.608 (7)	0.632 (6)	0.437 (4)	0.038 (12)*
H2	0.760 (6)	0.500 (5)	0.490 (4)	0.028 (11)*
H3	0.625 (6)	0.625 (5)	−0.011 (4)	0.025 (10)*
H4	0.789 (7)	0.491 (6)	0.012 (4)	0.037 (12)*
H5	0.341 (8)	0.724 (6)	0.193 (5)	0.056 (15)*
H6	0.369 (7)	0.865 (6)	0.135 (4)	0.046 (12)*
H7	0.649 (7)	0.328 (6)	0.358 (4)	0.043 (11)*
H8	0.640 (6)	0.344 (5)	0.245 (3)	0.019 (10)*
H9	0.705 (7)	0.183 (5)	0.113 (4)	0.035 (11)*
H10	0.561 (7)	0.151 (6)	0.181 (4)	0.048 (14)*
H11	0.741 (9)	0.167 (6)	0.554 (5)	0.053 (15)*
H12	0.589 (9)	0.131 (7)	0.522 (5)	0.069 (18)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Gd1	0.0156 (1)	0.0131 (1)	0.0141 (1)	−0.0049 (1)	0.0013 (1)	−0.0038 (1)
O1	0.0320 (12)	0.0258 (11)	0.0220 (11)	0.0000 (9)	−0.0012 (9)	−0.0096 (9)
O2	0.0307 (12)	0.0234 (11)	0.0190 (10)	0.0007 (9)	0.0036 (9)	−0.0040 (9)
O3	0.0254 (11)	0.0149 (10)	0.0366 (12)	0.0005 (9)	0.0011 (9)	−0.0006 (9)
O4	0.0246 (11)	0.0262 (11)	0.0298 (11)	−0.0082 (9)	0.0003 (9)	−0.0116 (9)
O5	0.0296 (12)	0.0181 (10)	0.0389 (13)	−0.0076 (9)	−0.0101 (10)	−0.0053 (9)
O6	0.0422 (14)	0.0241 (12)	0.0326 (12)	−0.0157 (11)	−0.0049 (10)	−0.0102 (10)
O7	0.0239 (11)	0.0200 (11)	0.0241 (10)	−0.0073 (9)	0.0020 (8)	−0.0024 (8)
O8	0.0213 (11)	0.0266 (12)	0.0495 (14)	−0.0051 (9)	0.0113 (10)	−0.0140 (11)
O9	0.0416 (13)	0.0205 (11)	0.0261 (11)	−0.0107 (10)	0.0118 (10)	0.0002 (9)

O10	0.0258 (12)	0.0208 (12)	0.0175 (11)	-0.0050 (10)	0.0025 (9)	-0.0072 (10)
O11	0.0260 (11)	0.0204 (11)	0.0162 (10)	-0.0040 (9)	-0.0005 (9)	-0.0061 (9)
O12	0.0182 (11)	0.0195 (12)	0.0354 (12)	-0.0055 (9)	-0.0007 (9)	-0.0058 (10)
O13	0.0445 (14)	0.0267 (12)	0.0178 (11)	-0.0233 (11)	0.0068 (9)	-0.0101 (10)
N1	0.0203 (12)	0.0136 (12)	0.0264 (13)	-0.0063 (10)	0.0027 (10)	-0.0029 (10)
N2	0.0305 (14)	0.0202 (13)	0.0182 (12)	-0.0125 (11)	-0.0018 (10)	-0.0023 (10)
N3	0.0247 (13)	0.0162 (12)	0.0222 (12)	-0.0070 (10)	0.0054 (10)	-0.0070 (10)
O14	0.0277 (12)	0.0189 (11)	0.0232 (11)	-0.0083 (10)	0.0008 (9)	-0.0057 (9)
O15	0.0328 (14)	0.0201 (12)	0.0245 (11)	-0.0118 (10)	0.0039 (10)	-0.0061 (9)

Geometric parameters (Å, °)

Gd1—O1	2.528 (2)	O7—N3	1.277 (3)
Gd1—O2	2.494 (3)	O8—N3	1.243 (4)
Gd1—O4	2.578 (2)	O9—N3	1.227 (3)
Gd1—O5	2.518 (3)	O10—H1	0.76 (5)
Gd1—O7	2.552 (2)	O10—H2	0.75 (4)
Gd1—O8	2.754 (2)	O11—H3	0.82 (4)
Gd1—O10	2.398 (2)	O11—H4	0.78 (5)
Gd1—O11	2.389 (2)	O12—H5	0.77 (6)
Gd1—O12	2.392 (3)	O12—H6	0.83 (5)
Gd1—O13	2.364 (3)	O13—H7	0.90 (4)
O1—N1	1.260 (4)	O13—H8	0.71 (4)
O2—N1	1.268 (3)	O14—H9	0.81 (5)
O3—N1	1.228 (4)	O14—H10	0.76 (5)
O4—N2	1.280 (4)	O15—H11	0.80 (6)
O5—N2	1.259 (4)	O15—H12	0.76 (6)
O6—N2	1.223 (4)		
O1—Gd1—O2	50.56 (7)	O8—Gd1—O12	68.70 (8)
O1—Gd1—O4	67.93 (8)	O8—Gd1—O13	130.50 (8)
O1—Gd1—O5	111.77 (8)	O10—Gd1—O11	140.10 (9)
O1—Gd1—O7	99.76 (8)	O10—Gd1—O12	79.72 (8)
O1—Gd1—O8	146.55 (9)	O10—Gd1—O13	71.04 (9)
O1—Gd1—O10	68.65 (9)	O11—Gd1—O12	78.30 (9)
O1—Gd1—O11	115.74 (9)	O11—Gd1—O13	71.47 (8)
O1—Gd1—O12	143.98 (8)	O12—Gd1—O13	75.61 (9)
O1—Gd1—O13	78.10 (9)	Gd1—O1—N1	95.96 (16)
O2—Gd1—O4	93.36 (8)	Gd1—O2—N1	97.41 (18)
O2—Gd1—O5	141.08 (9)	Gd1—O4—N2	95.25 (17)
O2—Gd1—O7	68.28 (8)	Gd1—O5—N2	98.75 (19)
O2—Gd1—O8	109.78 (8)	Gd1—O7—N3	101.83 (17)
O2—Gd1—O10	117.19 (8)	Gd1—O8—N3	92.94 (17)
O2—Gd1—O11	69.75 (8)	Gd1—O10—H2	129 (3)
O2—Gd1—O12	145.36 (9)	H1—O10—H2	110 (5)
O2—Gd1—O13	81.74 (9)	Gd1—O10—H1	118 (3)
O4—Gd1—O5	49.97 (8)	H3—O11—H4	105 (5)
O4—Gd1—O7	69.84 (7)	Gd1—O11—H3	122 (3)

O4—Gd1—O8	89.68 (8)	Gd1—O11—H4	131 (4)
O4—Gd1—O10	75.39 (8)	Gd1—O12—H6	128 (3)
O4—Gd1—O11	144.41 (8)	Gd1—O12—H5	122 (4)
O4—Gd1—O12	120.88 (9)	H5—O12—H6	109 (5)
O4—Gd1—O13	138.89 (7)	Gd1—O13—H8	122 (3)
O5—Gd1—O7	84.58 (8)	Gd1—O13—H7	120 (3)
O5—Gd1—O8	64.27 (7)	H7—O13—H8	104 (5)
O5—Gd1—O10	70.94 (8)	H9—O14—H10	110 (5)
O5—Gd1—O11	130.51 (8)	H11—O15—H12	110 (6)
O5—Gd1—O12	71.40 (9)	O1—N1—O2	116.1 (3)
O5—Gd1—O13	133.03 (9)	O1—N1—O3	122.2 (2)
O7—Gd1—O8	47.73 (8)	O2—N1—O3	121.7 (3)
O7—Gd1—O10	145.14 (8)	O4—N2—O5	116.0 (3)
O7—Gd1—O11	74.76 (8)	O4—N2—O6	122.0 (3)
O7—Gd1—O12	116.18 (8)	O5—N2—O6	122.0 (3)
O7—Gd1—O13	140.84 (8)	O8—N3—O9	122.0 (3)
O8—Gd1—O10	131.16 (8)	O7—N3—O8	117.5 (2)
O8—Gd1—O11	68.65 (8)	O7—N3—O9	120.5 (3)
O2—Gd1—O1—N1	-0.36 (17)	O11—Gd1—O5—N2	-134.37 (16)
O4—Gd1—O1—N1	-114.9 (2)	O12—Gd1—O5—N2	170.31 (18)
O5—Gd1—O1—N1	-139.49 (18)	O13—Gd1—O5—N2	122.60 (17)
O7—Gd1—O1—N1	-51.5 (2)	O1—Gd1—O7—N3	-171.29 (17)
O8—Gd1—O1—N1	-63.6 (3)	O2—Gd1—O7—N3	148.39 (19)
O10—Gd1—O1—N1	162.8 (2)	O4—Gd1—O7—N3	-109.28 (19)
O11—Gd1—O1—N1	26.2 (2)	O5—Gd1—O7—N3	-60.09 (18)
O12—Gd1—O1—N1	132.50 (19)	O8—Gd1—O7—N3	-0.29 (16)
O13—Gd1—O1—N1	88.7 (2)	O10—Gd1—O7—N3	-104.8 (2)
O1—Gd1—O2—N1	0.36 (17)	O11—Gd1—O7—N3	74.51 (18)
O4—Gd1—O2—N1	57.96 (19)	O12—Gd1—O7—N3	6.1 (2)
O5—Gd1—O2—N1	75.6 (2)	O13—Gd1—O7—N3	105.6 (2)
O7—Gd1—O2—N1	124.7 (2)	O1—Gd1—O8—N3	16.5 (3)
O8—Gd1—O2—N1	148.82 (18)	O2—Gd1—O8—N3	-30.59 (19)
O10—Gd1—O2—N1	-17.3 (2)	O4—Gd1—O8—N3	62.87 (18)
O11—Gd1—O2—N1	-154.2 (2)	O5—Gd1—O8—N3	107.53 (19)
O12—Gd1—O2—N1	-130.31 (19)	O7—Gd1—O8—N3	0.29 (16)
O13—Gd1—O2—N1	-80.99 (19)	O10—Gd1—O8—N3	132.99 (18)
O1—Gd1—O4—N2	-148.15 (17)	O11—Gd1—O8—N3	-88.21 (18)
O2—Gd1—O4—N2	167.14 (15)	O12—Gd1—O8—N3	-173.6 (2)
O5—Gd1—O4—N2	1.56 (14)	O13—Gd1—O8—N3	-126.70 (18)
O7—Gd1—O4—N2	101.75 (16)	Gd1—O1—N1—O2	0.6 (3)
O8—Gd1—O4—N2	57.34 (16)	Gd1—O1—N1—O3	179.5 (3)
O10—Gd1—O4—N2	-75.63 (16)	Gd1—O2—N1—O1	-0.6 (3)
O11—Gd1—O4—N2	108.04 (19)	Gd1—O2—N1—O3	-179.5 (3)
O12—Gd1—O4—N2	-7.40 (18)	Gd1—O4—N2—O5	-2.7 (2)
O13—Gd1—O4—N2	-111.58 (18)	Gd1—O4—N2—O6	177.6 (2)
O1—Gd1—O5—N2	28.63 (19)	Gd1—O5—N2—O4	2.7 (2)
O2—Gd1—O5—N2	-24.9 (2)	Gd1—O5—N2—O6	-177.5 (2)

O4—Gd1—O5—N2	-1.59 (14)	Gd1—O7—N3—O8	0.5 (3)
O7—Gd1—O5—N2	-69.73 (17)	Gd1—O7—N3—O9	-179.1 (2)
O8—Gd1—O5—N2	-114.96 (19)	Gd1—O8—N3—O7	-0.5 (3)
O10—Gd1—O5—N2	85.07 (17)	Gd1—O8—N3—O9	179.1 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O10—H1...O15 ⁱ	0.76 (5)	1.97 (5)	2.720 (4)	171 (5)
O10—H2...O4 ⁱⁱ	0.75 (4)	2.21 (4)	2.956 (3)	178 (6)
O11—H3...O14 ⁱⁱⁱ	0.82 (4)	1.85 (4)	2.671 (3)	176 (5)
O11—H4...O7 ^{iv}	0.78 (5)	2.19 (5)	2.967 (4)	173 (5)
O12—H5...O4 ^v	0.77 (6)	2.50 (6)	3.185 (3)	149 (5)
O12—H5...O7 ^v	0.77 (6)	2.56 (6)	3.156 (4)	135 (5)
O12—H6...O8 ^{vi}	0.83 (5)	2.27 (5)	3.074 (3)	162 (4)
O12—H6...O9 ^{vi}	0.83 (5)	2.39 (5)	3.062 (4)	139 (4)
O13—H7...O15	0.90 (4)	1.84 (4)	2.721 (3)	169 (5)
O13—H8...O14	0.71 (4)	2.04 (4)	2.738 (4)	168 (5)
O14—H9...O9 ^{iv}	0.81 (5)	2.03 (5)	2.826 (4)	167 (4)
O14—H10...O3 ^v	0.76 (5)	2.45 (5)	3.008 (4)	132 (5)
O14—H10...O5 ^{vii}	0.76 (5)	2.31 (5)	2.888 (4)	134 (5)
O15—H11...O6 ⁱⁱ	0.80 (6)	2.02 (6)	2.819 (4)	176 (6)
O15—H12...O3 ^{viii}	0.76 (6)	2.30 (6)	2.903 (4)	139 (5)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+1, -z+1$; (iii) $-x+1, -y+1, -z$; (iv) $-x+2, -y+1, -z$; (v) $x-1, y, z$; (vi) $-x+1, -y+2, -z$; (vii) $x, y-1, z$; (viii) $-x+2, -y, -z+1$.