

Bis(4-benzoyl-3-methyl-1-phenyl-1*H*-pyrazol-5-olato- κ^2O,O')bis(ethanol- κO)-cobalt(II)

 Omoruyi G. Idemudia^{a*} and Eric C. Hosten^b

^aUniversity of Fort Hare, Department of Chemistry, Private Bag X1314, Alice 5700, South Africa, and ^bNelson Mandela Metropolitan University, Department of Chemistry, PO Box 77000, Port Elizabeth 6031, South Africa
Correspondence e-mail: idemudiaog@yahoo.com


Received 27 June 2012; accepted 19 July 2012

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(C-C) = 0.003$ Å; disorder in main residue; R factor = 0.034; wR factor = 0.095; data-to-parameter ratio = 18.8.

The title compound, $[Co(C_{17}H_{13}N_2O_2)_2(C_2H_5OH)_2]$, is a Co^{II} complex with two 4-benzoyl-3-methyl-1-phenyl-1*H*-pyrazol-5-olate (BMPP) ligands and two coordinating ethanol molecules. In the asymmetric unit, there are two half molecules, with the Co^{II} atoms located on inversion centres. The two cobalt complexes have slightly different geometries and in one, the ethyl group of the ethanol is disordered over two sets of sites [occupancy ratio 0.757 (7):0.243 (7)]. Each BMPP ligand is deprotonated with the negative charge delocalized. The hydroxy group of each ethanol molecule forms hydrogen bonds with a pyrazole N atom in an adjacent BMPP ligand. Weaker $C-H\cdots O$ and $C-H\cdots N$ interactions link the molecules into a three-dimensional structure.

Related literature

For related structures, see: Raman *et al.* (2001); Yang *et al.* (2007). For general background and applications of acylpyrazolones, see: Idemudia *et al.* (2012); Marchetti *et al.* (2005); Parihar *et al.* (2012); Zhang *et al.* (2008).



Experimental

Crystal data

$[Co(C_{17}H_{13}N_2O_2)_2(C_2H_6O)_2]$
 $M_r = 705.65$

Triclinic, $P\bar{1}$
 $a = 11.0484$ (3) Å

$b = 11.2282$ (3) Å
 $c = 14.8425$ (4) Å
 $\alpha = 89.205$ (1)°
 $\beta = 87.678$ (1)°
 $\gamma = 76.997$ (1)°
 $V = 1792.56$ (8) Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.53$ mm⁻¹
 $T = 200$ K
 $0.49 \times 0.36 \times 0.12$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: numerical (*SADABS*; Bruker, 2008)
 $T_{min} = 0.84$, $T_{max} = 0.94$

32307 measured reflections
8869 independent reflections
7482 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.095$
 $S = 1.02$
8869 reflections

471 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.51$ e Å⁻³
 $\Delta\rho_{min} = -0.47$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O13—H13A \cdots N22 ⁱ	0.80	2.04	2.8314 (18)	175
O23—H23A \cdots N12 ⁱⁱ	0.89	1.90	2.7862 (16)	177
C16A—H16B \cdots O12	0.99	2.56	3.192 (3)	122
C24—H24C \cdots O11 ⁱⁱⁱ	0.98	2.46	3.361 (2)	154
C112—H112 \cdots O11	0.95	2.24	2.8478 (19)	121
C116—H116 \cdots O23 ^{iv}	0.95	2.60	3.458 (2)	151
C116—H116 \cdots N12	0.95	2.50	2.827 (2)	100
C212—H212 \cdots O21	0.95	2.31	2.848 (2)	115

Symmetry codes: (i) $x - 1, y, z$; (ii) $x + 1, y, z - 1$; (iii) $x + 1, y, z$; (iv) $x - 1, y, z + 1$.

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) and *SHELXL12* (Hübschle *et al.*, 2011); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2009) and *PUBLICIF* (Westrip, 2010).

The Department of Chemistry and the Govan Mbeki Research and Development Centre (GMRDC), both of the University of Fort Hare, are gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GG2088).

References

- Bruker (2008). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2010). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
Idemudia, O. G., Sadimenko, A. P., Afolayan, A. J. & Hosten, E. C. (2012). *Acta Cryst.* **E68**, o1280–o1281.
Marchetti, F., Pettinari, C. & Pettinari, R. (2005). *Coord. Chem. Rev.* **249**, 2909–2945.
Parihar, S., Pathan, S., Jadeja, R. N., Patel, A. & Gupta, V. K. (2012). *Inorg. Chem.* **51**, 1152–1161.
Raman, N., Kulandaisamy, A., Shunmugasundaram, A. & Jeyasubramanian, K. (2001). *Transition Met. Chem.* **26**, 131–135.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

Yang, Y., Zhang, L., Liu, L., Liu, G., Guo, J. & Jia, D. (2007). *Struct. Chem.* **18**, 909–915.

Zhang, L., Liu, L., Xu, G.-C. & Jia, D.-Z. (2008). *J. Chem. Crystallogr.* **38**, 837–843.

supporting information

Acta Cryst. (2012). E68, m1107–m1108 [https://doi.org/10.1107/S1600536812032837]

Bis(4-benzoyl-3-methyl-1-phenyl-1*H*-pyrazol-5-olato- κ^2 O,*O'*)bis(ethanol- κ O)cobalt(II)

Omoruyi G. Idemudia and Eric C. Hosten

S1. Comment

Acylpyrazolones are excellent spectroscopic chelating agents for the determination of metals in trace amounts (Zhang *et al.*, 2008); they also can be used as heterogeneous catalysts (Parihar *et al.*, 2012). As important β -diketones they react with amino groups to form biologically significant Schiff bases (Idemudia *et al.*, 2012). In our probe on transition metal complexes of Schiff bases, an acylpyrazolone Schiff base was treated with cobalt thiocyanate to obtain the title compound (I), instead of the proposed acylpyrazolone Schiff base cobalt complex. The molecular and crystal structure report is presented herein.

The title compound, C₃₈H₃₈CoN₄O₆, is the Co^{II} complex of two 4-benzoyl-3-methyl-1-phenyl-1*H*-pyrazol-5-olate (BMPP) and two ethanol ligands. There are two half molecules of the complex in the asymmetric unit (Figure 1) with Co located at an inversion centres. The ethanol ethyl groups exhibit disorder: in one complex in a ratio of 0.76:0.24 and to a much lesser extent in the other complex molecule (and hence not modelled).

Each BMPP is deprotonated with the negative charge delocalized over O21...C21...C22...C25...O22 and O11...C11...C12...C15...O12. The average Co—O bond length is 2.042 (39) Å and 2.135 (15) Å for the BMPP and ethanol oxygen atoms respectively. The least squares dihedral angles of the BMPP phenyl groups with the pyrazole ring are 16.61 (9) and 61.27 (9)° for one complex and 24.64 (9) and 62.86 (10)° for the other.


The hydrogen of the hydroxy group of each ethanol is hydrogen bonded to a pyrazole nitrogen in an adjacent BMPP (Figure 2). In this way each complex is hydrogen bonded to four adjacent complex molecules. In addition there are inter and intra molecular contacts and C—H...Cg π interactions.

S2. Experimental


A mixture of 4-benzoylphenylhydrazine-3-methyl-1-phenyl-2-pyrazolin-5-one and cobalt thiocyanate in methanol with a molar ratio of 2:1 respectively, was stirred under reflux for 4 h. The complex molecule of C₃₈H₃₈CoN₄O₆ as orange block-like single crystals and with a melting point 219–220°C suitable for X-ray diffraction analysis was obtained from slow evaporation of the final blue solution at room temperature.

S3. Refinement

C-bound H atoms were placed in calculated positions and refined as riding atoms, with C—H 0.95 (CH), 0.99 (CH₂), 0.98 (CH₃) Å and with $U_{iso}(H)=1.2(1.5 \text{ for methyl})U_{eq}(C)$. Hydroxy H atoms were located on a Fourier map and allowed to refine freely.

**Figure 1**

The molecular structure of the title compound (I), with anisotropic displacement ellipsoids drawn at the 50% probability level (for clarity not all atom labels are shown). [Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x + 2, -y + 1, -z$]

**Figure 2**

Hydrogen bonding in (I) (anisotropic displacements ellipsoids drawn at 50% probability level). [Symmetry codes: (i) $-x, -y, -z + 1$; (iii) $x - 1, y, z$; (vii) $-x + 1, -y, -z + 1$]




Figure 3

Crystal packing of (I), viewed along [0 1 0] (anisotropic displacements ellipsoids drawn at 50% probability level).

Bis(4-benzoyl-3-methyl-1-phenyl-1*H*-pyrazol-5-olato- κ^2O,O')bis(ethanol- κO)cobalt(II)

Crystal data

[Co(C₁₇H₁₃N₂O₂)₂(C₂H₆O)₂]

$M_r = 705.65$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 11.0484\ (3)\ \text{\AA}$

$b = 11.2282\ (3)\ \text{\AA}$

$c = 14.8425\ (4)\ \text{\AA}$

$\alpha = 89.205\ (1)^\circ$

$\beta = 87.678\ (1)^\circ$

$\gamma = 76.997\ (1)^\circ$

$V = 1792.56\ (8)\ \text{\AA}^3$

$Z = 2$

$F(000) = 738$

$D_x = 1.307\ \text{Mg m}^{-3}$

Melting point: 219 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 108 reflections

$\theta = 3.8\text{--}31.5^\circ$

$\mu = 0.53\ \text{mm}^{-1}$

$T = 200\ \text{K}$

Block, orange

$0.49 \times 0.36 \times 0.12\ \text{mm}$

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: sealed tube

Graphite monochromator

Detector resolution: $8.3333\ \text{pixels mm}^{-1}$

φ and ω scans

Absorption correction: numerical
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.84$, $T_{\max} = 0.94$

32307 measured reflections

8869 independent reflections

7482 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -14 \rightarrow 14$

$k = -14 \rightarrow 14$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.095$

$S = 1.02$

8869 reflections

471 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0422P)^2 + 0.9543P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.51 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Co1	0	0	0.5	0.02644 (7)	
Co2	1.0	0.5	0	0.03026 (8)	
O11	-0.06104 (10)	0.16165 (9)	0.56233 (6)	0.0315 (2)	
O12	0.11939 (10)	-0.05987 (10)	0.60361 (7)	0.0336 (2)	
O13	0.14538 (12)	0.06673 (12)	0.43001 (9)	0.0466 (3)	
H13A	0.1259	0.1075	0.3864	0.060 (7)*	
O21	1.07029 (12)	0.34177 (11)	0.06276 (7)	0.0394 (3)	
O22	0.87771 (11)	0.54948 (10)	0.11026 (7)	0.0359 (2)	
O23	0.87494 (12)	0.41461 (12)	-0.06579 (8)	0.0462 (3)	
H23A	0.8985	0.369	-0.1143	0.062 (7)*	
N11	-0.10305 (11)	0.27443 (11)	0.69400 (8)	0.0280 (2)	
N12	-0.05984 (13)	0.26925 (12)	0.78148 (8)	0.0341 (3)	
N21	1.11456 (13)	0.22367 (12)	0.19080 (8)	0.0336 (3)	
N22	1.07617 (14)	0.22544 (13)	0.28176 (8)	0.0375 (3)	
C11	-0.03970 (13)	0.17640 (12)	0.64384 (9)	0.0252 (3)	
C12	0.04931 (14)	0.10603 (13)	0.70179 (9)	0.0279 (3)	
C13	0.02963 (15)	0.17038 (14)	0.78577 (9)	0.0335 (3)	
C14	0.0899 (2)	0.1371 (2)	0.87390 (11)	0.0562 (6)	
H14A	0.0953	0.0502	0.8866	0.084*	
H14B	0.0402	0.1863	0.9221	0.084*	
H14C	0.1736	0.1531	0.8708	0.084*	
C15	0.13206 (13)	-0.00484 (13)	0.67443 (9)	0.0275 (3)	
C21	1.05355 (15)	0.32601 (14)	0.14632 (9)	0.0315 (3)	
C22	0.97286 (15)	0.39911 (14)	0.21254 (9)	0.0320 (3)	
C23	0.99378 (16)	0.32862 (15)	0.29435 (10)	0.0362 (3)	

C24	0.9383 (2)	0.35577 (17)	0.38780 (11)	0.0532 (5)
H24A	0.8479	0.3844	0.385	0.08*
H24B	0.9732	0.4192	0.4147	0.08*
H24C	0.9575	0.2814	0.4247	0.08*
C25	0.88599 (15)	0.50747 (14)	0.18943 (9)	0.0318 (3)
C26	0.7822 (3)	0.3650 (3)	−0.01816 (18)	0.0825 (9)
H26A	0.7871	0.2809	−0.0395	0.099*
H26B	0.7993	0.3604	0.0469	0.099*
C27	0.6610 (3)	0.4357 (5)	−0.0300 (4)	0.155 (2)
H27A	0.6598	0.5222	−0.0209	0.232*
H27B	0.6032	0.4099	0.0139	0.232*
H27C	0.6357	0.424	−0.0912	0.232*
C111	−0.19608 (13)	0.37569 (13)	0.66632 (10)	0.0286 (3)
C112	−0.25930 (15)	0.37105 (15)	0.58784 (11)	0.0371 (3)
H112	−0.2418	0.2989	0.5524	0.045*
C113	−0.34801 (17)	0.47218 (18)	0.56156 (13)	0.0481 (4)
H113	−0.39	0.4693	0.5073	0.058*
C114	−0.37636 (19)	0.57684 (18)	0.61289 (15)	0.0535 (5)
H114	−0.4389	0.6449	0.5952	0.064*
C115	−0.3129 (2)	0.58125 (18)	0.68994 (16)	0.0578 (5)
H115	−0.3311	0.6535	0.7252	0.069*
C116	−0.22273 (18)	0.48182 (16)	0.71716 (13)	0.0463 (4)
H116	−0.1793	0.4863	0.7705	0.056*
C121	0.24248 (14)	−0.06123 (14)	0.72766 (9)	0.0309 (3)
C122	0.2658 (2)	−0.18464 (17)	0.74798 (15)	0.0533 (5)
H122	0.2096	−0.2321	0.7307	0.064*
C123	0.3715 (2)	−0.2392 (2)	0.79362 (18)	0.0699 (7)
H123	0.3862	−0.3235	0.8089	0.084*
C124	0.45441 (19)	−0.1725 (2)	0.81670 (14)	0.0581 (5)
H124	0.5263	−0.2104	0.8481	0.07*
C125	0.43434 (17)	−0.0511 (2)	0.79472 (13)	0.0521 (5)
H125	0.4936	−0.0055	0.8094	0.063*
C126	0.32726 (17)	0.00593 (17)	0.75092 (12)	0.0428 (4)
H12A	0.3125	0.0907	0.737	0.051*
C211	1.19389 (14)	0.11779 (14)	0.15297 (10)	0.0318 (3)
C212	1.26245 (15)	0.12585 (17)	0.07298 (10)	0.0382 (3)
H212	1.2579	0.2026	0.044	0.046*
C213	1.33730 (16)	0.02042 (19)	0.03630 (12)	0.0463 (4)
H213	1.3826	0.0252	−0.0189	0.056*
C214	1.34711 (18)	−0.09108 (19)	0.07852 (14)	0.0516 (5)
H214	1.3995	−0.1624	0.053	0.062*
C215	1.28032 (19)	−0.09846 (17)	0.15823 (15)	0.0514 (5)
H215	1.2875	−0.1751	0.1879	0.062*
C216	1.20287 (17)	0.00521 (16)	0.19519 (12)	0.0418 (4)
H216	1.1559	−0.0008	0.2494	0.05*
C221	0.79264 (16)	0.57662 (15)	0.25689 (10)	0.0375 (3)
C222	0.66813 (19)	0.5958 (2)	0.23946 (14)	0.0560 (5)
H222	0.6429	0.5648	0.1858	0.067*

C223	0.5791 (2)	0.6608 (3)	0.30081 (18)	0.0757 (8)	
H223	0.4931	0.6719	0.2901	0.091*	
C224	0.6170 (3)	0.7090 (2)	0.37703 (16)	0.0757 (8)	
H224	0.5565	0.7535	0.4187	0.091*	
C225	0.7401 (3)	0.69350 (19)	0.39334 (14)	0.0635 (6)	
H225	0.765	0.7285	0.4455	0.076*	
C226	0.8290 (2)	0.62677 (16)	0.33398 (11)	0.0459 (4)	
H226	0.9147	0.6151	0.3458	0.055*	
C16A	0.2742 (3)	-0.0041 (4)	0.4279 (2)	0.0533 (9)	0.757 (7)
H16A	0.2836	-0.0705	0.3832	0.064*	0.757 (7)
H16B	0.2927	-0.0424	0.4878	0.064*	0.757 (7)
C17A	0.3656 (3)	0.0719 (3)	0.4045 (2)	0.0640 (10)	0.757 (7)
H17A	0.45	0.0206	0.4048	0.096*	0.757 (7)
H17B	0.3573	0.1374	0.4488	0.096*	0.757 (7)
H17C	0.3496	0.1078	0.3443	0.096*	0.757 (7)
C16B	0.2583 (12)	0.0716 (13)	0.4401 (8)	0.070 (4)	0.243 (7)
H16C	0.2663	0.1027	0.5012	0.084*	0.243 (7)
H16D	0.2814	0.1302	0.3958	0.084*	0.243 (7)
C17B	0.3450 (12)	-0.0480 (17)	0.4284 (10)	0.106 (6)	0.243 (7)
H17D	0.3669	-0.062	0.3642	0.159*	0.243 (7)
H17E	0.3053	-0.1125	0.4519	0.159*	0.243 (7)
H17F	0.4204	-0.0493	0.4614	0.159*	0.243 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.03579 (15)	0.02519 (13)	0.01517 (12)	0.00070 (10)	-0.00458 (10)	-0.00412 (9)
Co2	0.04097 (16)	0.03298 (15)	0.01484 (12)	-0.00448 (12)	0.00239 (10)	-0.00391 (10)
O11	0.0441 (6)	0.0282 (5)	0.0180 (4)	0.0022 (4)	-0.0068 (4)	-0.0049 (4)
O12	0.0445 (6)	0.0306 (5)	0.0216 (5)	0.0021 (4)	-0.0091 (4)	-0.0060 (4)
O13	0.0443 (7)	0.0425 (7)	0.0467 (7)	0.0012 (5)	0.0090 (5)	0.0110 (6)
O21	0.0552 (7)	0.0384 (6)	0.0188 (5)	0.0007 (5)	0.0073 (4)	-0.0016 (4)
O22	0.0456 (6)	0.0379 (6)	0.0209 (5)	-0.0031 (5)	0.0045 (4)	-0.0029 (4)
O23	0.0487 (7)	0.0593 (8)	0.0324 (6)	-0.0162 (6)	0.0078 (5)	-0.0235 (5)
N11	0.0322 (6)	0.0298 (6)	0.0199 (5)	-0.0019 (5)	-0.0034 (4)	-0.0068 (4)
N12	0.0400 (7)	0.0393 (7)	0.0200 (5)	-0.0013 (6)	-0.0062 (5)	-0.0098 (5)
N21	0.0415 (7)	0.0373 (7)	0.0195 (5)	-0.0049 (6)	0.0052 (5)	-0.0018 (5)
N22	0.0508 (8)	0.0387 (7)	0.0205 (6)	-0.0065 (6)	0.0079 (5)	-0.0001 (5)
C11	0.0298 (7)	0.0255 (6)	0.0196 (6)	-0.0045 (5)	-0.0013 (5)	-0.0041 (5)
C12	0.0329 (7)	0.0309 (7)	0.0186 (6)	-0.0038 (6)	-0.0047 (5)	-0.0045 (5)
C13	0.0380 (8)	0.0381 (8)	0.0218 (6)	-0.0018 (6)	-0.0063 (6)	-0.0079 (6)
C14	0.0680 (13)	0.0625 (12)	0.0253 (8)	0.0159 (10)	-0.0182 (8)	-0.0149 (8)
C15	0.0324 (7)	0.0293 (7)	0.0199 (6)	-0.0046 (5)	-0.0042 (5)	-0.0005 (5)
C21	0.0389 (8)	0.0337 (7)	0.0217 (6)	-0.0081 (6)	0.0035 (5)	-0.0028 (5)
C22	0.0426 (8)	0.0335 (7)	0.0200 (6)	-0.0094 (6)	0.0062 (6)	-0.0035 (5)
C23	0.0493 (9)	0.0364 (8)	0.0220 (7)	-0.0093 (7)	0.0072 (6)	-0.0013 (6)
C24	0.0847 (15)	0.0415 (9)	0.0252 (8)	-0.0007 (9)	0.0183 (8)	0.0016 (7)
C25	0.0412 (8)	0.0339 (7)	0.0211 (6)	-0.0108 (6)	0.0052 (6)	-0.0051 (5)

C26	0.0781 (17)	0.118 (2)	0.0662 (15)	-0.0562 (17)	0.0244 (13)	-0.0505 (15)
C27	0.066 (2)	0.212 (6)	0.173 (5)	-0.009 (3)	0.028 (3)	-0.017 (4)
C111	0.0271 (7)	0.0295 (7)	0.0274 (7)	-0.0031 (5)	0.0015 (5)	-0.0030 (5)
C112	0.0369 (8)	0.0387 (8)	0.0316 (7)	0.0007 (7)	-0.0040 (6)	-0.0059 (6)
C113	0.0428 (10)	0.0523 (11)	0.0417 (9)	0.0066 (8)	-0.0103 (7)	-0.0007 (8)
C114	0.0479 (11)	0.0429 (10)	0.0596 (12)	0.0119 (8)	-0.0060 (9)	0.0010 (8)
C115	0.0609 (13)	0.0366 (9)	0.0669 (13)	0.0100 (9)	-0.0095 (10)	-0.0157 (9)
C116	0.0484 (10)	0.0384 (9)	0.0476 (10)	0.0022 (7)	-0.0113 (8)	-0.0150 (7)
C121	0.0332 (7)	0.0337 (7)	0.0232 (6)	-0.0008 (6)	-0.0048 (5)	-0.0050 (5)
C122	0.0572 (12)	0.0383 (9)	0.0645 (12)	-0.0062 (8)	-0.0304 (10)	0.0048 (8)
C123	0.0739 (15)	0.0438 (11)	0.0865 (17)	0.0045 (10)	-0.0421 (13)	0.0066 (11)
C124	0.0432 (10)	0.0688 (14)	0.0531 (11)	0.0107 (9)	-0.0203 (9)	-0.0118 (10)
C125	0.0369 (9)	0.0748 (14)	0.0462 (10)	-0.0138 (9)	-0.0103 (8)	-0.0100 (9)
C126	0.0444 (9)	0.0461 (10)	0.0392 (9)	-0.0117 (8)	-0.0092 (7)	-0.0016 (7)
C211	0.0302 (7)	0.0381 (8)	0.0268 (7)	-0.0072 (6)	0.0002 (5)	-0.0065 (6)
C212	0.0333 (8)	0.0508 (10)	0.0278 (7)	-0.0039 (7)	-0.0001 (6)	-0.0014 (6)
C213	0.0337 (8)	0.0670 (12)	0.0339 (8)	-0.0023 (8)	0.0031 (6)	-0.0119 (8)
C214	0.0435 (10)	0.0516 (11)	0.0573 (11)	-0.0056 (8)	0.0050 (8)	-0.0246 (9)
C215	0.0536 (11)	0.0357 (9)	0.0656 (12)	-0.0125 (8)	0.0071 (9)	-0.0110 (8)
C216	0.0468 (10)	0.0374 (9)	0.0424 (9)	-0.0138 (7)	0.0093 (7)	-0.0062 (7)
C221	0.0493 (9)	0.0329 (8)	0.0263 (7)	-0.0035 (7)	0.0128 (6)	0.0001 (6)
C222	0.0509 (11)	0.0668 (13)	0.0429 (10)	0.0005 (10)	0.0082 (8)	0.0024 (9)
C223	0.0549 (13)	0.0874 (18)	0.0668 (15)	0.0166 (12)	0.0210 (11)	0.0144 (13)
C224	0.095 (2)	0.0621 (14)	0.0505 (13)	0.0167 (13)	0.0379 (13)	0.0018 (10)
C225	0.0972 (18)	0.0469 (11)	0.0376 (10)	-0.0019 (11)	0.0259 (11)	-0.0102 (8)
C226	0.0682 (12)	0.0375 (9)	0.0301 (8)	-0.0107 (8)	0.0154 (8)	-0.0065 (6)
C16A	0.0383 (16)	0.055 (2)	0.0609 (17)	0.0003 (14)	0.0033 (12)	0.0128 (14)
C17A	0.0478 (17)	0.072 (2)	0.073 (2)	-0.0154 (14)	0.0098 (14)	-0.0167 (16)
C16B	0.083 (8)	0.059 (7)	0.083 (7)	-0.043 (6)	-0.025 (6)	0.007 (6)
C17B	0.047 (7)	0.152 (15)	0.098 (10)	0.022 (8)	-0.013 (6)	0.011 (9)

Geometric parameters (Å, °)

Co1—O11	2.0125 (10)	C112—H112	0.95
Co1—O11 ⁱ	2.0125 (10)	C113—C114	1.378 (3)
Co1—O12	2.0721 (10)	C113—H113	0.95
Co1—O12 ⁱ	2.0721 (10)	C114—C115	1.372 (3)
Co1—O13	2.1460 (12)	C114—H114	0.95
Co1—O13 ⁱ	2.1461 (12)	C115—C116	1.387 (3)
Co2—O21	2.0047 (11)	C115—H115	0.95
Co2—O21 ⁱⁱ	2.0047 (11)	C116—H116	0.95
Co2—O22	2.0786 (10)	C121—C122	1.382 (2)
Co2—O22 ⁱⁱ	2.0786 (10)	C121—C126	1.384 (2)
Co2—O23	2.1245 (12)	C122—C123	1.389 (3)
Co2—O23 ⁱⁱ	2.1245 (12)	C122—H122	0.95
O11—C11	1.2622 (16)	C123—C124	1.363 (3)
O12—C15	1.2550 (16)	C123—H123	0.95
O13—C16B	1.276 (11)	C124—C125	1.368 (3)

O13—C16A	1.465 (3)	C124—H124	0.95
O13—H13A	0.7956	C125—C126	1.394 (3)
O21—C21	1.2624 (17)	C125—H125	0.95
O22—C25	1.2593 (18)	C126—H12A	0.95
O23—C26	1.431 (3)	C211—C216	1.388 (2)
O23—H23A	0.8858	C211—C212	1.395 (2)
N11—C11	1.3746 (17)	C212—C213	1.386 (2)
N11—N12	1.3974 (16)	C212—H212	0.95
N11—C111	1.4189 (18)	C213—C214	1.376 (3)
N12—C13	1.313 (2)	C213—H213	0.95
N21—C21	1.370 (2)	C214—C215	1.380 (3)
N21—N22	1.3976 (16)	C214—H214	0.95
N21—C211	1.4165 (19)	C215—C216	1.385 (2)
N22—C23	1.313 (2)	C215—H215	0.95
C11—C12	1.4252 (19)	C216—H216	0.95
C12—C15	1.422 (2)	C221—C222	1.378 (3)
C12—C13	1.4336 (18)	C221—C226	1.394 (2)
C13—C14	1.496 (2)	C222—C223	1.397 (3)
C14—H14A	0.98	C222—H222	0.95
C14—H14B	0.98	C223—C224	1.380 (4)
C14—H14C	0.98	C223—H223	0.95
C15—C121	1.4934 (19)	C224—C225	1.363 (4)
C21—C22	1.434 (2)	C224—H224	0.95
C22—C25	1.417 (2)	C225—C226	1.385 (3)
C22—C23	1.438 (2)	C225—H225	0.95
C23—C24	1.500 (2)	C226—H226	0.95
C24—H24A	0.98	C16A—C17A	1.491 (5)
C24—H24B	0.98	C16A—H16A	0.99
C24—H24C	0.98	C16A—H16B	0.99
C25—C221	1.499 (2)	C17A—H17A	0.98
C26—C27	1.411 (5)	C17A—H17B	0.98
C26—H26A	0.99	C17A—H17C	0.98
C26—H26B	0.99	C16B—C17B	1.47 (2)
C27—H27A	0.98	C16B—H16C	0.99
C27—H27B	0.98	C16B—H16D	0.99
C27—H27C	0.98	C17B—H17D	0.98
C111—C116	1.387 (2)	C17B—H17E	0.98
C111—C112	1.390 (2)	C17B—H17F	0.98
C112—C113	1.386 (2)		
O11—Co1—O11 ⁱ	180.0	H27A—C27—H27B	109.5
O11—Co1—O12	90.12 (4)	C26—C27—H27C	109.5
O11 ⁱ —Co1—O12	89.88 (4)	H27A—C27—H27C	109.5
O11—Co1—O12 ⁱ	89.88 (4)	H27B—C27—H27C	109.5
O11 ⁱ —Co1—O12 ⁱ	90.12 (4)	C116—C111—C112	119.39 (15)
O12—Co1—O12 ⁱ	180.0	C116—C111—N11	119.69 (14)
O11—Co1—O13	90.93 (5)	C112—C111—N11	120.91 (13)
O11 ⁱ —Co1—O13	89.07 (5)	C113—C112—C111	119.63 (15)

O12—Co1—O13	89.14 (5)	C113—C112—H112	120.2
O12 ⁱ —Co1—O13	90.86 (5)	C111—C112—H112	120.2
O11—Co1—O13 ⁱ	89.07 (5)	C114—C113—C112	121.06 (17)
O11 ⁱ —Co1—O13 ⁱ	90.93 (5)	C114—C113—H113	119.5
O12—Co1—O13 ⁱ	90.86 (5)	C112—C113—H113	119.5
O12 ⁱ —Co1—O13 ⁱ	89.14 (5)	C115—C114—C113	119.04 (17)
O13—Co1—O13 ⁱ	180.0	C115—C114—H114	120.5
O21—Co2—O21 ⁱⁱ	180.0	C113—C114—H114	120.5
O21—Co2—O22	88.73 (4)	C114—C115—C116	121.02 (18)
O21 ⁱⁱ —Co2—O22	91.27 (4)	C114—C115—H115	119.5
O21—Co2—O22 ⁱⁱ	91.27 (4)	C116—C115—H115	119.5
O21 ⁱⁱ —Co2—O22 ⁱⁱ	88.73 (4)	C115—C116—C111	119.84 (17)
O22—Co2—O22 ⁱⁱ	179.9980 (10)	C115—C116—H116	120.1
O21—Co2—O23	89.54 (5)	C111—C116—H116	120.1
O21 ⁱⁱ —Co2—O23	90.46 (5)	C122—C121—C126	119.27 (15)
O22—Co2—O23	92.76 (4)	C122—C121—C15	119.56 (14)
O22 ⁱⁱ —Co2—O23	87.24 (4)	C126—C121—C15	120.97 (14)
O21—Co2—O23 ⁱⁱ	90.46 (5)	C121—C122—C123	120.02 (19)
O21 ⁱⁱ —Co2—O23 ⁱⁱ	89.54 (5)	C121—C122—H122	120.0
O22—Co2—O23 ⁱⁱ	87.24 (4)	C123—C122—H122	120.0
O22 ⁱⁱ —Co2—O23 ⁱⁱ	92.76 (4)	C124—C123—C122	120.4 (2)
O23—Co2—O23 ⁱⁱ	180.0	C124—C123—H123	119.8
C11—O11—Co1	121.85 (9)	C122—C123—H123	119.8
C15—O12—Co1	128.43 (9)	C123—C124—C125	120.19 (18)
C16B—O13—Co1	139.3 (5)	C123—C124—H124	119.9
C16A—O13—Co1	121.07 (15)	C125—C124—H124	119.9
C16B—O13—H13A	104.2	C124—C125—C126	120.19 (18)
C16A—O13—H13A	115.2	C124—C125—H125	119.9
Co1—O13—H13A	116.1	C126—C125—H125	119.9
C21—O21—Co2	122.73 (10)	C121—C126—C125	119.86 (18)
C25—O22—Co2	128.58 (10)	C121—C126—H12A	120.1
C26—O23—Co2	122.93 (12)	C125—C126—H12A	120.1
C26—O23—H23A	106.9	C216—C211—C212	119.73 (15)
Co2—O23—H23A	121.5	C216—C211—N21	119.95 (14)
C11—N11—N12	111.13 (11)	C212—C211—N21	120.32 (14)
C11—N11—C111	128.62 (11)	C213—C212—C211	119.18 (17)
N12—N11—C111	120.16 (11)	C213—C212—H212	120.4
C13—N12—N11	106.47 (11)	C211—C212—H212	120.4
C21—N21—N22	111.42 (12)	C214—C213—C212	121.13 (17)
C21—N21—C211	127.65 (12)	C214—C213—H213	119.4
N22—N21—C211	120.37 (12)	C212—C213—H213	119.4
C23—N22—N21	106.32 (12)	C213—C214—C215	119.52 (17)
O11—C11—N11	122.94 (12)	C213—C214—H214	120.2
O11—C11—C12	131.05 (13)	C215—C214—H214	120.2
N11—C11—C12	105.98 (11)	C214—C215—C216	120.40 (19)
C15—C12—C11	122.89 (12)	C214—C215—H215	119.8
C15—C12—C13	132.25 (13)	C216—C215—H215	119.8
C11—C12—C13	104.83 (12)	C215—C216—C211	120.02 (16)

N12—C13—C12	111.57 (12)	C215—C216—H216	120.0
N12—C13—C14	118.60 (13)	C211—C216—H216	120.0
C12—C13—C14	129.73 (14)	C222—C221—C226	119.56 (17)
C13—C14—H14A	109.5	C222—C221—C25	118.70 (16)
C13—C14—H14B	109.5	C226—C221—C25	121.67 (16)
H14A—C14—H14B	109.5	C221—C222—C223	119.9 (2)
C13—C14—H14C	109.5	C221—C222—H222	120.0
H14A—C14—H14C	109.5	C223—C222—H222	120.0
H14B—C14—H14C	109.5	C224—C223—C222	119.5 (2)
O12—C15—C12	122.74 (12)	C224—C223—H223	120.2
O12—C15—C121	115.73 (13)	C222—C223—H223	120.2
C12—C15—C121	121.50 (12)	C225—C224—C223	120.8 (2)
O21—C21—N21	122.37 (13)	C225—C224—H224	119.6
O21—C21—C22	131.49 (15)	C223—C224—H224	119.6
N21—C21—C22	106.11 (12)	C224—C225—C226	120.0 (2)
C25—C22—C21	122.18 (13)	C224—C225—H225	120.0
C25—C22—C23	133.13 (14)	C226—C225—H225	120.0
C21—C22—C23	104.25 (13)	C225—C226—C221	120.1 (2)
N22—C23—C22	111.88 (13)	C225—C226—H226	120.0
N22—C23—C24	117.70 (14)	C221—C226—H226	120.0
C22—C23—C24	130.42 (15)	O13—C16A—C17A	112.7 (3)
C23—C24—H24A	109.5	O13—C16A—H16A	109.1
C23—C24—H24B	109.5	C17A—C16A—H16A	109.1
H24A—C24—H24B	109.5	O13—C16A—H16B	109.1
C23—C24—H24C	109.5	C17A—C16A—H16B	109.1
H24A—C24—H24C	109.5	H16A—C16A—H16B	107.8
H24B—C24—H24C	109.5	O13—C16B—C17B	112.6 (11)
O22—C25—C22	122.65 (13)	O13—C16B—H16C	109.1
O22—C25—C221	115.15 (14)	C17B—C16B—H16C	109.1
C22—C25—C221	122.12 (13)	O13—C16B—H16D	109.1
C27—C26—O23	112.4 (3)	C17B—C16B—H16D	109.1
C27—C26—H26A	109.1	H16C—C16B—H16D	107.8
O23—C26—H26A	109.1	C16B—C17B—H17D	109.5
C27—C26—H26B	109.1	C16B—C17B—H17E	109.5
O23—C26—H26B	109.1	H17D—C17B—H17E	109.5
H26A—C26—H26B	107.9	C16B—C17B—H17F	109.5
C26—C27—H27A	109.5	H17D—C17B—H17F	109.5
C26—C27—H27B	109.5	H17E—C17B—H17F	109.5
O12—Co1—O11—C11	-14.75 (12)	N21—N22—C23—C24	-178.81 (16)
O12 ⁱ —Co1—O11—C11	165.25 (12)	C25—C22—C23—N22	172.28 (17)
O13—Co1—O11—C11	-103.89 (12)	C21—C22—C23—N22	0.03 (19)
O13 ⁱ —Co1—O11—C11	76.11 (12)	C25—C22—C23—C24	-8.4 (3)
O11—Co1—O12—C15	2.27 (13)	C21—C22—C23—C24	179.35 (19)
O11 ⁱ —Co1—O12—C15	-177.73 (13)	Co2—O22—C25—C22	-12.6 (2)
O13—Co1—O12—C15	93.20 (13)	Co2—O22—C25—C221	170.58 (11)
O13 ⁱ —Co1—O12—C15	-86.80 (13)	C21—C22—C25—O22	-3.2 (2)
O11—Co1—O13—C16B	90.8 (10)	C23—C22—C25—O22	-174.33 (17)

O11 ⁱ —Co1—O13—C16B	-89.2 (10)	C21—C22—C25—C221	173.43 (15)
O12—Co1—O13—C16B	0.7 (10)	C23—C22—C25—C221	2.3 (3)
O12 ⁱ —Co1—O13—C16B	-179.3 (10)	Co2—O23—C26—C27	108.5 (3)
O11—Co1—O13—C16A	131.2 (2)	C11—N11—C111—C116	-160.85 (16)
O11 ⁱ —Co1—O13—C16A	-48.8 (2)	N12—N11—C111—C116	15.4 (2)
O12—Co1—O13—C16A	41.1 (2)	C11—N11—C111—C112	17.8 (2)
O12 ⁱ —Co1—O13—C16A	-138.9 (2)	N12—N11—C111—C112	-165.92 (14)
O22—Co2—O21—C21	-17.43 (13)	C116—C111—C112—C113	-0.1 (3)
O22 ⁱⁱ —Co2—O21—C21	162.58 (13)	N11—C111—C112—C113	-178.76 (16)
O23—Co2—O21—C21	-110.20 (13)	C111—C112—C113—C114	-1.2 (3)
O23 ⁱⁱ —Co2—O21—C21	69.80 (13)	C112—C113—C114—C115	1.7 (3)
O21—Co2—O22—C25	19.90 (13)	C113—C114—C115—C116	-0.9 (4)
O21 ⁱⁱ —Co2—O22—C25	-160.10 (13)	C114—C115—C116—C111	-0.3 (3)
O23—Co2—O22—C25	109.38 (13)	C112—C111—C116—C115	0.8 (3)
O23 ⁱⁱ —Co2—O22—C25	-70.62 (13)	N11—C111—C116—C115	179.53 (18)
O21—Co2—O23—C26	62.54 (19)	O12—C15—C121—C122	-49.2 (2)
O21 ⁱⁱ —Co2—O23—C26	-117.46 (19)	C12—C15—C121—C122	132.44 (18)
O22—Co2—O23—C26	-26.17 (19)	O12—C15—C121—C126	125.62 (16)
O22 ⁱⁱ —Co2—O23—C26	153.83 (19)	C12—C15—C121—C126	-52.7 (2)
C11—N11—N12—C13	0.52 (17)	C126—C121—C122—C123	1.9 (3)
C111—N11—N12—C13	-176.35 (14)	C15—C121—C122—C123	176.9 (2)
C21—N21—N22—C23	-1.08 (18)	C121—C122—C123—C124	-1.6 (4)
C211—N21—N22—C23	-173.12 (14)	C122—C123—C124—C125	-0.3 (4)
Co1—O11—C11—N11	-164.27 (10)	C123—C124—C125—C126	1.9 (3)
Co1—O11—C11—C12	17.8 (2)	C122—C121—C126—C125	-0.3 (3)
N12—N11—C11—O11	-179.12 (13)	C15—C121—C126—C125	-175.20 (16)
C111—N11—C11—O11	-2.6 (2)	C124—C125—C126—C121	-1.6 (3)
N12—N11—C11—C12	-0.73 (16)	C21—N21—C211—C216	-150.36 (17)
C111—N11—C11—C12	175.81 (14)	N22—N21—C211—C216	20.3 (2)
O11—C11—C12—C15	-2.9 (3)	C21—N21—C211—C212	29.1 (2)
N11—C11—C12—C15	178.94 (13)	N22—N21—C211—C212	-160.23 (15)
O11—C11—C12—C13	178.84 (16)	C216—C211—C212—C213	0.9 (2)
N11—C11—C12—C13	0.64 (16)	N21—C211—C212—C213	-178.58 (15)
N11—N12—C13—C12	-0.09 (18)	C211—C212—C213—C214	-1.5 (3)
N11—N12—C13—C14	-176.89 (16)	C212—C213—C214—C215	0.7 (3)
C15—C12—C13—N12	-178.42 (16)	C213—C214—C215—C216	0.7 (3)
C11—C12—C13—N12	-0.35 (18)	C214—C215—C216—C211	-1.2 (3)
C15—C12—C13—C14	-2.1 (3)	C212—C211—C216—C215	0.4 (3)
C11—C12—C13—C14	176.00 (19)	N21—C211—C216—C215	179.93 (16)
Co1—O12—C15—C12	9.8 (2)	O22—C25—C221—C222	54.7 (2)
Co1—O12—C15—C121	-168.49 (10)	C22—C25—C221—C222	-122.18 (19)
C11—C12—C15—O12	-12.5 (2)	O22—C25—C221—C226	-122.06 (17)
C13—C12—C15—O12	165.27 (16)	C22—C25—C221—C226	61.1 (2)
C11—C12—C15—C121	165.69 (14)	C226—C221—C222—C223	-2.6 (3)
C13—C12—C15—C121	-16.5 (3)	C25—C221—C222—C223	-179.47 (19)
Co2—O21—C21—N21	-171.43 (11)	C221—C222—C223—C224	2.2 (4)
Co2—O21—C21—C22	10.5 (3)	C222—C223—C224—C225	-0.2 (4)
N22—N21—C21—O21	-177.42 (15)	C223—C224—C225—C226	-1.3 (4)

C211—N21—C21—O21	-6.1 (3)	C224—C225—C226—C221	0.8 (3)
N22—N21—C21—C22	1.09 (18)	C222—C221—C226—C225	1.2 (3)
C211—N21—C21—C22	172.42 (15)	C25—C221—C226—C225	177.89 (17)
O21—C21—C22—C25	4.3 (3)	C16B—O13—C16A—C17A	-27.7 (10)
N21—C21—C22—C25	-173.99 (14)	Co1—O13—C16A—C17A	-160.64 (19)
O21—C21—C22—C23	177.64 (17)	C16A—O13—C16B—C17B	-1.5 (7)
N21—C21—C22—C23	-0.67 (17)	Co1—O13—C16B—C17B	72.6 (14)
N21—N22—C23—C22	0.61 (19)		

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+2, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O13—H13 <i>A</i> \cdots N22 ⁱⁱⁱ	0.80	2.04	2.8314 (18)	175
O23—H23 <i>A</i> \cdots N12 ^{iv}	0.89	1.90	2.7862 (16)	177
C16 <i>A</i> —H16 <i>B</i> \cdots O12	0.99	2.56	3.192 (3)	122
C24—H24 <i>C</i> \cdots O11 ^v	0.98	2.46	3.361 (2)	154
C112—H112 \cdots O11	0.95	2.24	2.8478 (19)	121
C116—H116 \cdots O23 ^{vi}	0.95	2.60	3.458 (2)	151
C116—H116 \cdots N12	0.95	2.50	2.827 (2)	100
C212—H212 \cdots O21	0.95	2.31	2.848 (2)	115

Symmetry codes: (iii) $x-1, y, z$; (iv) $x+1, y, z-1$; (v) $x+1, y, z$; (vi) $x-1, y, z+1$.